arXiv:math/0304411v1 [math.PR] 25 Apr 2003

ADDITIVE FUNCTIONALS ON RANDOM SEARCH TREES

by

Nevin Kapur

A dissertation submitted to The Johns Hopkins University in conformity with the
requirements for the degree of Doctor of Philosophy

Baltimore, Maryland
May 2003

© Nevin Kapur 2003
All rights reserved

Abstract

Search trees are fundamental data structures in computer science. We study functionals on random search trees that satisfy recurrence relations of a simple additive form. Many important functionals including the space requirement, internal path length, and the so-called shape functional fall under this framework. Our goal is to derive asymptotics of moments and identify limiting distributions of these functionals under two commonly studied probability models—the random permutation model and the uniform model.

For the random permutation model, our approach is based on establishing *transfer theorems* that link the order of growth of the input into a particular (deterministic) recurrence to the order of growth of the output. For the uniform model, our approach is based on the complex-analytic tool of *singularity analysis*. To facilitate a systematic analysis of these additive functionals we extend singularity analysis, a class of methods by which one can translate on a term-by-term basis an asymptotic expansion of a functional around its dominant singularity into a corresponding expansion for the Taylor coefficients of the function. The most important extension is the determination of how singularities are composed under the operation of Hadamard product of analytic power series.

The transfer theorems derived are used in conjunction with the method of moments to establish limit laws for m-ary search trees under the random permutation model. For the uniform model on binary search trees, the extended singularity analysis toolkit is employed to establish the asymptotic behavior of the moments of a wide class of functionals. These asymptotics are used, again in conjunction with the method of moments, to derive limit laws.

Advisor: James Allen Fill
Readers: James Allen Fill and John C. Wierman

Acknowledgments

It is impossible for me to express satisfactorily in words the gratitude I owe my advisor, Jim Fill. This dissertation is the result of many, many sessions of joint toil where I learned the art of research from him. For that and much more, I am indebted to him.

This work would have been nothing more than a graduate student's distant dream were it not for the contribution of Philippe Flajolet. His outline concerning singularity analysis for Hadamard products blossomed into this dissertation's central Chapter 4, and his vision throughout the project was invaluable.

I thank Professors Goldman, Priebe, Scheinerman, and Wierman for serving on my dissertation defense committee and offering suggestions to improve this work.

I am thankful to the faculty, staff, and students of the Department of Mathematical Sciences at The Johns Hopkins University for their support.

My stay in Baltimore has been most enjoyable thanks to the excellent friends I found here. I will always have fond memories of the times I spent with Carolyn Cooper, helen and Jason DeVinney, and EJ Valitutto.

My mother taught me to value education and my parents sacrificed much so that I could pursue my dream. To them I owe much.

Nevin Kapur

Contents

Abstract ii

Acknowledgments iii

Contents iv

List of Figures vi

I Preliminaries 1

1 Introduction 2
- 1.1 Background . . . 2
- 1.2 Probability models . . . 4
- 1.3 Roadmap . . . 4

2 Additive tree functionals 6
- 2.1 Notation and definitions . . . 6
- 2.2 Examples . . . 7
- 2.3 A common framework for all moments . . . 7

II Theory 11

3 Random permutation model: transfer theorems 12
- 3.1 The Exact Transfer Theorem . . . 13
- 3.2 The Asymptotic Transfer Theorem . . . 13

4 Singularity analysis of Hadamard products 19
- 4.1 Background . . . 19
- 4.2 Singular expansions and differentiation . . . 21
- 4.3 Hadamard products . . . 25

III Applications 34

5 Random permutation model: shape functional 35
- 5.1 Complete asymptotic expansion of the mean . . . 35
- 5.2 The variance for binary search trees . . . 45

6 Random permutation model: limiting distributions 53
- 6.1 Small toll functions . . . 53
- 6.2 Moderate and large toll functions . . . 60

7 Catalan model: the toll function n^α 66
7.1 Preliminaries 66
7.2 Asymptotics of the mean 68
7.3 Higher moments 70
7.4 Asymptotics of moments 78
7.5 The limiting distributions 79
7.5.1 $\alpha \neq 1/2$ 79
7.5.2 $\alpha = 1/2$ 84
7.5.3 A unified result 85

8 Catalan model: the shape functional 88
8.1 Mean 88
8.2 Second moment and variance 88
8.3 Higher moments 90
8.4 A central limit theorem 92
8.5 Sufficient conditions for asymptotic normality 93

IV Appendices and Bibliography 94

A Solution of an Euler differential equation 95

B Identities involving the indicial polynomial 101

Bibliography 107

Vita 112

List of Figures

1.1.1 A quaternary (i.e., "4-ary") tree. 3
1.1.2 Sequential construction of a quaternary search tree. 3
1.2.1 The distinct ternary trees and the permutations that generate them, for $n = 4$. . 5

2.1.1 Example of notation for m-ary search trees. 6

4.1.1 The region $\Delta(\phi, \eta)$ in Definition 4.1.1. 20
4.2.1 The contour γ used in the proof of the differentiation theorem. 22
4.2.2 The contour used in the proof of the integration theorem. 24
4.3.1 The contour γ_0. 27
4.3.2 The contour γ. 28
4.3.3 The geometry of the contour γ. 28

7.5.1 σ^2 of (7.5.3) as a function of α. 82
7.5.2 $\mathbf{E}\,[Y - \mathbf{E}\,Y]^3$ of Proposition 7.5.2 as a function of α. 83
7.5.3 f_Y of Remark 7.5.4. 84

“The time has come,” the Walrus said,
“To talk of many things: . . . ”

The Walrus and The Carpenter
LEWIS CARROLL

Part I

Preliminaries

Chapter 1

Introduction

1.1 Background

We start by giving a brief overview of search trees, the basic object under scrutiny in this dissertation. Search trees are fundamental data structures in computer science, their primary applications being in searching and sorting. For integer $m \geq 2$, the m-ary search tree, or multiway tree, generalizes the binary search tree. The quantity m is called the *branching factor*. According to [49], search trees of branching factors higher than 2 were first suggested by Muntz and Uzgalis [56] "to solve internal memory problems with large quantities of data." For more background we refer the reader to [43, 44] and [49].

An *m-ary tree* is a rooted tree with at most m "children" for each *node (vertex)*, each child of a node being distinguished as one of m possible types. Recursively expressed, an m-ary tree either is empty or consists of a distinguished node (called the *root*) together with an ordered m-tuple of *subtrees*, each of which is an m-ary tree.

An *m-ary search tree* is an m-ary tree in which each node has the capacity to contain $m-1$ elements of some linearly ordered set, called the set of *keys*. In typical implementations of m-ary search trees, the keys at each node are stored in increasing order and at each node one has m pointers to the subtrees. By spreading the input data in m directions instead of only 2, as is the case for a binary search tree, one seeks to have shorter path lengths and thus quicker searches.

We consider the space of m-ary search trees on n keys, and assume that the keys are linearly ordered. Hence, without loss of generality, we can take the set of keys to be $[n] := \{1, 2, \dots, n\}$. We construct an m-ary search tree from a sequence s of n distinct keys in the following way:

1. If $n < m$, then all the keys are stored in the root node in increasing order.

2. If $n \geq m$, then the first $m-1$ keys in the sequence are stored in the root in increasing order, and the remaining $n-(m-1)$ keys are stored in the subtrees subject to the condition that if $\sigma_1 < \sigma_2 < \cdots < \sigma_{m-1}$ denotes the ordered sequence of keys in the root, then the keys in the jth subtree are those that lie between σ_{j-1} and σ_j, where $\sigma_0 := 0$ and $\sigma_m := n+1$, sequenced as in s.

3. All the subtrees are m-ary search trees that satisfy conditions 1, 2, and 3.

Figure 1.1.1 shows the quaternary tree associated with the sequence

$$s := (10, 7, 12, 4, 1, 8, 5, 6, 9, 14, 11, 2, 15, 13, 3).$$

We may also build an m-ary search tree by inserting the successive elements of a sequence into an initially empty tree. This process is illustrated for the sequence s in Figure 1.1.2.

It is our goal to study functionals defined on m-ary search trees under two natural probability models. The functionals we consider are of additive type, each induced by a "toll function." Informally speaking, the value of an additive functional on an m-ary search tree is recursively defined as the sum of the functional's value on the subtrees rooted at the children of the root of the tree and a term (the *toll*) that is a function of the size of the tree. Additive functionals



Figure 1.1.1: The quaternary (i.e., “4-ary”) tree associated with the sequence s.



Figure 1.1.2: Sequential construction of the quaternary search tree from the sequence s.

defined in this manner represent the cost of divide-and-conquer algorithms, where the inherent recursive nature of the algorithms lends itself naturally to such a formulation.

1.2 Probability models

The two most common probability models on the space of m-ary search trees are the *uniform* model and the *random permutation*, or *random insertion*, model. In the case $m = 2$, the number of search trees on n keys is enumerated by the nth Catalan number. Thus the uniform model on binary search trees is also referred to as the *Catalan* model.

Under the uniform model, the probability of obtaining any particular tree is just the reciprocal of the number of m-ary search trees on n keys. Fill and Dobrow [22] have studied problems associated with determining this number. Fill [21] argued that the functional corresponding to the toll sequence $(\ln n)_{n\geq 1}$ serves as a crude measure of the "shape" of a binary search tree, and explained how this functional arises in connection with the move-to-root self-organizing scheme for dynamic maintenance of binary search trees. He obtained asymptotic information about the mean and variance under the Catalan model. (The latter results were rederived in the extension [55] from binary search trees to simply generated rooted trees.)

In [32, Prop. 2] Flajolet and Steyaert gave order-of-growth information about the mean of functionals induced by tolls of the form n^{α}. Takács established the limiting (Airy) distribution of path length in Catalan trees [70–72], which is the additive functional for the toll $n - 1$. The additive functional for the toll n^2 arises in the study of the Wiener index of the tree and has been analyzed by Janson [41].

The uniform model on binary search trees has been also been used recently by Janson [40] in the analysis of an algorithm of Koda and Ruskey [47] for listing ideals in a forest poset.

Now we turn our attention to the random permutation model. Let π be a random permutation of $[n]$: each of the $n!$ permutations is equally likely to be the value of π. The distribution of trees under the random permutation model is the distribution induced by constructing an m-ary search tree from π in the manner we have described in Section 1.1. For example, for $m = 3$ and $n = 4$, there are 6 distinct trees each occurring with probability 1/6, as depicted in Figure 1.2.1. We note that it is not the case in general that all trees are equiprobable. For example, if $m = 2$ and $n = 3$, there are 5 distinct trees; 4 of these have probability 1/6 each, and 1 (corresponding to permutations 213 and 231) has probability 2/6.

Unlike under the uniform model, functionals of search trees under the random permutation model have been studied a great deal, the most attention being paid to binary search trees. Indeed, much of the motivation for studying additive functionals comes from the fact that the number of key comparisons in Quicksort is such a functional and under the random permutation model it corresponds to the cost of Quicksort on uniformly random input. There is an extensive literature on the analysis of Quicksort; see, for example, [2, 10, 24–27, 33, 35, 36, 38, 42, 60, 63, 67, 73]. Many researchers have studied various other functionals under the random permutation model; see [7, 11–16, 18, 19, 21, 29, 39, 53, 57–59, 61, 62, 66] for treatment of functionals defined on binary search trees. For m-ary search trees see [8, 22, 48, 50]. Chapter 3 of [49] is a rich source of results on the properties of m-ary search trees.

1.3 Roadmap

In Chapter 2 we will formally define the problem of additive tree functionals, introducing the concept of a toll function. The recurrences obtained will be translated to relations among generating functions. The resulting relations will serve as a motivation for our main theoretical results—*transfer theorems* and the extension of the *singularity analysis* toolkit. In Chapter 3 we consider the random permutation model and prove general results that relate the asymptotics of the toll function to that of the corresponding tree functional. We will then proceed to extend



$1234, 1243, 2134, 2143$

$1324, 1342, 3124, 3142$

$1423, 1432, 4123, 4132$

$2314, 2341, 3214, 3241$

$2413, 2431, 4213, 4231$

$3412, 3421, 4312, 4321$

Figure 1.2.1: The distinct ternary trees and the permutations that generate them, for $n = 4$.

singularity analysis in Chapter 4 so that it is readily applicable to recurrences obtained under the uniform model.

Next we will consider various applications. In Chapter 5 we will combine the use of the extended singularity analysis toolkit and the transfer theorems derived in Chapter 3 to derive a complete asymptotic expansion of the mean of the shape functional of an m-ary search tree under the random permutation model and the variance for binary search trees. Then in Chapter 6 we will use the transfer theorems of Chapter 3 to determine limiting distributions of additive tree functionals under the random permutation model. Finally, in Chapters 7 and 8, the extension of singularity analysis to Hadamard products will be applied to the Catalan model for binary search trees, resulting in limit laws under two different classes of toll functions.

Chapter 2

Additive tree functionals

We now formally introduce additive functionals on m-ary search trees. We will provide some motivating examples and derive the basic recurrence that governs the behavior of these functions under each of the random permutation model and the uniform model.

2.1 Notation and definitions

We first establish some notation. Let T be an m-ary search tree. We use $|T|$ to denote the number of keys in T. Call a node *full* if it contains $m - 1$ keys. For $1 \leq j \leq m$, let $L_j(T)$ denote the jth subtree pendant from the root of T. [The numbering of $L_1(T), \ldots, L_m(T)$ will correspond to left-to-right ordering in our figures.] When it is clear to which tree T we are referring we will denote the subtrees simply by $L_1, \ldots, L_m$. For x a node in T, write T_x for the subtree of T consisting of x and its descendants, with x as root. This notation is illustrated in Figure 2.1.1.



Figure 2.1.1: Example of notation for the quaternary tree T of Figures 1.1.1 and 1.1.2. In this example, $|T| = 15$ and $T_x = L_1$.

Definition 2.1.1. Fix $m \geq 2$. We will call a functional f on m-ary search trees an *additive tree functional* if it satisfies the recurrence

$$f(T) = \sum_{i=1}^{m} f(L_i(T)) + t_{|T|}, \tag{2.1.1}$$

for any tree T with $|T| \geq m - 1$. Here $(t_n)_{n \geq m-1}$ is a given sequence, henceforth called the *toll function.*

Note that the recurrence (2.1.1) does not make any reference to t_n for $0 \leq n \leq m - 2$ nor specify $f(T)$ for $0 \leq |T| \leq m - 2$.

2.2 Examples

Several interesting examples can be cast as additive functionals.

Example 2.2.1. If we specify $f(T)$ arbitrarily for $0 \le |T| \le m-2$ and take $t_n \equiv c$ for $n \ge m-1$, we obtain the "additive functional" framework of [49, §3.1]. (Our definition of an additive functional generalizes this notion.) In particular if we define $f(\emptyset) := 0$ and $f(T) := 1$ for the unique m-ary search tree T on n keys for $1 \le n \le m-2$ and let $t_n \equiv 1$ for $n \ge m-1$, then $f(T)$ counts the number of nodes in T and thus gives the *space requirement* functional discussed in [49, §3.4].

Example 2.2.2. If we define $f(T) := 0$ when $0 \le |T| \le m-2$ and $t_n := n-(m-1)$ for $n \ge m-1$ then f is the *internal path length* functional discussed in [49, §3.5]: $f(T)$ is the sum of all root-to-key distances in T.

Example 2.2.3. As noted in Chapter 1, we can associate with each permutation of $[n]$ an m-ary search tree. Let Q denote the probability mass function of the distribution of trees induced by associating an m-ary search tree with each of the $n!$ (equally likely) permutations of $[n]$. Dobrow and Fill [17] noted that

$$Q(T) = \frac{1}{\prod_x \binom{|T_x|}{m-1}}, \tag{2.2.1}$$

where the product in (2.2.1) is over all full nodes in T. This functional is sometimes called the *shape functional* as it serves as a crude measure of the "shape" of the tree, with "full" trees (like the complete tree) achieving larger values of Q. For further discussions along these lines, consult [17] and [21]. If we define $f(T) := 0$ for $0 \le |T| \le m-2$ and $t_n := \ln \binom{n}{m-1}$ for $n \ge m-1$, then $f(T) = -\ln Q(T)$. In this work we will refer to $-\ln Q$ (rather than Q) as the shape functional.

2.3 A common framework for all moments

Our aim is to study the distribution of the functional $f(T)$ satisfying (2.1.1) under the two aforementioned probability models. Under the uniform model T is distributed uniformly over all m-ary search trees with n keys, while under the random permutation model T has the distribution Q defined at (2.2.1). In either case we will use X_n to denote a random variable having the distribution of $f(T)$ and define $\mu_n(k) := \mathbf{E}\, X_n^k$ for $k \ge 1$, the choice of model being clear from context. To address the problem recursively we examine the joint distribution of the subtree sizes $|L_1|, \ldots, |L_m|$.

Random permutation model

Under the random permutation model the joint distribution of the subtree sizes $|L_1|, \ldots, |L_m|$ is uniform over all $\binom{n}{m-1}$ m-tuples of nonnegative integers that sum to $n-(m-1)$: see [49, Exercise 3.8]. We now apply the law of total expectation by conditioning on $(|L_1|, \ldots, |L_m|)$. Let $\sum_{\mathbf{j}}$ denote the sum over all m-tuples $(j_1, \ldots, j_m)$ that sum to $n-(m-1)$ and $\sum_{\mathbf{k}}$ the sum over all $(m+1)$-tuples $(k_1, \ldots, k_{m+1})$ of nonnegative integers that sum to k. Then, letting $\oplus$ denote sums of mutually independent random variables, for $n \ge m-1$ we have

$$\begin{aligned}\mu_n(k) = \mathbf{E}\,\mathbf{E}\,(X_n^k \bigm| |L_1|, \ldots, |L_m|) &= \frac{1}{\binom{n}{m-1}} \sum_{\mathbf{j}} \mathbf{E}\,(X_{j_1} \oplus \cdots \oplus X_{j_m} + t_n)^k \\ &= \frac{1}{\binom{n}{m-1}} \sum_{\mathbf{j}} \sum_{\mathbf{k}} \binom{k}{k_1, \ldots, k_m, k_{m+1}} \mu_{j_1}(k_1) \cdots \mu_{j_m}(k_m) t_n^{k_{m+1}}.\end{aligned}$$

We can rewrite this equation as

$$\mu_n(k) = \frac{m}{\binom{n}{m-1}} \sum_{j=0}^{n-(m-1)} \binom{n-1-j}{m-2} \mu_j(k) + r_n(k), \tag{2.3.1}$$

where

$$r_n(k) := \sum_{\mathbf{k}}{}^{*} \binom{k}{k_1, \ldots, k_m, k_{m+1}} t_n^{k_{m+1}} \frac{1}{\binom{n}{m-1}} \sum_{\mathbf{j}} \mu_{j_1}(k_1) \cdots \mu_{j_m}(k_m), \tag{2.3.2}$$

with $\sum_{\mathbf{k}}^{*}$ denoting the same sum as $\sum_{\mathbf{k}}$ with the additional restriction that $k_i < k$ for $i = 1, \ldots, m$. We have thus established that the moments $\mu_n(k)$ each satisfy the same basic recurrence in n, differing as k varies only in the non-homogeneous term $r_n(k)$. Observe that $r_n(1) = t_n$, the toll function. We record this important fact as

Proposition 2.3.1. *Under the random permutation model all moments of an additive functional satisfy the basic recurrence*

$$a_n = b_n + \frac{m}{\binom{n}{m-1}} \sum_{j=0}^{n-(m-1)} \binom{n-1-j}{m-2} a_j, \qquad n \geq m-1, \tag{2.3.3}$$

with specified initial conditions (say) $a_j := b_j$, $0 \leq j \leq m-2$. [Recall the statement following Definition 2.1.1 about the initial conditions for the recurrence (2.1.1)*.]*

In order to analyze the recurrence relation (2.3.3) we introduce generating functions

$$A(z) := \sum_{n=0}^{\infty} a_n z^n \quad \text{and} \quad B(z) := \sum_{n=0}^{\infty} b_n z^n.$$

Furthermore, let

$$x^{\overline{r}} := \prod_{k=0}^{r-1} (x+k)$$

denote the rth rising factorial power of x and

$$x^{\underline{r}} := \prod_{k=0}^{r-1} (x-k)$$

the rth falling factorial power of x. Multiplying (2.3.3) by $n^{\underline{m-1}} z^{n-(m-1)}$ and summing over $n \geq m-1$ we get the differential equation

$$A^{(m-1)}(z) = B^{(m-1)}(z) + m!(1-z)^{-(m-1)} A(z). \tag{2.3.4}$$

Equations of the form (2.3.4) are members of a class known as *Euler differential equations.* Using the method of variation of parameters and combinatorial identities one can obtain the general solution to this equation. We only state the results here, referring the interested reader to Appendix A.

Theorem 2.3.2. *Let A and B denote the respective ordinary generating functions of the se-*

quences (a_n) *and* (b_n) *in Proposition 2.3.1. Then*

$$A(z) = \sum_{j=1}^{m-1} c_j (1-z)^{-\lambda_j} + \sum_{j=1}^{m-1} \frac{(1-z)^{-\lambda_j}}{\psi'(\lambda_j)} \int_0^z B^{(m-1)}(\zeta)(1-\zeta)^{\lambda_j+m-2}\, d\zeta, \tag{2.3.5}$$

where ψ *is the indicial polynomial*

$$\psi(\lambda) := \lambda^{\overline{m-1}} - m! = \lambda(\lambda+1)\cdots(\lambda+m-2) - m! \tag{2.3.6}$$

with roots $2 =: \lambda_1, \lambda_2, \ldots, \lambda_{m-1}$ *in nonincreasing order of real parts. In* (2.3.5)*, the coefficients* $c_1, c_2, \ldots, c_{m-1}$ *can be written explicitly in terms of the initial conditions* $b_0, \ldots, b_{m-2}$ *as*

$$c_j = \frac{m!}{\psi'(\lambda_j)} \sum_{k=0}^{m-2} b_k \frac{k!}{\lambda_j^{\overline{k+1}}}, \qquad j = 1, \ldots, m-1. \tag{2.3.7}$$

Proof. See Appendix A, in particular Corollary A.3 and Proposition A.4. □

The indicial polynomial (2.3.6) is well studied; see [8, 49, 51] and Appendix B. We will exploit the expression (2.3.5) for $A(z)$ to relate the asymptotic properties of the sequence (b_n) to those of (a_n) in Chapter 3 and then, in Chapter 6, use the transfer theorems of Chapter 3 to derive limiting distributions of additive functionals.

Uniform model for binary trees

We now turn our attention to the uniform model for binary search trees. The techniques we develop in this dissertation can be extended to handle the uniform model on m-ary search trees as well, although much of that analysis will involve singularity analysis of implicitly defined functions. Our techniques can also be extended to handle any simply generated family of trees [54, 69].

It is well known that the number of binary trees on n nodes is counted by the nth Catalan number (see, e.g., [68, Exercise 6.1.19c])

$$\beta_n := \frac{1}{n+1}\binom{2n}{n},$$

with generating function

$$\mathrm{CAT}(z) := \sum_{n=0}^{\infty} \beta_n z^n = \frac{1}{2z}(1 - \sqrt{1-4z}). \tag{2.3.8}$$

Let $\mu_n(k)$ be defined as the kth moment of the random variable X_n and $\bar{\mu}_n(k) := \beta_n \mu_n(k)/4^n$. By conditioning on the key stored at the root, noting that the probability is $\beta_{l-1}\beta_{n-l}/\beta_n$ that the key stored at the root is l, $1 \le l \le n$, and multiplying throughout by $\beta_n/4^n$ we get, for $n \ge 1$,

$$\bar{\mu}_n(k) = \frac{1}{2} \sum_{j=1}^{n} \frac{\beta_{n-j}}{4^{n-j}} \bar{\mu}_{j-1}(k) + \bar{r}_n(k), \tag{2.3.9}$$

where

$$\bar{r}_n(k) := \frac{1}{4} \sum_{\substack{k_1+k_2+k_3=k\\ k_1,k_2<k}} \binom{k}{k_1,k_2,k_3} t_n^{k_3} \sum_{j=1}^{n} \bar{\mu}_{j-1}(k_1)\bar{\mu}_{n-j}(k_2). \tag{2.3.10}$$

Again we observe that moments of all order satisfy the same basic recurrence, with

$$\bar{r}_n(1) = \frac{t_n}{4^n} \sum_{j=1}^{n} \beta_{j-1}\beta_{n-j} = t_n \frac{\beta_n}{4^n}.$$

We define generating functions

$$\overline{M}_k(z) := \sum_{n=0}^{\infty} \bar{\mu}_n(k) z^n, \qquad \overline{R}_k(z) := \sum_{n=0}^{\infty} \bar{r}_n(k) z^n, \qquad T(z) := \sum_{n=0}^{\infty} t_n z^n. \tag{2.3.11}$$

Then, using (2.3.9) and (2.3.8), we get

$$\overline{M}_k(z) = \frac{\overline{R}_k(z)}{\sqrt{1-z}}, \tag{2.3.12}$$

so to analyze $\overline{M}_k(z)$ it is sufficient to analyze $\overline{R}_k(z)$. With this in mind we introduce the Hadamard product of two power series.

Definition 2.3.3. The *Hadamard product* of two power series F and G, denoted by $F(z) \odot G(z)$ or $(F \odot G)(z)$, is the power series defined by

$$F(z) \odot G(z) := \sum_n f_n g_n z^n,$$

where

$$F(z) = \sum_n f_n z^n \text{ and } G(z) = \sum_n g_n z^n.$$

In light of this definition, the defining relation for $\overline{R}_k(z)$ at (2.3.10) leads to the following relation in terms of generating functions:

$$\overline{R}_k(z) = \sum_{\substack{k_1+k_2+k_3=k \\ k_1,k_2<k}} \binom{k}{k_1, k_2, k_3} (T(z))^{\odot k_3} \odot [\frac{z}{4} \overline{M}_{k_1}(z) \overline{M}_{k_2}(z)], \tag{2.3.13}$$

where, for k a nonnegative integer,

$$T(z)^{\odot k} := \underbrace{T(z) \odot \cdots \odot T(z)}_{k}.$$

Our approach for getting asymptotic expansions for moments of all orders relies on *singularity analysis* of generating functions [31], a tool that, under suitable conditions, allows the translation of asymptotics of generating functions to asymptotics of the underlying sequences. In order to analyze $\overline{M}_k(z)$ and $\overline{R}_k(z)$ inductively (in k), we would like to be able to predict how singularities are composed under Hadamard products. This is precisely the problem studied in Chapter 4. We will, in Chapters 7 and 8, use the results derived in Chapter 4 to get limiting distributions induced by two types of toll functions.

Part II

Theory

CHAPTER 3

Random permutation model: transfer theorems

In Proposition 2.3.1 we have seen that under the random permutation model, moments of each specified order for any additive functional on m-ary search trees satisfy a recurrence relation of the form

$$a_n = b_n + \frac{m}{\binom{n}{m-1}} \sum_{j=0}^{n-(m-1)} \binom{n-1-j}{m-2} a_j, \qquad n \geq m-1, \tag{3.0.1}$$

with specified initial conditions $a_j := b_j$ for $j = 0, \ldots, m-2$. By Theorem 2.3.2 the output generating function

$$A(z) := \sum_{n=0}^{\infty} a_n z^n$$

is given in terms of the input generating function

$$B(z) := \sum_{n=0}^{\infty} b_n z^n$$

by

$$A(z) = \sum_{j=1}^{m-1} c_j (1-z)^{-\lambda_j} + \sum_{j=1}^{m-1} \frac{(1-z)^{-\lambda_j}}{\psi'(\lambda_j)} \int_0^z B^{(m-1)}(\zeta)(1-\zeta)^{\lambda_j+m-2}\, d\zeta,$$

where ψ is the indicial polynomial

$$\psi(\lambda) := \lambda^{\overline{m-1}} - m!$$

with roots $2 =: \lambda_1, \lambda_2, \ldots, \lambda_{m-1}$ listed in nonincreasing order of real parts. We refer the reader to Appendix B for relevant facts about this polynomial, and recall from there the notation

$$\alpha := \max_{2 \leq j \leq m-1} \Re(\lambda_j).$$

The coefficients $c_1, c_2, \ldots, c_{m-1}$ can be written explicitly in terms of the initial conditions $b_0, \ldots, b_{m-2}$. In particular, by Proposition A.4 with $j = 1$ and $g_{\mathrm{c}}^{(k)}(0) = k! b_k$ there, we get [also using (B.8)]

$$c_1 = \frac{1}{H_m - 1} \sum_{j=0}^{m-2} \frac{b_j}{(j+1)(j+2)}. \tag{3.0.2}$$

Here H_m is the mth Harmonic number: $H_m = \sum_{j=1}^{m} \frac{1}{j}$.

In this chapter we will obtain results of both exact and asymptotic natures that link the behavior of the output sequence (a_n) to that of the input sequence (b_n) in (3.0.1).

Note. In the sequel whenever multi-valued functions are encountered, we will assume that we follow the principal branch.

3.1 The Exact Transfer Theorem

Our main result repeats (2.3.5) and then gives an alternative exact transfer from B to A:

Theorem 3.1.1 (Exact Transfer Theorem (ETT)). *Let*

$$\widehat{B}(z) := B(z) - \sum_{j=0}^{m-2} b_j z^j = \sum_{n=m-1}^{\infty} b_n z^n. \tag{3.1.1}$$

Then

$$A(z) = \sum_{j=1}^{m-1} c_j (1-z)^{-\lambda_j} + \sum_{j=1}^{m-1} \frac{(1-z)^{-\lambda_j}}{\psi'(\lambda_j)} \int_0^z B^{(m-1)}(\zeta)(1-\zeta)^{\lambda_j+m-2}\, d\zeta \tag{3.1.2}$$

$$= \sum_{j=1}^{m-1} c_j (1-z)^{-\lambda_j} + \widehat{B}(z) + m! \sum_{j=1}^{m-1} \frac{(1-z)^{-\lambda_j}}{\psi'(\lambda_j)} \int_0^z \widehat{B}(\zeta)(1-\zeta)^{\lambda_j-1}\, d\zeta. \tag{3.1.3}$$

Proof. We begin with (3.1.2), repeated from Theorem 2.3.2, and note that B there can be replaced by $\widehat{B}$. We then use repeated integration by parts and invoke Identity B.3. Denoting

$$\widehat{A}(z) := A(z) - \sum_{j=1}^{m-1} c_j (1-z)^{-\lambda_j},$$

after $m-2$ integrations by parts we find

$$\widehat{A}(z) = \sum_{j=1}^{m-1} \frac{(1-z)^{-\lambda_j}}{\psi'(\lambda_j)} (\lambda_j + m - 2) \cdots (\lambda_j + 1) \int_0^z \widehat{B}'(\zeta)(1-\zeta)^{\lambda_j}\, d\zeta.$$

But

$$(\lambda_j + m - 2) \cdots (\lambda_j + 1) = \frac{\lambda_j^{\overline{m-1}}}{\lambda_j} = \frac{\psi(\lambda_j) + m!}{\lambda_j} = \frac{m!}{\lambda_j},$$

so

$$\widehat{A}(z) = m! \sum_{j=1}^{m-1} \frac{(1-z)^{-\lambda_j}}{\lambda_j \psi'(\lambda_j)} \int_0^z \widehat{B}'(\zeta)(1-\zeta)^{\lambda_j}\, d\zeta.$$

We obtain (3.1.3) by performing one more integration by parts and utilizing Identity B.1 with $\lambda = 0$. □

Remark 3.1.2. For computations, (3.1.2) might be easier to use when it is no bother to compute derivatives of B; otherwise, (3.1.3) is easier. Equation (3.1.3) will be used in establishing part (a) of the Asymptotic Transfer Theorem in Section 3.2; the proof of part (b) will use (3.1.2).

3.2 The Asymptotic Transfer Theorem

It is quite easy to transfer asymptotics for B to asymptotics for A using the ETT. We give three examples important for applications to moments of additive functionals in the next theorem, proved independently in [8] using a quite different approach.

Theorem 3.2.1 (Asymptotic Transfer Theorem (ATT)). *Suppose that* (a_n) *and* (b_n) *satisfy the recurrence* (3.0.1).

(a) If

$$b_n = o(n) \quad \text{and} \quad \sum_{n=0}^{\infty} \frac{b_n}{(n+1)(n+2)} \text{ converges (conditionally),} \tag{3.2.1}$$

then

$$a_n = \frac{K_1}{H_m - 1} n + o(n), \quad \text{where} \quad K_1 := \sum_{j=0}^{\infty} \frac{b_j}{(j+1)(j+2)}. \tag{3.2.2}$$

(b) If $b_n \equiv K_2(n+1) + h_n$ *where* h_n *satisfies* (3.2.1) *[with* (b_n) *replaced by* (h_n)*], then*

$$a_n = \frac{K_2}{H_m - 1} n H_n + \frac{K_3}{H_m - 1} n + o(n), \tag{3.2.3}$$

where

$$K_3 := \sum_{j=0}^{\infty} \frac{h_j}{(j+1)(j+2)} + K_2 \left[\frac{H_m - 1}{2} - 1 + \frac{H_m^{(2)} - 1}{2(H_m - 1)} \right]. \tag{3.2.4}$$

(c) If $b_n = K_4 n^v + o(n^{\Re(v)})$ *with* $\Re(v) > 1$*, then*

$$a_n = \frac{K_4}{1 - \frac{m!\Gamma(v+1)}{\Gamma(v+m)}} n^v + o(n^{\Re(v)}). \tag{3.2.5}$$

Before giving the proof we establish a few lemmas. The first result is found in the proof of Lemma 6 of [8]. For completeness we include a proof. Here, as throughout, $[z^n]F(z)$ is the coefficient of z^n in the series expansion of F.

Lemma 3.2.2. *Let* $Y(z)$ *denote the ordinary generating function of the sequence* (y_n) *with* $y_0 = 0$*. For any* $\lambda \in \mathbb{C}$*,*

$$[z^n] \left((1-z)^{-\lambda} \int_0^z (1-\zeta)^{\lambda-1} Y(\zeta)\, d\zeta \right) = \sum_{k=0}^{n-1} \frac{y_k}{k+1} \prod_{j=k+2}^{n} \left(1 + \frac{\lambda - 1}{j} \right), \quad n \geq 0. \tag{3.2.6}$$

Proof. The function

$$W(z) := (1-z)^{-\lambda} \int_0^z (1-\zeta)^{\lambda-1} Y(\zeta)\, d\zeta$$

is the unique solution with $W(0) = 0$ to the differential equation

$$W'(z) = \lambda(1-z)^{-1} W(z) + (1-z)^{-1} Y(z) :$$

that is, $w_n := [z^n]W(z)$ for $n \geq 0$ satisfies $w_0 = 0$ and

$$w_n = \frac{\lambda}{n} \sum_{k=0}^{n-1} w_k + \frac{1}{n} \sum_{k=0}^{n-1} y_k, \qquad n \geq 1. \tag{3.2.7}$$

To solve the recurrence (3.2.7) we compute $n w_n - (n-1) w_{n-1} = \lambda w_{n-1} + y_{n-1}$ so that

$$w_n = \left(1 + \frac{\lambda - 1}{n} \right) w_{n-1} + \frac{1}{n} y_{n-1}, \qquad n \geq 1.$$

Now we iterate to get the result. ☐

The product in (3.2.6) may be written (when $\lambda \in \mathbb{C} \setminus \{0, -1, -2, \ldots\}$) as

$$\frac{\Gamma(\lambda+n)\Gamma(2+k)}{\Gamma(1+n)\Gamma(\lambda+k+1)},$$

which by Stirling's formula equals

$$\frac{n^{\lambda-1}[1+O(n^{-1})]}{(k+1)^{\lambda-1}[1+O((k+1)^{-1})]} \tag{3.2.8}$$

for $n \geq 1$ and $k \geq 1$. [The product in (3.2.6) equals (3.2.8) even if $\lambda \in \{0, -1, -2, \ldots\}$.] Also, of special interest is the case $\lambda = 2$, in which case (3.2.6) reduces to

$$[z^n]\left((1-z)^{-2}\int_0^z (1-\zeta)Y(\zeta)\,d\zeta\right) = (n+1)\sum_{k=0}^{n-1}\frac{y_k}{(k+1)(k+2)}, \qquad n \geq 0. \tag{3.2.9}$$

For part (a) of the ATT we use the following estimates from [8]. These follow readily from Lemma 3.2.2, (3.2.8), and (3.2.9).

Lemma 3.2.3. *(a) If* $\Re(\lambda) < 2$ *and*

$$Y(z) = \sum_{n=0}^{\infty} y_n z^n$$

satisfies $y_0 = 0$ *and* $y_n = o(n)$*, then*

$$[z^n]\left((1-z)^{-\lambda}\int_0^z Y(\zeta)(1-\zeta)^{\lambda-1}\,d\zeta\right) = o(n).$$

(b) With $\widehat{B}$ *defined at* (3.1.1)*, if* (3.2.1) *holds, then*

$$[z^n]\left((1-z)^{-2}\int_0^z \widehat{B}(\zeta)(1-\zeta)\,d\zeta\right) = n\sum_{j=m-1}^{\infty}\frac{b_j}{(j+1)(j+2)} + o(n).$$

For part (c) of the ATT, we will need the following simple comparison lemma.

Lemma 3.2.4. *If* (b_n) *and* (b'_n) *are two toll sequences such that*

$$|b_n| \leq b'_n \quad \text{for all } n \geq 0,$$

then the corresponding output sequences (a_n) *and* (a'_n) *in* (3.0.1) *(with the initial conditions stated there) satisfy*

$$|a_n| \leq a'_n \quad \text{for all } n \geq 0.$$

Proof. This follows immediately by induction. □

Proof of Theorem 3.2.1. Part (a) follows immediately from (3.1.3), (3.2.1), Lemma 3.2.3, (3.0.2), and (B.8).

For part (b) suppose first that $b_n \equiv n+1$. Then $B(z) = (1-z)^{-2}$, so

$$B^{(m-1)}(z) = m!(1-z)^{-(m+1)}.$$

Plugging this into (3.1.2) we find

$$a_n = (n+1)\left[c_1 + m!\sum_{j=2}^{m-1}\frac{1}{(2-\lambda_j)\psi'(\lambda_j)}\right] + \frac{m!}{\psi'(2)}[z^n][(1-z)^{-2}\ln((1-z)^{-1})] + o(n).$$

Recall Identity B.4; and since $b_n \equiv n+1$, we have in this case by (3.0.2), that $c_1 = 1$. Therefore

$$\begin{aligned} a_n &= (n+1)\left(1 + \frac{1}{2}\left[\frac{H_m^{(2)}-1}{(H_m-1)^2} - 1\right]\right) + \frac{1}{H_m-1}[(n+1)H_n - n] + o(n) \\ &= \frac{1}{H_m-1}nH_n + \left[\frac{1}{2} - \frac{1}{H_m-1} + \frac{H_m^{(2)}-1}{2(H_m-1)^2}\right]n + o(n). \end{aligned}$$

This completes the proof of part (b) for our special case, and the general case follows from this and part (a) using the superposition principle.

Finally for part (c) suppose first that $b_n \equiv (v+1)^{\overline{n}}/n! \sim n^v/\Gamma(v+1)$, so that $B(z) = (1-z)^{-(v+1)}$ and $B^{(m-1)}(z) = (v+1)^{\overline{m-1}}(1-z)^{-(v+m)}$. Plugging this into (3.1.2) and utilizing Identity B.1 with $\lambda = v+1$ and the calculation

$$\frac{(v+1)^{\overline{m-1}}}{(v+1)^{\overline{m-1}} - m!} = \left[1 - \frac{m!\Gamma(v+1)}{\Gamma(v+m)}\right]^{-1},$$

we find

$$A(z) = \left[1 - \frac{m!\Gamma(v+1)}{\Gamma(v+m)}\right]^{-1}(1-z)^{-(v+1)} + O(|1-z|^{-2}).$$

By singularity analysis (see Theorem 4.1.2), this completes the proof of part (c) for our special case.

To complete the proof (by superposition) in the general case, we need only show that if $b_n = o(n^v)$ for real $v > 1$, then $a_n = o(n^v)$. Indeed, fix $\epsilon > 0$; then there exists a sequence (b'_n) such that $|b_n| \le b'_n$ for all n and $b'_n = \epsilon(v+1)^{\overline{n}}/n!$ for all large n. This toll sequence is but a slight modification of our special-case toll sequence, and we see that its output generating function has coefficients

$$a'_n = \epsilon' n^v + o(n^v), \qquad \text{where} \qquad \epsilon' := \frac{\epsilon}{\Gamma(v+1)}\left[1 - \frac{m!\Gamma(v+1)}{\Gamma(v+m)}\right]^{-1}.$$

Now Lemma 3.2.4 implies that

$$\limsup_n |a_n| n^{-v} \le \epsilon';$$

since ϵ (and hence ϵ') can be made arbitrarily small, this completes the proof. □

The conditions (3.2.1) on b_n are not only sufficient but also necessary for asymptotic linearity of a_n. Indeed, here is a converse:

Proposition 3.2.5. *If $a_n = Kn + o(n)$ for some constant K, then* (3.2.1) *holds.*

Proof. Proposition 7 of [8] provides the simple proof that $b_n = o(n)$. Moreover, from (3.1.3), the value of c_1 at (3.0.2), and (3.2.9) with Y taken to be $\widehat{B}$, we find that

$$\sum_{j=0}^{n-1}\frac{b_j}{(j+1)(j+2)} = K(H_m-1) + o(1),$$

i.e., the series

$$\sum_{n=0}^{\infty} \frac{b_n}{(n+1)(n+2)}$$

converges [conditionally, to $K(H_m - 1)$]. $\square$

The following additional asymptotic transfer results are established by calculations similar to those in the proof of the ATT.

Definition 3.2.6. A nonnegative function $L(n)$ defined for $n \geq n_0 \geq 0$ and not identically zero is called *slowly varying* if

$$\frac{L(n)}{L(\lfloor tn \rfloor)} \to 1 \quad \text{as } n \to \infty$$

for every $t > 0$. If $n_0 > 0$, we define $L(n) = 0$ for $0 \leq n < n_0$.

Examples of slowly varying functions include any powers of $\ln n$ or $\ln \ln n$. The key property of slowly varying functions that we employ is the regular variation fact [4, 1.5.9a], that

$$L(n) = o\left(\sum_{k \leq n} \frac{L(k)}{k} \right). \tag{3.2.10}$$

Theorem 3.2.7. *Consider the initial value problem* (2.3.3).

(a) If $2 \leq m \leq 26$ *and* $b_n = o(\sqrt{n})$*, then we can refine* (3.2.2) *to*

$$a_n = \frac{K_1}{H_m - 1} n + o(\sqrt{n}). \tag{3.2.11}$$

(b) If $\Re(\lambda_2) =: \alpha < 1 + \beta$ *and* $b_n \sim n^{\beta} L(n)$ *with* L *slowly varying, then we can refine* (3.2.2) *to*

$$a_n = \frac{K_1}{H_m - 1} n = \frac{(1+\beta)^{\overline{m-1}}}{m! - (1+\beta)^{\overline{m-1}}} n^{\beta} L(n) + o(n^{\beta} L(n)). \tag{3.2.12}$$

(c) If $b_n \sim nL(n)$ *with* L *slowly varying, then, with* K_1 *defined at* (3.2.2),

$$a_n \sim \begin{cases} \dfrac{K_1}{H_m - 1} n, & \textit{if } \displaystyle\sum^{\infty} \frac{L(k)}{k} < \infty, \\ \dfrac{1}{H_m - 1} n \displaystyle\sum_{k \leq n} \frac{L(k)}{k}, & \textit{if } \displaystyle\sum^{\infty} \frac{L(k)}{k} = \infty. \end{cases} \tag{3.2.13}$$

(d) Part (c) of the ATT can be extended as follows. If $b_n = K_4 n^v L(n) + o(n^{\Re(v) L(n)})$ *with* $\Re(v) > 1$ *and* L *slowly varying, then*

$$a_n = \frac{K_4}{1 - \frac{m!\Gamma(v+1)}{\Gamma(v+m)}} n^v L(n) + o(n^{\Re(v)} L(n)). \tag{3.2.14}$$

Proof sketch. Whenever the conditions (3.2.1) are met we have by the ETT and (3.2.9)

$$
\begin{aligned}
a_n - \frac{K_1}{H_m - 1}(n+1) = O(n^{\alpha-1}) + b_n - \frac{1}{H_m - 1}(n+1) \sum_{k=n}^{\infty} \frac{\hat{b}_k}{(k+1)(k+2)} \\
+ m! \sum_{j=2}^{m-1} \frac{1}{\psi'(\lambda_j)} [z^n] \left((1-z)^{-\lambda_j} \int_0^z \widehat{B}(\zeta)(1-\zeta)^{\lambda_j - 1}\, d\zeta \right), \quad (3.2.15)
\end{aligned}
$$

where $\alpha := \Re(\lambda_2)$ is the real part of the root of ψ with second largest real part, which is smaller than $3/2$ when $m \leq 26$ (see Appendix B). Simple estimates, including the use of (3.2.8), give cases (a) and (b); for (b), the coefficient of $n^\beta L(n)$ in (3.2.12) indeed is, using Identity B.1 and (B.8),

$$
1 - \frac{1}{(1-\beta)H_m - 1} + m! \sum_{j=2}^{m-1} \frac{1}{((1+\beta) - \lambda_j)\psi'(\lambda_j)} = -\frac{(1+\beta)^{\overline{m-1}}}{m! - (1+\beta)^{\overline{m-1}}}.
$$

In case (c), from the ETT result (3.1.3) and simple estimates we find

$$
a_n = (n+1) \frac{1}{H_m - 1} \sum_{k=0}^{m-1} \frac{b_k}{(k+1)(k+2)} + O(nL(n)).
$$

To finish, we use the regular variation fact (3.2.10).

In case (d), again from (3.1.3) and simple estimates we find

$$
a_n = O(n) + b_n + K_4 n^v L(n) m! \sum_{j=1}^{m-1} \frac{1}{\psi'(\lambda_j)(v + 1 - \lambda_j)} + o(n^v L(n)).
$$

The proof is completed by using Identity B.1. □

Chapter 4

Singularity analysis of Hadamard products

4.1 Background

Singularity analysis is "a class of methods by which one can translate, on a term-by-term basis, an asymptotic expansion of a function around a dominant singularity into a corresponding asymptotic expansion for the Taylor coefficients of the function" [31]. In general, it provides sufficient conditions for the validity of the implications

$$\begin{aligned} F(z) = O(G(z)) &\implies f_n = O(g_n) \\ F(z) = o(G(z)) &\implies f_n = o(g_n) \\ F(z) \sim G(z) &\implies f_n \sim g_n, \end{aligned}$$

where F and G are the ordinary generating functions of the sequences (f_n) and (g_n) respectively. The sufficiency conditions are expressed in terms of the following analytic continuation condition.

Definition 4.1.1. A function defined by a Taylor series with radius of convergence equal to 1 is said to be Δ-*regular* if, with the sole exception of $z = 1$, it can be analytically continued in a domain

$$\Delta(\phi, \eta) := \{z : |z| \leq 1 + \eta,\ |\arg(z-1)| \geq \phi\}, \tag{4.1.1}$$

for some $\eta > 0$ and $0 < \phi < \pi/2$.

A function f is said to admit a Δ-*singular expansion* at $z = 1$ if it is Δ-regular and

$$f(z) = \sum_{n=0}^{N} c_n (1-z)^{\alpha_n} + O(|1-z|^A) \tag{4.1.2}$$

as $z \to 1$ in $\Delta(\phi, \eta)$, for a sequence of complex numbers $(c_n)_{n=0}^N$ and a sequence of real numbers $(\alpha_n)_{n=0}^N$ satisfying $\alpha_n < A$.

Note that under these assumptions the $O(\cdot)$ error term in (4.1.2) holds uniformly in $z \in \Delta(\phi, \eta)$. We also allow in the usual way infinite asymptotic expansions

$$f(z) \sim \sum_{n \geq 0} c_n (1-z)^{\alpha_n}$$

to represent an infinite collection of expansions of type (4.1.2). When it is convenient, we will also allow powers of $\ln\left((1-z)^{-1}\right)$ in singular expansions. The region of analytic continuation in the complex plane is depicted in Figure 4.1.1. We will sometimes refer to such a region as an *indented crown.*

For completeness we state relevant results from [31] and [28]. These will be used frequently from here on.

Theorem 4.1.2 (Theorem 1 in [31]). *Let $f(z)$ be a Δ-regular function in $\Delta(\phi, \eta)$ and assume that, as $z \to 1$ in $\Delta(\phi, \eta)$,*

$$f(z) = O(|1-z|^{-\alpha})$$



Figure 4.1.1: The region $\Delta(\phi, \eta)$ in Definition 4.1.1.

for some real number α. Then the nth Taylor coefficient of $f(z)$ satisfies

$$f_n = [z^n]f(z) = O(n^{\alpha-1}).$$

Flajolet also showed [28] that singularity analysis is likewise applicable to the *generalized polylogarithm* $\mathrm{Li}_{\alpha,r}(z)$ defined for a nonnegative integer r and an arbitrary complex number α and $|z| < 1$ as

$$\mathrm{Li}_{\alpha,r}(z) := \sum_{n=1}^{\infty} \frac{(\ln n)^r}{n^\alpha} z^n. \tag{4.1.3}$$

Theorem 4.1.3 (Theorem 1 in [28]). *The function $\mathrm{Li}_{\alpha,0}(z)$ is analytic in the slit plane $\mathbb{C} \setminus [1, +\infty)$. For $\alpha \notin \{1, 2, \ldots\}$ the function $\mathrm{Li}_{\alpha,0}(z)$ satisfies the singular expansion*

$$\mathrm{Li}_{\alpha,0}(z) \sim \Gamma(1-\alpha)t^{\alpha-1} + \sum_{j\geq 0} \frac{(-1)^j}{j!}\zeta(\alpha - j)t^j,$$

with

$$t = -\ln z = \sum_{l=1}^{\infty} \frac{(1-z)^l}{l}$$

as $z \to 1$ in the sector $-\pi + \epsilon < \arg(1-z) < \pi - \epsilon$ for any $\epsilon > 0$.

For $r = 1, 2, \ldots$, the singular expansion of $\mathrm{Li}_{\alpha,r}(z)$ is obtained by

$$\mathrm{Li}_{\alpha,r}(z) = (-1)^r \frac{\partial^r}{\partial \alpha^r} \mathrm{Li}_{\alpha,0}(z).$$

The case when $\alpha = m \in \{1, 2, \ldots\}$ is also handled in [28, p. 380]:

$$\mathrm{Li}_{m,r}(z) = \operatorname{Res}\left((-1)^r \zeta^{(r)}(s+m)\Gamma(s)t^{-s}\right)_{s=1-m} + \sum_{j\geq 0, j\neq m-1} \frac{(-1)^{j+r}}{j!}\zeta^{(r)}(m-j)t^j \tag{4.1.4}$$

with $t = -\ln z$, where $\operatorname{Res}(\cdot)$ denotes a residue.

Singularity analysis is a very useful tool in the analysis of combinatorial structures mainly because generating functions of structures that are constructed combinatorially readily admit the technical conditions needed for its application. In such cases precise asymptotic expansions of the coefficients can be derived. In the next two sections we will extend singularity analysis so that it is applicable to the right-hand side of (2.3.13).

4.2 Singular expansions and differentiation

In preparation for the treatment of Hadamard products in Section 4.3, we need a theorem that enables us to differentiate local expansions of analytic functions around a singularity. Such a result cannot of course be unconditionally true; see, for example, (4.2.5). However, it turns out that the class of functions amenable to singularity analysis satisfy this property.

Theorem 4.2.1. *If $f(z)$ is Δ-regular and admits a Δ-expansion near its singularity in the sense of Definition 4.1.1, then for each integer $k > 0$, $\frac{d^k}{dz^k}f(z)$ is also Δ-regular and admits a Δ-expansion obtained through term-by-term differentiation:*

$$\frac{d^k}{dz^k}f(z) = \sum_{n\geq 0} c_n(-1)^k \frac{\Gamma(\alpha_n+1)}{\Gamma(\alpha_n+1-k)}(1-z)^{\alpha_n-k} + O(|1-z|^{A-k}).$$

Proof. Clearly, all that is required is to establish

$$\frac{d}{dz}O(|1-z|^A) = O(|1-z|^{A-1}).$$

Let us define $g(z) := f(z)/(1-z)^A$ where $f(z) = O(|1-z|^A)$. By differentiation, we find

$$g'(z) = \frac{f'(z)}{(1-z)^A} + A\frac{f(z)}{(1-z)^{A+1}}.$$

This can be rewritten as

$$f'(z) = g'(z)(1-z)^A - A\frac{f(z)}{1-z}.$$

Now $g(z)$ is by assumption $O(1)$ at $z = 1$. If we assume that $g'(z) = O(|1-z|^{-1})$ at $z = 1$, we obtain the sought result: $f'(z) = O(|1-z|^{A-1})$. Thus we need only prove the implication

$$g(z) = O(1) \implies g'(z) = O(|1-z|^{-1}), \tag{4.2.1}$$

or, in more symbolic terms,

$$\frac{d}{dz}O(1) = O\left(\frac{1}{1-z}\right).$$

We are going to prove (4.2.1) in a domain $\Delta(\phi+\epsilon, \eta)$, where $\epsilon > 0$ should be fixed but can be taken as small as we please. The starting point is Cauchy integral formula

$$g(z) = \frac{1}{2\pi i}\int_\gamma g(w)\frac{dw}{w-z},$$

which, through differentiation, provides

$$g'(z) = \frac{1}{2\pi i}\int_\gamma g(w)\frac{dw}{(w-z)^2}.$$



Figure 4.2.1: The contour γ used in the proof of the differentiation theorem.

We select a small $\epsilon > 0$ and restrict z to the domain $\Delta(\phi+\epsilon,\eta)$. The integral is estimated using the contour γ depicted in Figure 4.2.1: $\gamma := \gamma_1 \cup \gamma_2 \cup \gamma_3 \cup \gamma_4$, with

$$\begin{aligned}
\gamma_1 &:= \{w \colon |w-1| = \tfrac{1}{2}|z-1|,\ |\arg(w-1)| \geq \phi\},\\
\gamma_2 &:= \{w \colon \tfrac{1}{2}|z-1| \leq |w-1|,\ |w| \leq 1+\eta, \arg(w-1) = \phi\},\\
\gamma_3 &:= \{w \colon |w| = 1+\eta,\ |\arg(w-1)| \geq \phi\},\\
\gamma_4 &:= \{w \colon \tfrac{1}{2}|z-1| \leq |w-1|,\ |w| \leq 1+\eta, \arg(w-1) = -\phi\}.
\end{aligned}$$

We then decompose the integral giving $g'(z)$ as $\int_\gamma = \int_{\gamma_1} + \int_{\gamma_2} + \int_{\gamma_3} + \int_{\gamma_4}$. Along γ_1 we apply trivial upper bounds:

$$\left|\int_{\gamma_1}\right| = O(1)\cdot O\left(\frac{1}{|1-z|^2}\right)\cdot O(|1-z|) = O(|1-z|^{-1}). \tag{4.2.2}$$

The integral along the external circle γ_3 satisfies trivially

$$\left|\int_{\gamma_3}\right| = O(1) = O(|1-z|^{-1}). \tag{4.2.3}$$

If we set $|1-z| = \delta$, we find that the integrals along γ_2 and γ_4 are of no bigger order than

$$\int_{\delta/2}^{\infty} = \frac{du}{\delta^2+u^2} = O\left(\frac{1}{\delta}\right) = O(|1-z|^{-1}).$$

Thus we have

$$\left|\int_{\gamma_2}\right|,\ \left|\int_{\gamma_4}\right| = O\left(|1-z|^{-1}\right). \tag{4.2.4}$$

Combining the estimates (4.2.2)–(4.2.4) yields the asymptotic form of $g'(z)$ as $O(|1-z|^{-1})$, as required at (4.2.1). ☐

Thus, for instance, taking

$$g(z) = \cos\ln\left(\frac{1}{1-z}\right) \quad \text{and} \quad g'(z) = -\frac{1}{1-z}\sin\ln\left(\frac{1}{1-z}\right),$$

we correctly predict that $g(z) = O(1) \Rightarrow g'(z) = O(|1-z|^{-1})$. On the other hand, the apparent paradox given by the pair

$$g(z) = \cos\exp\left(\frac{1}{1-z}\right) \quad \text{and} \quad g'(z) = -\frac{1}{(1-z)^2}\exp\left(\frac{1}{1-z}\right)\sin\exp\left(\frac{1}{1-z}\right), \tag{4.2.5}$$

is resolved by observing that in no nondegenerate sector around $z = 1$ do we have $g(z) = O(1)$.

It is also well known that integration of asymptotic expansions is usually easier than differentiation, and we can indeed prove the following theorem.

Theorem 4.2.2. *If $f(z)$ has a Δ-expansion that reduces to $f(z) = O(|1-z|^A|\log(1-z)|^B)$, then we have*

$$\int_0^z f(t)\,dt = \begin{cases} O(|1-z|^{A+1}|\log(1-z)|^B) & \text{if } A < -1 \\ O(|\log(1-z)|^{B+1}) & \text{if } A = -1,\ B \neq -1 \\ O(|\log\log(1-z)^{-1}|) & \text{if } A = -1,\ B = -1 \\ \displaystyle\int_0^1 f(t)\,dt + O(|1-z|^{A+1}|\log(1-z)|^B) & \text{if } A > -1. \end{cases}$$

Proof. The Δ-expansion condition tells us that $|f(z)| \le K|1-z|^A|\ln(1-z)|^B$ for some $0 \le K < \infty$.

Suppose $A < -1$. By Cauchy integral formula, we can choose any path of integration that stays within the region of analyticity of f. We choose the contour $\gamma := \gamma_1 \cup \gamma_2$, shown in Figure 4.2.2.

Then

$$\begin{aligned}\left|\int_\gamma f(t)\,dt\right| &= \left|\int_{\gamma_1} f(t)\,dt + \int_{\gamma_2} f(t)\,dt\right| \le \left|\int_{\gamma_1} f(t)\,dt\right| + \left|\int_{\gamma_2} f(t)\,dt\right| \\ &\le K\int_{\gamma_1} |1-t|^A|\ln(1-t)|^B\,dt + K\int_{\gamma_2} |1-t|^A|\ln(1-t)|^B\,|dt|.\end{aligned}$$

The last integral along γ_2 is

$$|1-z|^A|\ln(1-z)|^B\int_{\gamma_2}|dt| = O(|1-z|^{A+1}|\log(1-z)|^B).$$

The last integral along γ_1 is a real integral. Define $x := 1-|1-z|$, so that the last integral



Figure 4.2.2: The contour used in the proof of the integration theorem.

along γ_1 equals

$$\begin{aligned}
&\int_0^x (1-t)^A(\ln(1-t)^{-1})^B\,dt\\
&= \frac{1}{|A|-1}\left[(1-x)^{A+1}(\ln(1-x)^{-1})^B - B\int_0^x (1-t)^A(\ln(1-t)^{-1})^{B-1}\,dt\right]\\
&= \frac{1}{|A|-1}(1-x)^{A+1}(\ln(1-x)^{-1})^B + o\left(\int_0^x (1-t)^A(\log(1-t)^{-1})^B\,dt\right).
\end{aligned}$$

We have shown that

$$\begin{aligned}
\int_{\gamma_1} |1-t|^A|\ln(1-t)|^B\,dt &= (1+o(1))\frac{1}{|A|-1}(1-x)^{A+1}(\ln(1-x)^{-1})^B\\
&= (1+o(1))\frac{1}{|A|-1}|1-z|^{A+1}(\ln|1-z|^{-1})^B\\
&= (1+o(1))\frac{1}{|A|-1}|1-z|^{A+1}|\ln(1-z)^{-1}|^B\\
&= \Theta(|1-z|^{A+1}|\log(1-z)^{-1}|^B)\\
&= O(|1-z|^{A+1}|\log(1-z)^{-1}|^B).
\end{aligned}$$

Hence

$$\int_\gamma f(t)\,dt = O(|1-z|^{A+1}|\log(1-z)|^B).$$

The case $A=-1$ is similar. As above,

$$\left|\int_\gamma f(t)\,dt\right| \le K\int_{\gamma_1} |1-t|^{-1}|\ln(1-t)|^B\,dt + K\int_{\gamma_2} |1-t|^{-1}|\ln(1-t)|^B\,|dt|.$$

The integral here along γ_2 is

$$|1-z|^{-1}|\ln(1-z)|^B\int_{\gamma_2}|dt| = O(|\log(1-z)|^B) = o(|\log(1-z)|^{B+1}).$$

The integral along γ_1 is a real integral. If $B \neq -1$, then setting $x := 1-|1-z|$, we integrate to get

$$\int_0^x (1-t)^{-1}(\ln(1-t)^{-1})^B\,dt = \frac{1}{B+1}(\ln(1-x)^{-1})^{B+1} = O(|\log(1-z)|^{B+1}).$$

If $B=-1$,

$$\left|\int_{\gamma_1} f(t)\,dt\right| \le O(1) + K\int_{1/2}^x (1-t)^{-1}(\ln(1-t)^{-1})^{-1}\,dt = O(|\log\log(1-z)^{-1}|).$$

Combining the two integrals gives us the result.

The case $A > -1$ is obtained by observing that the asymptotic condition guarantees the existence of $\int_0^1$ and by decomposing $\int_0^z = \int_0^1 + \int_1^z$. The contour of integration is the line segment joining 1 to z. It is enough to show that

$$\left|\int_z^1 f(t)\,dt\right| = O(|1-z|^{A+1}|\log(1-z)|^B).$$

Indeed, performing the change of variable $t = uz + (1-u)$, we can bound the absolute value of this integral by K times

$$\begin{aligned}&\int_0^1 |1-z|^A u^A |\ln(u(1-z))|^B |1-z|\,du\\ &= |1-z|^{A+1}|\ln(1-z)|^B\int_0^1 u^A\left(\frac{|\ln(u(1-z))|}{|\ln(1-z)|}\right)^B du = O(|1-z|^{A+1}|\log(1-z)|^B),\end{aligned}$$

since the last integral is bounded. □

4.3 Hadamard products

In this section we examine the way singular expansions are composed under Hadamard products. The Hadamard product is a bilinear form. So if we have a set of functions admitting known singular expansions, we need to establish their composition law, and this will give composition rules for finite terminating expansions. In order to extend this to asymptotic expansions, either finite with $O(|1-z|^A)$ error terms or infinite, we need to establish a theorem providing the form of

$$O(|1-z|^A)\odot O(|1-z|^B).$$

The composition rule for polylogarithms [see (4.1.3)] is trivial, since

$$\mathrm{Li}_{\alpha,r}(z)\odot\mathrm{Li}_{\beta,s}(z) = \mathrm{Li}_{\alpha+\beta,r+s}(z).$$

However, polylogarithms do not have a simple composition rule with respect to ordinary products.

We thus turn to the composition rule for the basis formed by functions of the form $(1-z)^{-\alpha}$, where α may be any complex number. From the expansion

$$(1-z)^{-\alpha} = 1 + \frac{\alpha}{1}z + \frac{\alpha(\alpha+1)}{2!}z^2 + \cdots$$

around the origin, we get through term-by-term multiplication,

$$(1-z)^{-\alpha} \odot (1-z)^{-\beta} = 1 + \frac{\alpha\beta}{(1!)^2} z + \frac{\alpha(\alpha+1)\beta(\beta+1)}{(2!)^2} z^2 + \cdots . \tag{4.3.1}$$

The classical *hypergeometric function* of Gauss is defined for γ not a nonpositive integer by

$${}_2F_1[\alpha, \beta; \gamma; z] = 1 + \frac{\alpha\beta}{\gamma} \frac{z}{1!} + \frac{\alpha(\alpha+1)\beta(\beta+1)}{\gamma(\gamma+1)} \frac{z^2}{2!} + \cdots ,$$

so that (4.3.1) reduces to

$$(1-z)^{-\alpha} \odot (1-z)^{-\beta} = {}_2F_1[\alpha, \beta; 1; z]. \tag{4.3.2}$$

From the transformation theory of hypergeometrics—see, e.g., [37, p. 163]—we know that, in general, hypergeometrics can be expanded in the vicinity of $z = 1$ by means of the $z \mapsto 1 - z$ transformation. If we instantiate this transformation with $\gamma = 1$, we get

$$\begin{aligned} {}_2F_1[\alpha, \beta; 1; z] = {} & \frac{\Gamma(1-\alpha-\beta)}{\Gamma(1-\alpha)\Gamma(1-\beta)}\, {}_2F_1[\alpha, \beta; \alpha+\beta; 1-z] \\ & + \frac{\Gamma(\alpha+\beta-1)}{\Gamma(\alpha)\Gamma(\beta)} (1-z)^{-\alpha-\beta+1}\, {}_2F_1[1-\alpha, 1-\beta; 2-\alpha-\beta; 1-z]. \end{aligned} \tag{4.3.3}$$

In other words, we can state the following theorem.

Proposition 4.3.1. *When α, β, and $\alpha+\beta$ are not integers, the Hadamard product*

$$(1-z)^{-\alpha} \odot (1-z)^{-\beta}$$

has an infinite Δ-expansion with exponent scale $\{0, 1, 2, \ldots\} \cup \{-\alpha-\beta+1, -\alpha-\beta+2, \ldots\}$, namely,

$$(1-z)^{-\alpha} \odot (1-z)^{-\beta} \sim \sum_{k \geq 0} \lambda_k^{(\alpha,\beta)} \frac{(1-z)^k}{k!} + \sum_{k \geq 0} \mu_k^{(\alpha,\beta)} \frac{(1-z)^{-\alpha-\beta+1+k}}{k!},$$

where the coefficients λ and μ are given by

$$\begin{aligned} \lambda_k^{(\alpha,\beta)} &= \frac{\Gamma(1-\alpha-\beta)}{\Gamma(1-\alpha)\Gamma(1-\beta)} \frac{\alpha^{\bar{k}} \beta^{\bar{k}}}{(\alpha+\beta)^{\bar{k}}}, \\ \mu_k^{(\alpha,\beta)} &= \frac{\Gamma(\alpha+\beta-1)}{\Gamma(\alpha)\Gamma(\beta)} \frac{(1-\alpha)^{\bar{k}} (1-\beta)^{\bar{k}}}{(2-\alpha-\beta)^{\bar{k}}}. \end{aligned}$$

We are thus facing a situation where polylogarithms are simple for Hadamard products and complicated for ordinary products, with the reverse situation for power functions.

We now examine how $O(\cdot)$ terms get composed. The task is easier when the resulting function becomes large at its singularity. Fortunately, thanks to the results of Section 4.2, all cases can be reduced (see Proposition 4.3.4) to the following.

Proposition 4.3.2. *Assume that $f(z)$ and $g(z)$ are Δ-regular in $\Delta(\psi_0, \eta)$. Additionally, suppose that*

$$f(z) = O(|1-z|^a) \text{ and } g(z) = O(|1-z|^b) \text{ as } z \to 1,$$

where a and b satisfy $a + b + 1 < 0$. Then the Hadamard product $(f \odot g)(z)$ is Δ-regular and *admits the expansion*

$$(f \odot g)(z) = O(|1-z|^{a+b+1}).$$

Proof. We first give the proof in the simpler case when$f(z)$ and $g(z)$ are analytic in the whole of the complex plane slit along $z \geq 1$.

Assume that s is a complex number satisfying $|s| < 1$. Then consider the integral

$$I = \frac{1}{2\pi i} \int_{\gamma_0} f(w) g\left(\frac{s}{w}\right) \frac{dw}{w}, \tag{4.3.4}$$

taken (counterclockwise) along a contour γ_0 (see Figure 4.3.1) which is simply a circle of radius R centered at the origin and such that $|s| < R < 1$. Setting

$$f(z) = \sum_{n\geq 0} f_n z^n \qquad \text{and} \qquad g(z) = \sum_{n \geq 0} g_n z^n,$$

and computing the integral (4.3.4) by expanding the functions, we find

$$I = \sum_{n\geq 0} f_n g_n s^n.$$

This is the classical formula of Hadamard for Hadamard products.



Figure 4.3.1: The contour γ_0.

We propose to continue $f \odot g$ to a point z such that $|z| > 1$ where z/s is real and positive. Because of the analytic continuation properties of f and g, we can take as the contour of integration in (4.3.4) the contour γ depicted in Figure 4.3.2. The quantity $f \odot g$ evaluated at s is still equal to the integral form along γ since there is a continuous deformation from γ_0 to γ such that, along the intermediate contours, both w and s/w stay within the domains of analyticity of f and g.

We now consider the integral in (4.3.4) as a function of the parameter s, and let s vary continuously from z along the ray from the origin to z. The integrand remains an analytic



Figure 4.3.2: The contour γ.

function of both s and z so that the integral (4.3.4) provides the analytic continuation of $f \odot g$. Using this contour, we thus capture the singular behaviors of both f and g.

In order to obtain the asymptotics for $f \odot g$, we fully specify in Figure 4.3.3 the geometry of the contour γ. Without loss of generality, we assume that z is in the upper half-plane.



Figure 4.3.3: The geometry of the contour γ.

Define

$$\theta := \arg z, \qquad \phi := \arcsin \frac{c\delta}{R},$$

$$\alpha_1 := Re^{i(\theta-\phi)}, \qquad \beta_1 := z - c\delta e^{i(\theta+\frac{\pi}{2})},$$

$$\alpha_2 := Re^{i(\theta+\phi)}, \qquad \beta_2 := z + c\delta e^{i(\theta+\frac{\pi}{2})},$$

where we can (it is not hard to see) and do choose $0 < c \leq \frac{1}{4}\sin\psi_0$ small enough that γ (and its interior) stays entirely within the interior of $\Delta(\psi_0, \eta)$. (The reason for the restriction $c \leq \frac{1}{4}\sin\psi_0$ will become clear later.)

Then, the contour γ is $\gamma_1 \cup \gamma_2 \cup \gamma_3 \cup \gamma_4$, where

$$\begin{aligned}
\gamma_1 &= \{w \colon |w| = R,\ \arg w \in [0, \theta - \phi] \cup [\theta + \phi, 2\pi]\},\\
\gamma_2 &= \{w \colon w = (1-t)\alpha_1 + t\beta_1,\ 0 \leq t \leq 1\},\\
\gamma_3 &= \{w \colon |w - z| = \delta/2,\ \arg(w - z) \in [0, \pi]\},\\
\gamma_4 &= \{w \colon w = t\alpha_2 + (1-t)\beta_2,\ 0 \leq t \leq 1\}.
\end{aligned}$$

By the asymptotic conditions on f and g, we know that there are positive constants K_1 and K_2 such that $f(w) \leq K_1|1-w|^a$ and $g(w) \leq K_2|1-w|^b$ for all w on γ. We also note that $R \leq |w| \leq 1 + \eta$, so we can find a constant K such that

$$\begin{aligned}
|(f \odot g)(z)| = \left|\frac{1}{2\pi i}\int_\gamma f(w) g\left(\frac{z}{w}\right)\frac{dw}{w}\right| &\leq \frac{K}{2\pi}\int_\gamma |1-w|^a |w-z|^b \, |dw|\\
&\leq \frac{K}{2\pi}\left(\int_{\gamma_1} + \int_{\gamma_2} + \int_{\gamma_3} + \int_{\gamma_4}\right) |1-w|^a |w-z|^b \, |dw|. \qquad (4.3.5)
\end{aligned}$$

We now show that each of the four integrals in (4.3.5) is $O(\delta^{a+b+1}) = O(|1-z|^{a+b+1})$, completing the proof of (4.3.2).

1. Uniformly along γ_1, the integrand is $\Theta(1)$. Hence the integral is $\Theta(1)$, which is $O(\delta^{a+b+1})$ since $a + b + 1 < 0$.

2. Along γ_2, we can express the variable of integration w as

   $$w \equiv w(u) = (\alpha_1 - \beta_1)u + \beta_1, \quad 0 \leq u \leq 1,$$

   so that

   $$dw = (\alpha_1 - \beta_1)du.$$

   Note that $|\beta_1 - \alpha_1| = (1 + o(1))(1 - R)$. The integral of interest is then bounded by a constant times

   $$\int_0^\infty |1 - w(u)|^a |w(u) - z|^b \, du.$$

   We split this integral into two pieces, one where $u \leq M\delta$, say γ_{21}, and the other where $u > M\delta$, say γ_{22}, where M is a suitably chosen large constant. We will show that each of these integrals is $O(\delta^{a+b+1})$. To this end, observe that

   $$\begin{aligned}
   1 - w = 1 - \beta_1 + (\beta_1 - \alpha_1)u &= 1 - z + c\delta e^{i(\theta + \frac{\pi}{2})} + u|\beta_1 - \alpha_1|e^{i\theta}\\
   &= e^{i\theta}(e^{-i\theta} - |z| + ic\delta + u|\beta_1 - \alpha_1|), \qquad (4.3.6)
   \end{aligned}$$

   We defer to Lemma 4.3.3 a calculation that establishes $|1 - w| = \Omega(\delta)$ uniformly along all of γ_2. To get an upper bound, using the triangle inequality,

   $$|1 - w| \leq |1 - \beta_1| + |\beta_1 - w| \leq \delta + c\delta + |\beta_1 - w| = (c+1)\delta + |\beta_1 - w|.$$

   Restricting attention to γ_{21} gives $|\beta_1 - w| \leq |\beta_1 - \alpha_1|M\delta$. Hence, uniformly along γ_{21}, $|1 - w| = \Theta(\delta)$.

   For $|w - z|$, a trivial lower bound is $|w - z| \geq |\beta_1 - z| = c\delta$. Along γ_{21}, an upper bound is established by the triangle inequality as

$$|w-z| \leq |\beta_1 - z| + |w - \beta_1| \leq c\delta + |\beta_1 - \alpha_1| M\delta.$$

Hence $|w-z| = \Theta(\delta)$ uniformly along γ_{21}, so that the integral along γ_{21} is $\Theta(\delta^{a+b+1})$ since the integrand is uniformly $\Theta(\delta^{a+b})$ and the path of integration is of length $\Theta(\delta)$.

For the integral along γ_{22}, we will establish that $|1-w| = \Theta(u)$ and $|w-z| = \Theta(u)$ for suitably large M. The desired bound on the integral will then follow, since $a+b+1<0$.

We use (4.3.6) and the fact that the real part of a complex number is a lower bound on its absolute value to get

$$|1-w| \geq \cos\theta - |z| + |\beta_1 - \alpha_1| u.$$

Writing $z = 1 + \delta e^{i\psi}$, we note that

$$\cos\theta = \frac{1+\delta\cos\psi}{|z|} \geq \frac{1+\delta\cos\psi}{1+\delta} \geq \frac{1-\delta}{1+\delta} \geq 1 - 2\delta.$$

Hence

$$|1-w| \geq 1 - 2\delta - (1+\delta) + (1+o(1))(1-R)u = (1+o(1))(1-R)u - 3\delta.$$

Since $u \geq M\delta$, if we choose M large enough, we get $|1-w| = \Omega(u)$.

For the upper bound,

$$|1-w| \leq |w-\beta_1| + |1-\beta_1| = |\alpha_1 - \beta_1| u + |1-\beta_1| \leq (1+o(1))(1-R)u + (c+1)\delta = O(u).$$

Hence $|1-w| = \Theta(u)$.

The estimate for $|w-z|$ is similar. For the upper bound, observe that

$$w - z = \beta_1 - z + (\alpha_1 - \beta_1)u,$$

whence

$$|w-z| \leq |\beta_1 - z| + |\alpha_1 - \beta_1| uc\delta + (1+o(1))(1-R)u = O(u).$$

For the lower bound, we express

$$w - z = -c\delta e^{i(\theta + \frac{\pi}{2})} + u|\alpha_1 - \beta_1| e^{i\theta} = e^{i\theta}(u|\alpha_1 - \beta_1| - ic\delta),$$

so that

$$|w-z| = |u|\alpha_1 - \beta_1| - ic\delta| \geq u|\alpha_1 - \beta_1| = (1+o(1))(1-R)u.$$

This establishes $|w-z| = \Omega(u)$ and hence $|w-z| = \Theta(u)$.

3. Along γ_3, $|w-z| = c\delta$, and $(1-c)\delta = \delta - c\delta \leq |1-w| \leq \delta + c\delta = (1+c)\delta$ using the triangle inequality. The contour of integration has length $\pi c\delta$. Hence the integral is $\Theta(\delta^{a+b+1}) = O(|1-z|^{a+b+1})$.

4. Along γ_4, the argument is similar to (and somewhat simpler than) that for the integral along γ_2.

□

We now present the proof of the lower-bound on $|1-w|$ along γ_2 used in the proof of Proposition 4.3.2.

Lemma 4.3.3. *In the notation of the proof of Proposition 4.3.2,* $|1-w| = \Omega(\delta)$ *uniformly along all of* γ_2*.*

Proof. If $\sin\theta > 2c\delta$, we lower-bound $|1-w|$ using (4.3.6) and the fact that the imaginary part of a complex number is a lower bound on the its absolute value,

$$|1-w| \geq -\Im(e^{-i\theta} - |z| + ic\delta + u|\beta_1 - \alpha_1|) = \sin\theta - c\delta > c\delta.$$

Otherwise, we use the real part as a lower bound:

$$|1-w| \geq \Re(e^{-i\theta} - |z| + ic\delta + u|\beta_1 - \alpha_1|) \geq \cos\theta - |z| + u|\beta_1 - \alpha_1|.$$

We choose c small enough so that if $\Re(z) = 1 - |z|\cos\theta < 0$, then $\sin\theta > 2c\delta$. Indeed, if $\Re(z) > 1$, then $z = 1 + \delta e^{i\psi}$ where $\psi_0 < \psi < \pi/2$, and so

$$\sin\theta = \frac{\delta\sin\psi}{|z|} > \frac{\delta\sin\psi_0}{2}$$

and any $c \leq \frac{1}{4}\sin\psi_0$ will suffice. So we may assume

$$\sin\theta \leq 2c\delta \text{ and } 1 - |z|\cos\theta \geq 0.$$

Now $\cos^2\theta \geq 1 - 4c^2\delta^2$, whence $\cos\theta = 1 - o(\delta)$, and we need only prove that $1 - |z| \geq 0$ and $1 - |z| \geq \Omega(\delta)$. To this end, observe that $\delta^2 = |1-z|^2 = (1 - |z|\cos\theta)^2 + (|z|\sin\theta)^2$, so that

$$\begin{aligned} 1 = |z|\cos\theta + (1 - |z|\cos\theta) &\geq |z|\sqrt{1 - 4c^2\delta^2} + \delta\sqrt{1 - 4c^2|z|^2} \\ &\geq |z|(1 - 4c^2\delta^2) + \delta(1 - 4c^2|z|^2), \end{aligned}$$

and hence, for all sufficiently small δ,

$$\begin{aligned} 1 - |z| &\geq \delta(1 - 4c^2|z|^2) - 4|z|c^2\delta^2 \\ &\geq \delta(1 - 16c^2) - 8c^2\delta^2 \quad (\text{since } |z| \leq 2 \text{ for } \delta \leq 1) \\ &\geq p\delta, \end{aligned}$$

where p is a positive constant depending only on ψ_0, since $c \leq \frac{1}{4}\sin\psi_0 < \frac{1}{4}$. □

Next we show how to modify Proposition 4.3.2 whenever $a + b > -2$.

Proposition 4.3.4. *Assume that* $f(z)$ *and* $g(z)$ *are* Δ*-regular and that*

$$f(z) = O(|1-z|^a) \text{ and } g(z) = O(|1-z|^b) \text{ as } z \to 1.$$

(a) If $k < a + b + 1 < k + 1$ *for some integer* $-1 \leq k < \infty$*, then*

$$(f \odot g)(z) = \sum_{j=0}^{k} \frac{(-1)^j}{j!}(f \odot g)^{(j)}(1)(1-z)^j + O(|1-z|^{a+b+1}).$$

(b) If $a + b + 1$ *is a nonnegative integer then*

$$(f \odot g)(z) = \sum_{j=0}^{a+b} \frac{(-1)^j}{j!}(f \odot g)^{(j)}(1)(1-z)^j + O(|1-z|^{a+b+1}|\log((1-z)^{-1})|).$$

Proof. Let D denote the operator $\frac{d}{dz}$ and let θ denote the operator zD.

(a) We first show that

$$(f \odot g)(z) = \sum_{j=0}^{k} \frac{(-1)^j}{j!} \left[\theta^j (f \odot g)\right](1)(-\ln z)^j + O(|1-z|^{a+b+1}). \tag{4.3.7}$$

The proof is by induction on k. The basis case of $k = -1$ is a special case of Proposition 4.3.2. Observe that

$$\theta(f \odot g) = (\theta f) \odot g$$

and by Theorem 4.2.1 that

$$(\theta f)(z) = zf'(z) = O(|1-z|^{a-1}).$$

If $k \geq 0$, then by the induction hypothesis

$$\left[\theta(f \odot g)\right](\zeta) = (\theta f \odot g)(\zeta) = \sum_{j=0}^{k-1} \frac{(-1)^j}{j!} \left[\theta^j (\theta f \odot g)\right](1)(-\ln \zeta)^j + O(|1-\zeta|^{a+b}).$$

We divide by ζ and then integrate both sides with respect to ζ from z to 1 to get

$$\begin{aligned}(f \odot g)(z) &= (f \odot g)(1) + \sum_{j=0}^{k-1} \frac{(-1)^{j+1}}{(j+1)!} \left[\theta^{j+1} (f \odot g)\right](1)(-\ln z)^{j+1} + O(|1-z|^{a+b+1}) \\ &= \sum_{j=0}^{k} \frac{(-1)^j}{j!} \left[\theta^j (f \odot g)\right](1)(-\ln z)^j + O(|1-z|^{a+b+1}),\end{aligned}$$

as required.

To obtain the expansion in the asserted form, we use the following two well-known facts:

a) For fixed nonnegative integers j and k, as $z \to 1$ we have

$$(-\ln z)^j = j! \sum_{r=0}^{k} \begin{bmatrix} r \\ j \end{bmatrix} \frac{(1-z)^r}{r!} + O(|1-z|^{k+1}),$$

where $\begin{bmatrix} r \\ j \end{bmatrix}$ denotes a Stirling number of the first kind (see 1.2.9-(26) in [43]).

b) The operators D and θ satisfy

$$z^r D^r = \sum_{j=1}^{r} (-1)^{r-j} \begin{bmatrix} r \\ j \end{bmatrix} \theta^j, \quad r \geq 1.$$

Using these facts in (4.3.7) and recalling that $a+b+1<k+1$, we get

$$\begin{aligned}
(f\odot g)(z) &= \sum_{j=0}^{k}(-1)^j\left[\theta^j(f\odot g)\right](1)\sum_{r=0}^{k}\begin{bmatrix} r\\ j\end{bmatrix}\frac{(1-z)^r}{r!}+O(|1-z|^{a+b+1})\\
&= (f\odot g)(1)+\sum_{r=1}^{k}(-1)^r\frac{(1-z)^r}{r!}\sum_{j=1}^{k}(-1)^{r-j}\begin{bmatrix} r\\ j\end{bmatrix}\left[\theta^j(f\odot g)\right](1)+O(|1-z|^{a+b+1})\\
&= \sum_{r=0}^{k}(-1)^r\frac{(1-z)^r}{r!}\left[w^r(f\odot g)^{(r)}(w)\right]\Big|_{w=1}(1)+O(|1-z|^{a+b+1})\\
&= \sum_{j=0}^{k}\frac{(-1)^j}{j!}(f\odot g)^{(j)}(1)(1-z)^j+O(|1-z|^{a+b+1}),
\end{aligned}$$

as desired.

(b) The structure of the proof is the same as that of part (a). One first proves a result similar to (4.3.7). Again, the proof is by induction on $k := a+b+1$. When $k=0$, after applying Theorem 4.2.1 and Proposition 4.3.2 to obtain $\theta(f\odot g)(z)=O(|1-z|^{-1})$ we can apply Theorem 4.2.2 to get $(f\odot g)(z)=O(|\log((1-z)^{-1})|)$. The induction step is similar to that in (a) above. Finally, using the same argument as for part (a), we obtain the expansion in the asserted form.

□

For convenience, we now combine Propositions 4.3.2 and 4.3.4 to get a result covering all possible values of a and b.

Theorem 4.3.5. *If f and g are amenable to singularity analysis and*

$$f(z)=O(|1-z|^a)\quad\text{and}\quad g(z)=O(|1-z|^b)\quad\text{as } z\to 1,$$

then $f\odot g$ is also amenable to singularity analysis. Furthermore,

(a) If $a+b+1<0$ then

$$f(z)\odot g(z)=O(|1-z|^{a+b+1}).$$

(b) If $k<a+b+1<k+1$ for some integer $-1\le k<\infty$, then

$$f(z)\odot g(z)=\sum_{j=0}^{k}\frac{(-1)^j}{j!}(f\odot g)^{(j)}(1)(1-z)^j+O(|1-z|^{a+b+1}).$$

(c) If $a+b+1$ is a nonnegative integer then

$$f(z)\odot g(z)=\sum_{j=0}^{a+b}\frac{(-1)^j}{j!}(f\odot g)^{(j)}(1)(1-z)^j+O(|1-z|^{a+b+1}|\log((1-z)^{-1})|).$$

We will make extensive use of Theorem 4.3.5 in the subsequent chapters. We mention in passing that the results of this chapter are also applicable to the analysis of the Cayley tree recurrence [45, 46], which models the evolution of a graph from being totally disconnected to being tree-like, when successive edges are added at random; see [23] for details.

Part III

Applications

Chapter 5

Random permutation model: shape functional

As our first application of the theoretical results of Chapters 3 and 4, we demonstrate how to get a complete asymptotic expansion for the mean of the shape functional (see Example 2.2.3) for m-ary search trees and we determine the asymptotics of its variance for binary search trees. Our results generalize those obtained in [21].

5.1 Complete asymptotic expansion of the mean

As noted in Example 2.2.3, the shape functional corresponds to the toll function $t_n := \ln \binom{n}{m-1}$. In order to obtain a complete asymptotic expansion of its mean using the ETT (Theorem 3.1.1), our first task is to obtain a complete asymptotic expansion for its generating function

$$T(z) := \sum_{n=m-1}^{\infty} t_n z^n.$$

The generating function of the toll sequence

By definition

$$\begin{aligned} T(z) &= \sum_{n=m-1}^{\infty} t_n z^n = \sum_{n=m-1}^{\infty} \left[\ln \binom{n}{m-1}\right] z^n \\ &= \sum_{n=m-1}^{\infty} \left[\sum_{l=0}^{m-2} \ln (n-l)\right] z^n - [\ln (m-1)!] \sum_{n=m-1}^{\infty} z^n = P(z) - Q(z), \end{aligned}$$

where

$$P(z) := \sum_{n=m-1}^{\infty} \left[\sum_{l=0}^{m-2} \ln (n-l)\right] z^n$$

and

$$Q(z) := [\ln (m-1)!] \sum_{n=m-1}^{\infty} z^n = [\ln (m-1)!] \left[(1-z)^{-1} - \sum_{n=0}^{m-2} z^n\right].$$

But

$$\begin{aligned} P(z) = \sum_{l=0}^{m-2} \sum_{n=m-1}^{\infty} [\ln (n-l)]\, z^n &= \sum_{l=0}^{m-2} z^l \sum_{n=l+1}^{\infty} [\ln (n-l)]\, z^{n-l} - \sum_{l=0}^{m-2} \sum_{n=l+1}^{m-2} [\ln (n-l)]\, z^n \\ &= \frac{1-z^{m-1}}{1-z} \operatorname{Li}_{0,1}(z) - \sum_{n=1}^{m-2} z^n \sum_{l=0}^{n-1} \ln (n-l), \end{aligned} \tag{5.1.1}$$

where Li denotes the generalized polylogarithm defined at (4.1.3). The term subtracted in (5.1.1) equals

$$\begin{aligned}\sum_{n=1}^{m-2} z^n \ln n! &= \sum_{n=1}^{m-2}\sum_{k=0}^{n}(-1)^k\binom{n}{k}(1-z)^k \ln n! \\ &= \sum_{k=0}^{m-2}(-1)^k(1-z)^k \sum_{n=\max\{k,1\}}^{m-2}\binom{n}{k}\ln n! = \sum_{k=0}^{m-2}(-1)^k(1-z)^k\sum_{n=k}^{m-2}\binom{n}{k}\ln n!, \qquad (5.1.2)\end{aligned}$$

while the first term in (5.1.1) is the product of $(1-z)^{-1}\,\mathrm{Li}_{0,1}(z)$ and

$$1 - z^{m-1} = 1 - \sum_{k=0}^{m-1}(-1)^k\binom{m-1}{k}(1-z)^k = \sum_{k=1}^{m-1}(-1)^{k+1}\binom{m-1}{k}(1-z)^k,$$

i.e., it is $\mathrm{Li}_{0,1}(z)$ times

$$\frac{1-z^{m-1}}{1-z} = \sum_{k=0}^{m-2}(-1)^k\binom{m-1}{k+1}(1-z)^k. \qquad (5.1.3)$$

Using Theorem 4.1.3 we know that

$$\mathrm{Li}_{0,1}(z) \sim t^{-1}\ln t^{-1} + \Gamma'(1)t^{-1} - \sum_{j\geq 0}\frac{(-1)^j}{j!}\zeta'(-j)t^j \qquad (5.1.4)$$

as $z \to 1$ in a suitable indented crown, with $t := -\ln z$. We note that $\Gamma'(1) = -\gamma$, where $\gamma \doteq 0.5772$ is Euler's constant.

In order to carry out the integration required in the ETT, we need to express $T(z)$ in terms of powers of $(1-z)$ and $\ln((1-z)^{-1})$. This is our first goal. [For notational convenience we will define $L(z) := \ln((1-z)^{-1})$.] To this end, expression (5.1.2) can be written as

$$\sum_{k=0}^{m-2} S_1(k)(1-z)^k \quad \text{where} \quad S_1(k) := (-1)^k\sum_{n=k}^{m-2}\binom{n}{k}\ln n!. \qquad (5.1.5)$$

Expression (5.1.3) can be written as

$$\sum_{k=0}^{m-2} S_2(k)(1-z)^k, \quad \text{where} \quad S_2(k) := (-1)^k\binom{m-1}{k+1}. \qquad (5.1.6)$$

For (5.1.4), we will need the following two facts: First,

$$t^j = \sum_{r=j}^{\infty} T_1(r,j)(1-z)^r, \quad j \geq -1 \quad \text{where} \quad T_1(r,i) := \begin{cases}[z^r]L^i(z) & i \neq 0\\ \delta_{r,0} & i = 0.\end{cases} \qquad (5.1.7)$$

Next,

$$\ln t^{-1} = L(z) + \sum_{r=1}^{\infty} T_2(r)(1-z)^r, \quad \text{where} \quad T_2(r) := \sum_{i=1}^{r}\frac{(-1)^i}{i}T_3(r+i,i), \qquad (5.1.8)$$

with

$$T_3(r,i) := [w^r]\,[L(w)-w]^i\,. \qquad (5.1.9)$$

The final ingredient in our expression for $T(z)$ is

$$\begin{aligned}\sum_{n=0}^{m-2} z^n = \sum_{n=0}^{m-2}(1-(1-z))^n &= \sum_{n=0}^{m-2}\sum_{k=0}^{n}\binom{n}{k}(-1)^k(1-z)^k \\ &= \sum_{k=0}^{m-2}(1-z)^k(-1)^k\sum_{n=k}^{m-2}\binom{n}{k} = \sum_{k=0}^{m-2} S_3(k)(1-z)^k, \end{aligned} \tag{5.1.10}$$

where

$$S_3(k) := (-1)^k \sum_{n=k}^{m-2}\binom{n}{k}. \tag{5.1.11}$$

We now express $\mathrm{Li}_{0,1}(z)$ in terms of $(1-z)$:

$$\begin{aligned}\mathrm{Li}_{0,1}(z) \sim{}& \left(\sum_{r=-1}^{\infty} T_1(r,-1)(1-z)^r\right)\left(L(z)+\sum_{r=1}^{\infty} T_2(r)(1-z)^r\right) \\ &+\Gamma'(1)\sum_{r=-1}^{\infty} T_1(r,-1)(1-z)^r - \sum_{j\geq 0}\frac{(-1)^j}{j!}\zeta'(-j)\sum_{r=j}^{\infty} T_1(r,j)(1-z)^r \\ ={}& \sum_{r=-1}^{\infty} T_1(r,-1)(1-z)^r L(z) + \sum_{r=-1}^{\infty} T_1(r,-1)(1-z)^r \sum_{j=1}^{\infty} T_2(j)(1-z)^j \\ &+\Gamma'(1)\sum_{r=-1}^{\infty} T_1(r,-1)(1-z)^r + \sum_{r=0}^{\infty}(1-z)^r\sum_{j=0}^{r}\frac{(-1)^j}{j!}\zeta'(-j)T_1(r,j) \\ ={}& \sum_{r=-1}^{\infty} T_1(r,-1)(1-z)^r L(z) + \sum_{r=-1}^{\infty}\sum_{j=1}^{\infty} T_1(r,-1)T_2(j)(1-z)^{r+j} \\ &+\Gamma'(1)\sum_{r=-1}^{\infty} T_1(r,-1)(1-z)^r + \sum_{r=0}^{\infty} S_4(r)(1-z)^r\end{aligned}$$

where

$$S_4(r) := \sum_{j=0}^{r}\frac{(-1)^j}{j!}\zeta'(-j)T_1(r,j). \tag{5.1.12}$$

Hence

$$\begin{aligned}\mathrm{Li}_{0,1}(z) \sim{}& \sum_{r=-1}^{\infty} T_1(r,-1)(1-z)^r L(z) + \sum_{r=-1}^{\infty}\sum_{j=r+1}^{\infty} T_1(r,-1)T_2(j-r)(1-z)^j \\ &+\Gamma'(1)\sum_{r=-1}^{\infty} T_1(r,-1)(1-z)^r + \sum_{r=0}^{\infty} S_4(r)(1-z)^r \\ ={}& \sum_{r=-1}^{\infty} T_1(r,-1)(1-z)^r L(z) + \sum_{j=0}^{\infty}(1-z)^j\sum_{r=-1}^{j-1} T_1(r,-1)T_2(j-r) \\ &+\Gamma'(1)\sum_{r=-1}^{\infty} T_1(r,-1)(1-z)^r + \sum_{r=0}^{\infty} S_4(r)(1-z)^r \\ ={}& \sum_{r=-1}^{\infty} T_1(r,-1)(1-z)^r L(z) + \sum_{r=-1}^{\infty} S_5(r)(1-z)^r,\end{aligned}$$

where

$$S_5(r) := \begin{cases} \Gamma'(1)T_1(-1,1) & r=-1 \\ \Gamma'(1)T_1(r,-1)+S_4(r)+\displaystyle\sum_{j=-1}^{r-1} T_1(j,-1)T_2(r-j) & r \geq 0. \end{cases} \tag{5.1.13}$$

Now we multiply (5.1.3) and (5.1.4), using (5.1.6) and (5.1.12), to get

$$\begin{aligned} &\left(\sum_{k=0}^{m-2} S_2(k)(1-z)^k\right)\left(\sum_{r=-1}^{\infty} T_1(r,-1)(1-z)^r L(z) + \sum_{r=-1}^{\infty} S_5(r)(1-z)^r\right) \\ &= \sum_{k=0}^{m-2}\sum_{r=-1}^{\infty} S_2(k)T_1(r,-1)(1-z)^{k+r}L(z) + \sum_{k=0}^{m-2}\sum_{r=-1}^{\infty} S_2(k)S_5(r)(1-z)^{k+r} \\ &= \sum_{k=0}^{m-2}\sum_{r=k-1}^{\infty} S_2(k)T_1(r-k,-1)(1-z)^r L(z) + \sum_{k=0}^{m-2}\sum_{r=k-1}^{\infty} S_2(k)S_5(r-k)(1-z)^r \\ &= \sum_{r=-1}^{\infty}(1-z)^r L(z) \sum_{k=0}^{\min\{r+1,m-2\}} S_2(k)T_1(r-k,-1) \\ &\quad + \sum_{r=-1}^{\infty}(1-z)^r \sum_{k=0}^{\min\{r+1,m-2\}} S_2(k)S_5(r-k) \\ &= \sum_{r=-1}^{\infty} S_6(r)(1-z)^r L(z) + \sum_{r=-1}^{\infty} S_7(r)(1-z)^r \end{aligned}$$

where

$$S_6(r) := \sum_{k=0}^{\min\{r+1,m-2\}} S_2(k)T_1(r-k,-1) \quad \text{and} \quad S_7(r) := \sum_{k=0}^{\min\{r+1,m-2\}} S_2(k)S_5(r-k). \tag{5.1.14}$$

We are now ready to plug (5.1.14), (5.1.5), and (5.1.10) into the expression for $T(z)$. This gives us

$$\begin{aligned} T(z) &\sim \sum_{r=-1}^{\infty} S_6(r)(1-z)^r L(z) + \sum_{r=-1}^{\infty} S_7(r)(1-z)^r \\ &\quad - \sum_{k=0}^{m-2} S_1(k)(1-z)^k - [\ln{(m-1)!}]\left[(1-z)^{-1} - \sum_{k=0}^{m-2} S_3(k)(1-z)^k\right] \\ &= \sum_{r=-1}^{\infty} S_6(r)(1-z)^r L(z) + \sum_{r=-1}^{\infty} S_8(r)(1-z)^r, \end{aligned} \tag{5.1.15}$$

where

$$S_8(r) := \begin{cases} S_7(-1) - \ln{(m-1)!} & r=-1 \\ S_7(r) - S_1(r) + [\ln{(m-1)!}]S_3(r) & 0 \leq r \leq m-2 \\ S_7(r) & r \geq m-1. \end{cases} \tag{5.1.16}$$

Application of the ETT

We now apply the ETT (Theorem 3.1.1). For our toll function,

$$\widehat{T}(z) := T(z) - \sum_{j=0}^{m-1} t_n z^j = T(z).$$

Thus, the generating function of the mean of the shape functional, call it G_1, is given by

$$G_1(z) = T(z) + m! \sum_{j=1}^{m-1} \frac{(1-z)^{-\lambda_j}}{\psi'(\lambda_j)} \int_{\zeta=0}^{z} T(\zeta)(1-\zeta)^{\lambda_j - 1}\, d\zeta. \tag{5.1.17}$$

We rewrite (5.1.15) as

$$T(z) \sim \sum_{r=-2}^{\infty} S_9(r)(1-z)^{r+1} L(z) + \sum_{r=-2}^{\infty} S_{10}(r)(1-z)^{r+1}, \tag{5.1.18}$$

where

$$S_9(r) := S_6(r+1) \quad \text{and} \quad S_{10}(r) := S_8(r+1). \tag{5.1.19}$$

For a fixed $\lambda \in \mathbb{C}$, the terms $(1-z)^{\lambda+r}L(z)$ and $(1-z)^{\lambda+r}$ are both integrable (resp., neither integrable) at $z = 1$ iff $r \geq -\lceil \Re(\lambda) \rceil$ (resp., iff $r < -\lceil \Re(\lambda) \rceil$).

We therefore write

$$\begin{aligned}
&\int_{\zeta=0}^{z} T(\zeta)(1-\zeta)^{\lambda-1} d\zeta \\
&= \sum_{-2 \leq r < -\lceil \Re(\lambda) \rceil} \int_{\zeta=0}^{z} [S_9(r)L(\zeta) + S_{10}(r)](1-\zeta)^{\lambda+r}\, d\zeta && \text{(5.1.20a)} \\
&\quad + \int_{\zeta=0}^{z} \left[T(\zeta)(1-\zeta)^{\lambda-1} - \sum_{-2 \leq r < -\lceil \Re(\lambda) \rceil} [S_9(r)L(\zeta) + S_{10}(r)](1-\zeta)^{\lambda+r} \right] d\zeta. && \text{(5.1.20b)}
\end{aligned}$$

The integrand in (5.1.20b) is integrable at $\zeta = 1$ and is asymptotically equivalent to

$$\sum_{r \geq -\lceil \Re(\lambda) \rceil} [S_9(r)L(\zeta) + S_{10}(r)](1-\zeta)^{\lambda+r},$$

as $\zeta \to 1$ in the sector $-\pi + \epsilon < \arg(1-\zeta) < \pi - \epsilon$ for any $\epsilon > 0$.

For (5.1.20a), we have the following formulas, valid for any α with $\Re(\alpha) > -1$:

$$\int_0^z (1-\zeta)^{\alpha}\, d\zeta = -\frac{1}{\alpha+1}[(1-z)^{\alpha+1} - 1],$$

and

$$\int_0^z (1-\zeta)^{\alpha} L(\zeta)\, d\zeta = -\frac{1}{\alpha+1}(1-z)^{\alpha+1}L(z) - \frac{1}{(\alpha+1)^2}[(1-z)^{\alpha+1} - 1].$$

In the special case that m is odd and $\lambda = \lambda_{m-1} = -m$ and $r = m-1$, we also need the formulas

$$\int_{\zeta=0}^{z} (1-\zeta)^{-1}\, d\zeta = L(z)$$

and

$$\int_{\zeta=0}^{z}(1-\zeta)^{-1}L(\zeta)\,d\zeta=\frac{1}{2}L^2(z).$$

We now conclude that (5.1.20a) equals, except in the special case described above,

$$\sum_{-2\le r<-\lceil\Re(\lambda)\rceil}\Bigg[-\frac{S_9(r)}{\lambda+r+1}(1-z)^{\lambda+r+1}L(z)-\frac{S_9(r)}{(\lambda+r+1)^2}(1-z)^{\lambda+r+1}$$
$$+\frac{S_9(r)}{(\lambda+r+1)^2}+\frac{S_{10}(r)}{\lambda+r+1}-\frac{S_{10}(r)}{\lambda+r+1}(1-z)^{\lambda+r+1}\Bigg]. \qquad (5.1.21)$$

In the special case (m is odd, $\lambda=-m$, $r=m-1$), (5.1.20a) equals

$$\sum_{r=-2}^{m-2}\Bigg[-\frac{S_9(r)}{\lambda+r+1}(1-z)^{\lambda+r+1}L(z)-\frac{S_9(r)}{(\lambda+r+1)^2}(1-z)^{\lambda+r+1}+\frac{S_9(r)}{(\lambda+r+1)^2}$$
$$+\frac{S_{10}(r)}{\lambda+r+1}-\frac{S_{10}(r)}{\lambda+r+1}(1-z)^{\lambda+r+1}\Bigg]+\frac{1}{2}S_9(m-1)L^2(z)+S_{10}(m-1)L(z). \qquad (5.1.22)$$

In the case m is even,

$$\sum_{j=1}^{m-1}\frac{(1-z)^{-\lambda_j}}{\psi'(\lambda_j)}\times(5.1.20\text{a})\Bigg|_{\lambda=\lambda_j}$$
$$=\sum_{j=1}^{m-1}\frac{1}{\psi'(\lambda_j)}\sum_{-2\le r<-\lceil\Re(\lambda_j)\rceil}\Bigg[-\frac{S_9(r)}{\lambda_j+r+1}(1-z)^{r+1}L(z)$$
$$-\left(\frac{S_{10}(r)}{\lambda_j+r+1}+\frac{S_9(r)}{(\lambda_j+r+1)^2}\right)(1-z)^{r+1}+\frac{S_9(r)}{(\lambda_j+r+1)^2}+\frac{S_{10}(r)}{\lambda_j+r+1}\Bigg]$$
$$=\sum_{r=-1}^{m-1}\left[S_{11}(r)(1-z)^rL(z)+S_{12}(r)(1-z)^r\right]+S_{13}, \qquad (5.1.23)$$

where

$$S_{11}(r):=-\sum_{j:\Re(\lambda_j)\le -r}\frac{S_9(r-1)}{\psi'(\lambda_j)(\lambda_j+r)},$$

$$S_{12}(r):=-\sum_{j:\Re(\lambda_j)\le -r}\frac{1}{\psi'(\lambda_j)}\left(\frac{S_9(r-1)}{(\lambda_j+r)^2}+\frac{S_{10}(r-1)}{\lambda_j+r}\right),$$

and

$$S_{13}:=\sum_{r=-1}^{m-1}\sum_{\Re(\lambda_j)\le -r}\frac{1}{\psi'(\lambda_j)}\left[\frac{S_9(r-1)}{(\lambda_j+r)^2}+\frac{S_{10}(r-1)}{\lambda_j+r+1}\right]. \qquad (5.1.24)$$

If m is odd, we need to add

$$\frac{(1-z)^m}{\psi'(-m)}\left(\frac{1}{2}S_9(m-1)L^2(z)+S_{10}(m-1)L(z)\right)$$

to the right-hand-side of (5.1.23) to get the left-hand-size of (5.1.23).

For (5.1.20b), we define

$$J_\lambda(\zeta) := T(\zeta)(1-\zeta)^{\lambda-1} - \sum_{-2\le r<-\lceil\Re(\lambda)\rceil} [S_9(r)L(z) + S_{10}(r)](1-\zeta)^{\lambda+r} \tag{5.1.25}$$

so that (5.1.20b) equals $\bar{K}(\lambda) - \int_{\zeta=z}^1 J_\lambda(\zeta)\,d\zeta$, where

$$\bar{K}(\lambda) := \int_{\zeta=0}^1 J_\lambda(\zeta)\,d\zeta \in \mathbb{C}. \tag{5.1.26}$$

Now we make use of

$$\int_{\zeta=z}^1 J_\lambda(\zeta)\,d\zeta \sim \sum_{r\ge-\lceil\Re(\lambda)\rceil} \int_{\zeta=z}^1 \Big[S_9(r)L(\zeta) + S_{10}(r)\Big](1-\zeta)^{\lambda+r}\,d\zeta. \tag{5.1.27}$$

Again, exact integration is possible: for all $\alpha\in\mathbb{C}$ with $\Re(\alpha) > -1$,

$$\int_{\zeta=z}^1 (1-\zeta)^\alpha\,d\zeta = \frac{1}{\alpha+1}(1-z)^{\alpha+1},$$

$$\int_{\zeta=z}^1 (1-\zeta)^\alpha L(\zeta)\,d\zeta = \frac{1}{\alpha+1}(1-z)^{\alpha+1}L(z) + \frac{1}{(\alpha+1)^2}(1-z)^{\alpha+1}.$$

Therefore

$$\begin{aligned}\int_{\zeta=z}^1 J_\lambda(\zeta)\,d\zeta \sim \sum_{r\ge-\lceil\Re(\lambda)\rceil} \Big[&\frac{S_9(r)}{\lambda+r+1}(1-z)^{\lambda+r+1}L(z) \\ &+ \frac{S_9(r)}{(\lambda+r+1)^2}(1-z)^{\lambda+r+1} + \frac{S_{10}(r)}{\lambda+r+1}(1-z)^{\lambda+r+1}\Big].\end{aligned}$$

We can thus conclude

$$\begin{aligned}&\sum_{j=1}^{m-1} \frac{(1-z)^{-\lambda_j}}{\psi'(\lambda_j)} \times (5.1.20\text{b})\Big|_{\lambda=\lambda_j} \\ &\sim \sum_{j=1}^{m-1} \frac{\bar{K}(\lambda_j)}{\psi'(\lambda_j)}(1-z)^{-\lambda_j} - \sum_{j=1}^{m-1}\frac{1}{\psi'(\lambda_j)} \sum_{r\ge-\lceil\Re(\lambda)\rceil} \Big[\frac{S_9(r)}{\lambda_j+r+1}(1-z)^{r+1}L(z) \\ &\qquad + \Big(\frac{S_9(r)}{(\lambda_j+r+1)^2} + \frac{S_{10}(r)}{\lambda_j+r+1}\Big)(1-z)^{r+1}\Big] \\ &= \sum_{j=1}^{m-1} \frac{\bar{K}(\lambda_j)}{\psi'(\lambda_j)}(1-z)^{-\lambda_j} + \sum_{r\ge-1} [S_{14}(r)(1-z)^r L(z) + S_{15}(r)(1-z)^r]\end{aligned} \tag{5.1.28}$$

where

$$S_{14}(r) := -\sum_{j:\Re(\lambda_j)>-r} \frac{S_9(r-1)}{(\lambda_j+r)\psi'(\lambda_j)}$$

$$S_{15}(r) := -\sum_{j:\Re(\lambda_j)>-r} \frac{1}{\psi'(\lambda_j)}\Big[\frac{S_9(r-1)}{(\lambda_j+r)^2} + \frac{S_{10}(r-1)}{\lambda_j+r}\Big].$$

Case m even

Using (5.1.23), (5.1.28), and (5.1.15) in (5.1.17), we get

$$G_1(z) = \sum_{j=1}^{m-1} \frac{K(\lambda_j)}{\psi'(\lambda_j)}(1-z)^{-\lambda_j} + M + \sum_{r\geq -1}[K_r(1-z)^r L(z) + L_r(1-z)^r], \tag{5.1.29}$$

where

$$K(\lambda) := m!\bar{K}(\lambda) \quad \text{and} \quad M := m!S_{13}, \tag{5.1.30}$$

$$\begin{aligned} K_r &:= m![S_{11}(r) + S_{14}(r)] + S_6(r) = -m!S_9(r-1)\sum_{j=1}^{m-1}\frac{1}{(\lambda_j + r)\psi'(\lambda_j)} + S_6(r), \\ &= \frac{m!S_9(r-1)}{(-r)^{\overline{m-1}} - m!} + S_6(r), \end{aligned} \tag{5.1.31}$$

and

$$\begin{aligned} L_r &:= m![S_{12}(r) + S_{15}(r)] + S_8(r) \\ &= -m!\left[S_9(r-1)\sum_{j=1}^{m-1}\frac{1}{(\lambda_j+r)^2\psi'(\lambda_j)} + S_{10}(r-1)\sum_{j=1}^{m-1}\frac{1}{(\lambda_j+r)\psi'(\lambda_j)}\right] + S_8(r) \\ &= -m!\left[S_9(r-1)\sum_{j=1}^{m-1}\frac{1}{(\lambda_j+r)^2\psi'(\lambda_j)} - \frac{S_{10}(r-1)}{(-r)^{\overline{m-1}} - m!}\right] + S_8(r). \end{aligned} \tag{5.1.32}$$

These constants can be obtained in a closed form using Identities B.1 and B.2.

Case m odd

In this case we have the same result as (5.1.29) except that the $r = m$ term must be replaced by

$$\begin{aligned} m!\Bigg\{ &\frac{(1-z)^m}{\psi'(-m)}\left[\frac{1}{2}S_9(m-1)L^2(z) + S_{10}(m-1)L(z)\right] \\ &+ S_{14}(m)(1-z)^m L(z) + S_{15}(m)(1-z)^m\Bigg\} + S_6(m)(1-z)^m L(z) + S_8(m)(1-z)^m \\ &\qquad = N_m(1-z)^m L^2(z) + K_m(1-z)^m L(z) + L_m(1-z)^m, \end{aligned}$$

where

$$N_m := \frac{m!S_9(m-1)}{2\psi'(-m)} = -\frac{S_9(m-1)}{2(H_m - 1)}, \quad K_m := m!\Big[\frac{S_{10}(m)}{\psi'(-m)} + S_{14}(m)\Big] + S_6(m),$$

and

$$L_m := m!S_{15}(m) + S_8(m).$$

(We note that K_m and L_m defined here for m odd are different [compare (5.1.31) and (5.1.32)] from those defined for m even.) The constants can be simplified using Identities B.5 and B.6.

Indeed,

$$\begin{aligned}S_{14}(m) &= -\sum_{j:\Re(\lambda_j)>-m} \frac{S_9(m-1)}{(\lambda_j+m)\psi'(\lambda_j)} = -S_9(m-1)\sum_{j\neq m-1}\frac{1}{(\lambda_j+m)\psi'(\lambda_j)}\\ &= -\frac{S_9(m-1)}{2m!}\Big[1-\frac{H_m^{(2)}-1}{(H_m-1)^2}\Big],\end{aligned}$$

and

$$\begin{aligned}S_{15}(m) &= -S_9(m-1)\sum_{j\neq m-1}\frac{1}{(\lambda_j+m)^2\psi'(\lambda_j)} - S_{10}(m-1)\sum_{j\neq m-1}\frac{1}{(\lambda_j+m)\psi'(\lambda_j)}\\ &= -\frac{S_9(m-1)}{m!}\left[\frac{H_m-1}{12}-\frac{1}{3}\frac{H_m^{(3)}-1}{(H_m-1)^2}+\frac{1}{4}\frac{(H_m^{(2)}-1)^2}{(H_m-1)^3}\right]\\ &\quad -\frac{S_{10}(m-1)}{2\cdot m!}\left[1-\frac{H_m^{(2)}-1}{(H_m-1)^2}\right].\end{aligned}$$

Some observations

The following facts can be established for the constants.

Remark 5.1.1. $K_r = 0$ for $0 \le r \le m-2$.

Proof.

$$K_r = \frac{m!S_9(r-1)}{(-r)^{\overline{m-1}}-m!}+S_6(r) = \frac{m!S_6(r)}{(-r)^{\overline{m-1}}-m!}+S_6(r) = S_6(r)\frac{(-r)^{\overline{m-1}}}{(-r)^{\overline{m-1}}-m!},$$

which contains the factor 0 if $0 \le r \le m-2$. □

Remark 5.1.2. $K_{-1} = -1$.

Proof.

$$\begin{aligned}K_{-1} &= S_6(-1)\frac{(m-1)!}{(m-1)!-m!} = \frac{S_6(-1)}{1-m} = \frac{(m-1)T_1(-1,-1)}{1-m}\\ &= -T_1(-1,-1) = -[z^{-1}]L(z) = -1.\end{aligned}$$

□

Remark 5.1.3. $K_{m-1} = 0$ for all odd values of m.

Proof. When m is odd,

$$K_{m-1} = \frac{S_6(m-1)(-1)^{m-1}(m-1)!}{(-1)^{m-1}(m-1)!-m!} = -\frac{S_6(m-1)}{m-1}$$

Now $S_6(m-1)$ is the coefficient of $(1-z)^{m-1}L(z)$ in the product (5.1.3) × (5.1.4), i.e.,

$$\begin{aligned}
S_6(m-1) &= [(1-z)^{m-1}L(z)]\,((5.1.3)\times(5.1.4))\\
&= [(1-z)^{m-1}L(z)]\left(\frac{1-z^{m-1}}{1-z}(-\ln z)^{-1}L(z)\right)\\
&= [(1-z)^{m-1}]\left(\frac{1-z^{m-1}}{1-z}(-\ln z)^{-1}\right)\\
&= [z^{m-1}]\left(\frac{1-(1-z)^{m-1}}{zL(z)}\right) = [z^m]\frac{1-(1-z)^{m-1}}{L(z)} = 0.
\end{aligned}$$

To see the last equality, observe that

$$\frac{1-(1-z)^{m-1}}{L(z)} = \int_{x=0}^{m-1}(1-z)^x\,dx$$

and

$$[z^m]\int_{x=0}^{m-1}(1-z)^x\,dx = [z^m]\int_{x=0}^{m-1}\sum_{n=0}^{\infty}\frac{(-x)^{\overline{n}}}{n!}z^n\,dx = \frac{1}{m!}\int_{x=0}^{m-1}(-x)^{\overline{m}}\,dx.$$

But the last integral vanishes because its integrand is antisymmetric about the midpoint of the interval of integration:

$$-(m-1-x)^{\overline{m}} = (-1)^m(m-1-x)_m = (-1)^m - x^{\overline{m}} = -(-x)^{\overline{m}},$$

the last equality following from the assumption that m is odd. □

Remark 5.1.4.

$$L_{-1} = \frac{\ln(m-1)!}{m-1} - \frac{m}{m-1}H_{m-1} + \gamma.$$

Proof.

$$\begin{aligned}
L_{-1} &= m!\Big[S_9(-2)\Big(-\frac{H_{m-1}}{(m-1)!(m-1)^2}\Big) + S_{10}(-2)\Big(\frac{1}{(m-1)!-m!}\Big)\Big] + S_8(-1)\\
&= m!\Big[-S_6(-1)\frac{H_{m-1}}{(m-1)!(m-1)^2} - S_8(-1)\Big(\frac{1}{(m-1)!(m-1)}\Big)\Big] + S_8(-1).
\end{aligned}$$

Now $S_6(-1) = m-1$ as noted in the proof of Remark 5.1.2, and $S_8(-1) = -(m-1)\gamma - \ln(m-1)!$. Plugging these in the display above, we get the desired result. □

The complete asymptotic expansion

The computations in this section have thus established that the generating function of the mean of the shape functional for m-ary search trees under the random permutation model, denoted by $G_1(z)$, has the complete asymptotic expansion

$$\begin{aligned}
G_1(z) \sim \sum_{j=1}^{m-1}\frac{K(\lambda_j)}{\psi'(\lambda_j)}(1-z)^{-\lambda_j} + M - (1-z)^{-1}L(z) + L_{-1}(1-z)^{-1}&\\
+ \sum_{r=0}^{m-2}L_r(1-z)^r + \sum_{r\geq m-1}\left[K_r(1-z)^rL(z) + L_r(1-z)^r\right]& \qquad (5.1.33)
\end{aligned}$$

when m is even and

$$G_1(z) \sim \sum_{j=1}^{m-1} \frac{K(\lambda_j)}{\psi'(\lambda_j)}(1-z)^{-\lambda_j} + M - (1-z)^{-1}L(z) + L_{-1}(1-z)^{-1} + \sum_{r=0}^{m-1} L_r(1-z)^r$$
$$+ N_m(1-z)^m L^2(z) + K_m(1-z)^m L(z) + L_m(1-z)^m + \sum_{r\geq m+1} [K_r(1-z)^r L(z) + L_r(1-z)^r]$$

when m is odd.

Using singularity analysis (Theorem 4.1.2), the complete asymptotic expansion can be translated on a term-by-term basis to a complete asymptotic expansion of the mean itself, denoted by μ_n. Thus

$$\mu_n \sim \sum_{j=1}^{m-1} \frac{K(\lambda_j)}{\psi'(\lambda_j)} \frac{\lambda_j^{\overline{n}}}{n!} - H_n + L_{-1} + \sum_{r\geq m-1} K_r \Delta^r\left(\left(\tfrac{1}{n}\right)\right) \tag{5.1.34}$$

when m is even and

$$\mu_n \sim \sum_{j=1}^{m-1} \frac{K(\lambda_j)}{\psi'(\lambda_j)} \frac{\lambda_j^{\overline{n}}}{n!} - H_n + L_{-1} + N_m \Delta^m\left(\left(\tfrac{2H_{n-1}}{n}\right)\right) + K_m \Delta^m\left(\left(\tfrac{1}{n}\right)\right) + \sum_{r\geq m+1} K_r \Delta^r\left(\left(\tfrac{1}{n}\right)\right)$$

when m is odd, where $\Delta^r(s)$ denotes the rth-order difference of a sequence $s = (s_n)$. Note that $\Delta^r((1/n)) = \Theta(n^{-(r+1)})$ and $\Delta^m((2H_{n-1}/n)) = \Theta(n^{-(m+1)}\log n)$.

The binary search tree

In preparation for the computation of the variance of the shape functional for the binary search trees we state a special case of the complete asymptotic expansion derived above. When $m = 2$, the toll function t_n equals $\ln n$ and the indicial polynomial $\psi(\lambda)$ equals $\lambda - 2$, with root $\lambda_1 = 2$. Thus (5.1.34), Remark 5.1.4, and the well-known expansion for the harmonic numbers [43, 1.2.7-(3)] immediately yield

$$\mu_n = K(2)\frac{2^{\overline{n}}}{n!} - H_n - 2 + \gamma + O(n^{-2}) = K(2)(n+1) - \ln n - 2 - \frac{1}{2n} + O(n^{-2}), \tag{5.1.35}$$

where, by (5.1.30), (5.1.26), and (5.1.25),

$$K(2) = 2\bar{K}(2) = 2\int_{\zeta=0}^{1} T(\zeta)(1-\zeta)\,d\zeta = 2\int_{\zeta=0}^{1} \sum_{k=1}^{\infty} (\ln k)\zeta^k(1-\zeta)\,d\zeta = 2\sum_{k=1}^{\infty} \frac{\ln k}{(k+1)(k+2)}.$$

5.2 The variance for binary search trees

Let X_n denote the mean of the shape functional for binary search trees under the random permutation model. In the sequel it will be convenient to work with the approximately centered random variable $\widetilde{X}_n$ defined as

$$\widetilde{X}_n := X_n - C_1(n+1) \quad \text{for } n \geq 0,$$

where $C_1 := K(2)$, and to define

$$\tilde{\mu}_n(k) := \mathbf{E}\,\widetilde{X}_n^k \quad \text{for } k \geq 1,$$

with $\tilde{\mu}_n := 0$. Then, following the arguments leading to (2.3.1) and (2.3.2), we have

$$\tilde{\mu}_n(2) = \frac{2}{n}\sum_{j=0}^{n-1}\tilde{\mu}_j(2) + r_n(2),$$

where

$$\begin{aligned}
r_n(2) &:= \sum_{\substack{k_1,k_2,k_3\geq 0\\ k_1+k_2+k_3=2\\ k_1<2,\ k_2<2}} \binom{2}{k_1,k_2,k_3}(\ln n)^{k_3}\frac{1}{n}\sum_{\substack{j_1+j_2=n-1\\ j_1,j_2\geq 0}}\tilde{\mu}_{j_1}(k_1)\tilde{\mu}_{j_2}(k_2)\\
&= (\ln n)^2 + 4(\ln n)\frac{1}{n}\sum_{j=0}^{n-1}\tilde{\mu}_j(1) + \frac{2}{n}\sum_{j=0}^{n-1}\tilde{\mu}_j(1)\tilde{\mu}_{n-1-j}(1)\\
&= (\ln n)^2 + 4(\ln n)\frac{1}{n}\sum_{j=0}^{n-1}(\mu_j - C_1(j+1))\\
&\quad + \frac{2}{n}\sum_{j=0}^{n-1}(\mu_j - C_1(j+1))(\mu_{n-1-j} - C_1(n-j)),
\end{aligned}$$

and $r_0(2) = \mathbf{E}\,\widetilde{X}_0^2 = C_1^2$. Note that knowing $\tilde{\mu}_n(1)$ and $\tilde{\mu}_n(2)$ is enough to compute the variance of the random variable X_n, call it σ_n^2, since

$$\sigma_n^2 = \mathbf{Var}\,\widetilde{X}_n = \tilde{\mu}_n(2) - \tilde{\mu}_n(1)^2. \tag{5.2.1}$$

Our proposal is to use the ETT (Theorem 3.1.1) again. To that end, let $B(z)$ be the generating function of $r_n(2)$. Then using the expression for $r_n(2)$ above we get

$$\begin{aligned}
B(z) = C_1^2 + \mathrm{Li}_{0,2}(z) + 4\,\mathrm{Li}_{0,1}(z) \odot \Big[\int_0^z (1-\zeta)^{-1}\big(G_1(\zeta) - C_1(1-\zeta)^{-2}\big)\,d\zeta\Big]&\\
+ 2\int_0^z [G_1(\zeta) - C_1(1-\zeta)^{-2}]^2\,d\zeta.&
\end{aligned}\tag{5.2.2}$$

Our goal is to get each term in this sum to $O(|1-z|^{-\epsilon})$ for arbitrarily small $\epsilon > 0$.

Using Theorem 4.1.3 we get

$$\mathrm{Li}_{\alpha,0}(z) \sim \Gamma(1-\alpha)t^{\alpha-1} + \sum_{j\geq 0}\frac{(-1)^j}{j!}\zeta(\alpha-j)t^j,$$

$$\mathrm{Li}_{\alpha,1}(z) \sim \Gamma'(1-\alpha)t^{\alpha-1} - \Gamma(1-\alpha)t^{\alpha-1}\ln t - \sum_{j\geq 0}\frac{(-1)^j}{j!}\zeta'(\alpha-j)t^j,$$

and

$$\begin{aligned}
\mathrm{Li}_{\alpha,2}(z) \sim \Gamma''(1-\alpha)t^{\alpha-1} - 2\Gamma'(1-\alpha)t^{\alpha-1}\ln t&\\
+ \Gamma(1-\alpha)t^{\alpha-1}(\ln t)^2 + \sum_{j\geq 0}\frac{(-1)^j}{j!}\zeta''(\alpha-j)t^j,&
\end{aligned}$$

where

$$t = -\ln z = (1-z) + \frac{(1-z)^2}{2} + O(|1-z|^3).$$

Hence

$$t^{-1} = (1-z)^{-1} - \frac{1}{2} + O(|1-z|)$$

and [recalling $L(z) := \ln((1-z)^{-1})$]

$$\ln t = -L(z) + \frac{1}{2}(1-z) + O(|1-z|^2).$$

This gives

$$(\ln t)^2 = L^2(z) - (1-z)L(z) + O(|(1-z)^2 L(z)|),$$

so that

$$t^{-1}(\ln t)^2 = (1-z)^{-1}L^2(z) - \frac{1}{2}L^2(z) - L(z) + O(|(1-z)L(z)|)$$

and

$$t^{-1}\ln t = -(1-z)^{-1}L(z) + \frac{1}{2}L(z) + O(1).$$

Combining these expansions we get

$$\mathrm{Li}_{0,1}(z) = (1-z)^{-1}L(z) + \Gamma'(1)(1-z)^{-1} - \frac{1}{2}L(z) + O(1) \tag{5.2.3}$$

and

$$\begin{aligned}\mathrm{Li}_{0,2}(z) &= (1-z)^{-1}L^2(z) - \frac{1}{2}L^2(z) - L(z) \\ &\quad - 2\Gamma'(1)\big[-(1-z)^{-1}L(z) + \frac{1}{2}L(z)\big] + \Gamma''(1)(1-z)^{-1} + O(1) \\ &= (1-z)^{-1}L^2(z) + 2\Gamma'(1)(1-z)^{-1}L(z) \\ &\quad + \Gamma''(1)(1-z)^{-1} - \frac{1}{2}L^2(z) - (\Gamma'(1)+1)L(z) + O(1).\end{aligned} \tag{5.2.4}$$

We also need an asymptotic expansion for the generating function $G_1(z) - C_1(1-z)^{-2}$ of $\tilde{\mu}_n(1)$. This is available to us using (5.1.33). Indeed, using $M = 0$ from (5.1.30), $L_0 = 1/4$ from (5.1.32), $L_{-1} = -(2-\gamma)$ from Remark 5.1.4, and $K_1 = -1/36$ from (5.1.31) [and the definitions (5.1.19), (5.1.14), (5.1.6), (5.1.7), (5.1.24)], we get

$$G_1(z) - C_1(1-z)^{-2} = -(1-z)^{-1}L(z) + L_{-1}(1-z)^{-1} + \frac{1}{4} - \frac{1}{36}(1-z)L(z) + O(|1-z|).$$

Thus, invoking Theorem 4.2.2, we get

$$\begin{aligned}&\int_0^z (1-\zeta)^{-1}[G_1(\zeta) - C_1(1-\zeta)^{-2}]\,d\zeta \\ &= \int_0^z [-(1-\zeta)^{-2}L(\zeta) + L_{-1}(1-\zeta)^{-2} + \frac{1}{4}(1-\zeta)^{-1} - \frac{1}{36}L(\zeta) + O(1)]\,d\zeta \\ &= -[(1-z)^{-1}L(z) - (1-z)^{-1}] + L_{-1}(1-z)^{-1} + \frac{1}{4}L(z) + O(1) \\ &= -(1-z)^{-1}L(z) + (L_{-1}+1)(1-z)^{-1} + \frac{1}{4}L(z) + O(1),\end{aligned}$$

and

$$[G_1(z) - C_1(1-z)^{-2}]^2 = (1-z)^{-2}L^2(z) - 2L_{-1}(1-z)^{-2}L(z) + L_{-1}^2(1-z)^{-2} - \frac{1}{2}(1-z)^{-1}L(z) + \frac{1}{2}L_{-1}(1-z)^{-1} + O(|L^2(z)|). \quad (5.2.5)$$

We now integrate (5.2.5) term by term. Integration by parts establishes

$$\int_0^z (1-\zeta)^r L^2(\zeta)\,d\zeta = \begin{cases} -\dfrac{1}{r+1}(1-z)^{r+1}L^2(z) - \dfrac{2}{(r+1)^2}(1-z)^{r+1}L(z) - \dfrac{2}{(r+1)^3}[(1-z)^{r+1}-1] & (r \neq -1) \\ \frac{1}{3}L^3(z) & (r=-1) \end{cases}$$

and

$$\int_0^z (1-\zeta)^r L(\zeta)\,d\zeta = \begin{cases} -\dfrac{1}{r+1}(1-z)^{r+1}L(z) - \dfrac{1}{(r+1)^2}(1-z)^{r+1} + \dfrac{1}{(r+1)^2} & (r \neq -1) \\ \dfrac{1}{2}L^2(z) & (r=-1). \end{cases}$$

By Theorem 4.2.2 the integral of the remainder in (5.2.5) is $O(1) = O(|1-z|^{-\epsilon})$. Hence

$$\begin{aligned} &\int_0^z [G_1(\zeta) - C_1(1-\zeta)^{-2}]^2\,d\zeta \\ &= (1-z)^{-1}L^2(z) - 2(1-z)^{-1}L(z) + 2(1-z)^{-1} \\ &\quad - 2L_{-1}[(1-z)^{-1}L(z) - (1-z)^{-1}] + L_{-1}^2(1-z)^{-1} + O(|1-z|^{-\epsilon}) \\ &= (1-z)^{-1}L^2(z) - 2(L_{-1}+1)(1-z)^{-1}L(z) + (2 + 2L_{-1} + L_{-1}^2)(1-z)^{-1} + O(|1-z|^{-\epsilon}). \end{aligned} \quad (5.2.6)$$

We now have an asymptotic expansion for each term in the sum (5.2.2) but the one involving the Hadamard product. We make use of the following facts.

1. $(1-z)^{-1}$ is the identity element for Hadamard products, i.e., $(1-z)^{-1} \odot f(z) = f(z)$.
2. $L(z) = O(|1-z|^{-\epsilon})$ for any $\epsilon > 0$, so that $(1-z)^{-1}L(z) = O(|1-z|^{-1-\epsilon})$. Hence, by Theorem 4.3.5, $[(1-z)^{-1}L(z)] \odot O(1) = O(|1-z|^{-\epsilon})$.
3. Similarly,

$$[(1-z)^{-1}L(z)] \odot L(z) = O(|1-z|^{-\epsilon}), \qquad L(z) \odot L(z) = O(1),$$

$$L(z) \odot O(1) = O(1), \qquad O(1) \odot O(1) = O(1).$$

4. The following lemma will also be used.

Lemma 5.2.1. $[(1-z)^{-1}L(z)] \odot [(1-z)^{-1}L(z)] = (1-z)^{-1}L^2(z) + (1-z)^{-1}\operatorname{Li}_{2,0}(z)$.

Proof. Let H_n denote the nth harmonic number. Then

$$\begin{aligned}
&[(1-z)^{-1}L(z)]\odot[(1-z)^{-1}L(z)] = \sum_{n\geq 1} H_n^2 z^n = \sum_{n\geq 1}\left(\sum_{k=1}^{n}\frac{1}{k}\right)^2 z^n\\
&= \sum_{n\geq 1}\left(\sum_{k=1}^{n}\frac{1}{k^2}\right) z^n + 2\sum_{n\geq 1}\left[\sum_{k=2}^{n}\frac{1}{k}\sum_{j=1}^{k-1}\frac{1}{j}\right] z^n = \sum_{k\geq 1}\frac{1}{k^2}\left(\sum_{n\geq k} z^n\right) + 2\sum_{n\geq 1}\left[\sum_{k=2}^{n}\frac{H_{k-1}}{k}\right] z^n\\
&= \sum_{k\geq 1}\frac{z^k}{k^2}(1-z)^{-1} + 2\sum_{n\geq 1}\left[\sum_{k=2}^{n}\frac{H_{k-1}}{k}\right] z^n.
\end{aligned}$$

The first sum is $(1-z)^{-1}\operatorname{Li}_{2,0}(z)$. For the second sum, observe that

$$\begin{aligned}
\sum_{n\geq 1}\left[\sum_{k=2}^{n}\frac{H_{k-1}}{k}\right] z^n &= \sum_{k\geq 2}\frac{H_{k-1}}{k}\sum_{n\geq k} z^n = (1-z)^{-1}\sum_{k\geq 1}\frac{H_k}{k+1}z^{k+1}\\
&= (1-z)^{-1}\int_0^z (1-\zeta)^{-1}L(\zeta)\,d\zeta = \frac{1}{2}(1-z)^{-1}L^2(z),
\end{aligned}$$

and the claim follows. □

In light of these facts we can now conclude that

$$\begin{aligned}
&\operatorname{Li}_{0,1}(z)\odot\int_0^z (1-\zeta)^{-1}[G_1(\zeta)-C_1(1-\zeta)^{-2}]\,d\zeta \qquad (5.2.7)\\
&= [(1-z)^{-1}L(z)+\Gamma'(1)(1-z)^{-1}-\frac{1}{2}L(z)+O(1)]\\
&\qquad \odot[-(1-z)^{-1}L(z)+(L_{-1}+1)(1-z)^{-1}+\frac{1}{4}L(z)+O(1)]\\
&= -\big[(1-z)^{-1}L^2(z)+(1-z)^{-1}\operatorname{Li}_{2,0}(z)\big]+(L_{-1}+1)(1-z)^{-1}L(z)\\
&\qquad -\Gamma'(1)(1-z)^{-1}L(z)+\Gamma'(1)(L_{-1}+1)(1-z)^{-1}+O(|1-z|^{-\epsilon}).
\end{aligned}$$

To get an asymptotic expansion for $\operatorname{Li}_{2,0}$ we use (4.1.4):

$$\operatorname{Li}_{2,0}(z) = t(\ln t - H_1)+\frac{\pi^2}{6}+O(|1-z|^2) = \frac{\pi^2}{6}-(1-z)L(z)+O(|1-z|).$$

Hence we see that the Hadamard product (5.2.7) has the asymptotic expansion

$$\begin{aligned}
&-(1-z)^{-1}L^2(z)-(1-z)^{-1}\Big(\frac{\pi^2}{6}-(1-z)L(z)+O(|1-z|)\Big)\\
&\quad +(L_{-1}+1-\Gamma'(1))(1-z)^{-1}L(z)+\Gamma'(1)(L_{-1}+1)(1-z)^{-1}+O(|1-z|^{-\epsilon})\\
&= -(1-z)^{-1}L^2(z)+(L_{-1}+1-\Gamma'(1))(1-z)^{-1}L(z)\\
&\quad +\Big[\Gamma'(1)(L_{-1}+1)-\frac{\pi^2}{6}\Big](1-z)^{-1}+O(|1-z|^{-\epsilon}). \qquad (5.2.8)
\end{aligned}$$

Using (5.2.4), (5.2.6), and (5.2.8) in (5.2.2), we get

$$\begin{aligned}
&B(z)\\
&= C_1^2 + (1-z)^{-1}L^2(z) + 2\Gamma'(1)(1-z)^{-1}L(z) + \Gamma''(1)(1-z)^{-1}\\
&+ 4\Big[-(1-z)^{-1}L^2(z) + (L_{-1}+1-\Gamma'(1))(1-z)^{-1}L(z) + \big[\Gamma'(1)(L_{-1}+1) - \frac{\pi^2}{6}\big](1-z)^{-1}\Big]\\
&+ 2\Big[(1-z)^{-1}L^2(z) - 2(L_{-1}+1)(1-z)^{-1}L(z) + (2+2L_{-1}+L_{-1}^2)(1-z)^{-1}\Big] + O(|1-z|^{-\epsilon})\\
&= -(1-z)^{-1}L^2(z) + V_1(1-z)^{-1}L(z) + V_2(1-z)^{-1} + O(|1-z|^{-\epsilon}), \qquad (5.2.9)
\end{aligned}$$

where

$$V_1 := -2\Gamma'(1) = 2\gamma \qquad (5.2.10)$$

and

$$V_2 := \Gamma''(1) + 4\Big(\Gamma'(1)(L_{-1}+1) - \frac{\pi^2}{6}\Big) + 2(2+2L_{-1}+L_{-1}^2) = 4 - \gamma^2 - \frac{\pi^2}{2}. \qquad (5.2.11)$$

Let $\widehat{B}(z) := B(z) - r_0(2) = B(z) - C_1^2$. By direct application of the ETT, $A(z)$, defined to be the generating function of $\big(\tilde{\mu}_n(2)\big)$, is given by

$$A(z) = c_1(1-z)^{-2} + \widehat{B}(z) + 2(1-z)^{-2}\int_0^z \widehat{B}(\zeta)(1-\zeta)\,d\zeta,$$

where $c_1 = r_0(2) = C_1^2$. The integral here can be broken up as

$$\int_0^z \widehat{B}(\zeta)(1-\zeta)\,d\zeta = V - \int_z^1 \widehat{B}(\zeta)(1-\zeta)\,d\zeta,$$

where the first term on the right is the constant

$$V := \int_0^1 \widehat{B}(\zeta)(1-\zeta)\,d\zeta = \int_0^1 \left(\sum_{k=1}^{\infty} r_k(2)\zeta^k\right)(1-\zeta)\,d\zeta = \sum_{k=1}^{\infty} \frac{r_k(2)}{(k+1)(k+2)}.$$

For the second term we use the following consequences of integration by parts:

$$\int_z^1 L^2(\zeta)\,d\zeta = (1-z)L^2(z) + 2(1-z)L(z) + 2(1-z),$$

$$\int_z^1 L(\zeta)\,d\zeta = (1-z)L(z) + (1-z)$$

and the crude remainder estimate

$$\int_z^1 O(|1-\zeta|^{1-\epsilon})\,d\zeta = O(|1-z|^{2-\epsilon}).$$

Thus

$$\begin{aligned}
&(1-z)^{-2}\int_z^1 \widehat{B}(\zeta)(1-\zeta)\,d\zeta \\
&\quad = (1-z)^{-2}\bigl[-(1-z)L^2(z) - 2(1-z)L(z) - 2(1-z) + V_1(1-z)L(z) \\
&\qquad\qquad + V_1(1-z) + V_2(1-z) + O(|1-z|^{2-\epsilon})\bigr] \\
&\quad = -(1-z)^{-1}L^2(z) + (V_1-2)(1-z)^{-1}L(z) + (V_1+V_2-2)(1-z)^{-1} + O(|1-z|^{-\epsilon}),
\end{aligned}$$

and

$$\begin{aligned}
A(z) &= C_1^2(1-z)^{-2} - (1-z)^{-1}L^2(z) + V_1(1-z)^{-1}L(z) + V_2(1-z)^{-1} + O(|1-z|^{-\epsilon}) \\
&\quad + 2\Bigl[V(1-z)^{-2} + (1-z)^{-1}L^2(z) - (V_1-2)(1-z)^{-1}L(z) \\
&\quad - (V_1+V_2-2)(1-z)^{-1} + O(|1-z|^{-\epsilon})\Bigr] \\
&= (C_1^2+2V)(1-z)^{-2} + (1-z)^{-1}L^2(z) + (V_1 - 2(V_1-2))(1-z)^{-1}L(z) \\
&\quad + (V_2 - 2(V_1+V_2-2))(1-z)^{-1} + O(|1-z|^{-\epsilon}) \\
&= (C_1^2+2V)(1-z)^{-2} + (1-z)^{-1}L^2(z) + (4-V_1)(1-z)^{-1}L(z) \\
&\quad + (4-2V_1-V_2)(1-z)^{-1} + O(|1-z|^{-\epsilon}).
\end{aligned} \tag{5.2.12}$$

We can now transfer this asymptotic expansion for the generating function to one for its Taylor coefficients, invoking Theorem 4.1.2 for the remainder estimate. We know that

$$[z^n](1-z)^{-1}L(z) = H_n = \ln n + \gamma + O(n^{-1})$$

and, from the proof of Lemma 5.2.1,

$$\begin{aligned}
[z^n]\bigl[(1-z)^{-1}L^2(z)\bigr] = H_n^2 - \sum_{k=1}^{n}\frac{1}{k^2} &= \Bigl(\ln n + \gamma + O(n^{-1})\Bigr)^2 - \sum_{k=1}^{\infty}\frac{1}{k^2} + O(n^{-1}) \\
&= (\ln n)^2 + 2\gamma\ln n + \left(\gamma^2 - \frac{\pi^2}{6}\right) + O\!\left(\frac{\ln n}{n}\right).
\end{aligned}$$

So, by singularity analysis (Theorem 4.1.2),

$$\begin{aligned}
\tilde{\mu}_n(2) = (C_1^2+2V)(n+1) + (\ln n)^2 + 2\gamma\ln n + \left(\gamma^2 - \frac{\pi^2}{6}\right) + (4-V_1)(\ln n + \gamma) \\
+ (4-2V_1-V_2) + O(n^{-1+\epsilon}).
\end{aligned} \tag{5.2.13}$$

In order to obtain the variance using (5.2.1) we also need [recall (5.1.35)]

$$\tilde{\mu}_n(1)^2 = (\ln n)^2 + 4\ln n + 4 + O(n^{-1+\epsilon}).$$

Hence, using (5.2.10) and (5.2.11) we obtain an expansion for the variance:

$$\begin{aligned}
\sigma_n^2 &= (C_1^2+2V)(n+1) + (2\gamma - V_1)\ln n + \left[\left(\gamma^2 - \frac{\pi^2}{6}\right) + (4-V_1)\gamma - 2V_1 - V_2\right] + O(n^{-1+\epsilon}) \\
&= (C_1^2+2V)(n+1) - \left(4 - \frac{\pi^2}{3}\right) + O(n^{-1+\epsilon}).
\end{aligned} \tag{5.2.14}$$

Numerical calculations using `Mathematica` suggest that the next term in the expansion is $-\frac{1}{2n}$.

One could possibly obtain this by carrying out the program of this section, keeping track of more terms in the asymptotic expansions.

Chapter 6

Random permutation model: limiting distributions

In this chapter we will apply the Asymptotic Transfer Theorem (Theorem 3.2.1) of Chapter 3 to derive limiting distributions for additive functionals defined on m-ary search trees under the random permutation model. Our main results are Theorems 6.1.1, 6.1.4, 6.2.2, and 6.2.4.

Throughout $\mathcal{N}(\mu, \sigma^2)$ denotes the normal distribution with mean μ and variance σ^2.

6.1 Small toll functions

We begin with two central limit theorems (Theorems 6.1.1 and 6.1.4) for "small" toll functions. Throughout, we write $\sum_{\mathbf{j}}$ as shorthand for the sum over m-tuples $(j_1, \ldots, j_m)$ of nonnegative integers summing to $n - (m-1)$.

Theorem 6.1.1. *If $2 \le m \le 26$ and the real toll sequence (t_n) satisfies*

$$\textit{(a) } t_n = o(\sqrt{n}) \quad \text{and} \quad \textit{(b) } \sum^{\infty} n^{-1} \max_{n^\delta \le k \le n} \frac{t_k^2}{k} < \infty \quad \text{for some } 0 < \delta < 1, \tag{6.1.1}$$

then the mean μ_n and the variance σ_n^2 of the corresponding additive functional X_n on m-ary search trees with the random permutation model satisfy, respectively,

$$\mu_n = \frac{K_1}{H_m - 1} n + o(\sqrt{n}) =: \mu n + o(\sqrt{n}), \tag{6.1.2}$$

with K_1 defined as

$$K_1 := \sum_{j=0}^{\infty} \frac{t_j}{(j+1)(j+2)}, \tag{6.1.3}$$

and

$$\sigma_n^2 = \sigma^2 n + o(n), \qquad \textit{where} \qquad \sigma^2 := \frac{1}{H_m - 1} \sum_{j=0}^{\infty} \frac{r_j}{(j+1)(j+2)}, \tag{6.1.4}$$

with the sequence (r_n) defined by $r_j := 0$ for $0 \le j \le m-2$ and

$$r_n := \frac{1}{\binom{n}{m-1}} \sum_{\mathbf{j}} [t_n + \mu_{j_1} + \cdots + \mu_{j_m} - \mu_n]^2, \qquad n \ge m-1. \tag{6.1.5}$$

Moreover,

$$\frac{X_n - \mu n}{\sqrt{n}} \xrightarrow{\mathcal{L}} \mathcal{N}(0, \sigma^2),$$

and there is convergence of moments of every order.

Proof. One can check (for *any* $2 \le m < \infty$) that the variance σ_n^2 vanishes for all $n \ge m-1$ if and only if the toll sequence is chosen as

$$t_n = t \min\{m-1, n\}, \qquad n \ge 0$$

for some constant $t \in \mathbb{R}$. In that case without loss of generality $t = 1$ and then $X_n \equiv n$ is just the number of keys and we have exact (though degenerate) normality. So we shall assume that $\sigma^2 > 0$.

We use the method of moments together with the asymptotic transfer results of Section 3.2.

Given the toll sequence (t_n) defining the sequence (X_n) of random functionals of interests, the means (μ_n) satisfy precisely the recurrence (2.3.3) with (b_n) replaced by (t_n). Thus (6.1.2) simply repeats the asymptotic transfer result (3.2.11).

According to the law of total variance, the sequence (σ_n^2) also satisfies the recurrence (2.3.3), but with (b_n) replaced by (r_n). Furthermore, according to Lemma 6.1.2 to follow, the sequence (r_n) satisfies the condition (3.2.1). Then (6.1.4) is immediate from part (a) of the ATT (Theorem 3.2.1).

Let $\widetilde{X}_n := X_n - \mu(n+1)$ for $n \geq 0$. We will complete the proof by showing by induction on k that

$$\tilde{\mu}_n(k) := \mathbf{E}\, \widetilde{X}_n^k, \quad k \geq 1 \quad [\text{with } \tilde{\mu}_n(0) := 1] \tag{6.1.6}$$

satisfies

$$\tilde{\mu}_n(2k) \sim \frac{(2k)!}{2^k k!}\sigma^{2k} n^k, \qquad \text{and} \qquad \tilde{\mu}_n(2k-1) = o(n^{k-\frac{1}{2}}), \quad k \geq 1. \tag{6.1.7}$$

Observe that (6.1.2) and (6.1.4) imply that (6.1.7) holds for $k = 1$.

The key to the induction step for (6.1.7) is to apply the law of total expectation to (6.1.6), by conditioning on the principal subtree sizes $|L_1|, \ldots, |L_m|$. Letting $\sum_{\mathbf{k}}$ denote the sum over $(m+1)$-tuples $(k_1, \ldots, k_{m+1})$ of nonnegative integers summing to k, for $n \geq m-1$ we have in a manner identical to that for (2.3.1),

$$\tilde{\mu}_n(k) = \frac{m}{\binom{n}{m-1}} \sum_{j=0}^{n-(m-1)} \binom{n-1-j}{m-2} \tilde{\mu}_j(k) + r_n(k), \tag{6.1.8}$$

where

$$r_n(k) := \sum_{\mathbf{k}}^{*} \binom{k}{k_1, \ldots, k_m, k_{m+1}} t_n^{k_{m+1}} \frac{1}{\binom{n}{m-1}} \sum_{\mathbf{j}} \tilde{\mu}_{j_1}(k_1) \cdots \tilde{\mu}_{j_m}(k_m),$$

with $\sum_{\mathbf{k}}^{*}$ denoting the same sum as $\sum_{\mathbf{k}}$ with the additional restriction that $k_i < k$ for $i = 1, \ldots, m$. Observe that (6.1.8) is again of the same basic form (2.3.3). We will apply the ATT after evaluating $r_n(k)$ asymptotically.

We will treat the induction step in detail only for $\tilde{\mu}_n(2k)$ at (6.1.7), the case for $\tilde{\mu}_n(2k-1)$ being similar and somewhat easier. Let $\sum_{\mathbf{k}}^{**}$ denote the sum over m-tuples $(k_1, \ldots, k_m)$ of nonnegative integers, each strictly less than k, summing to k (i.e., the same sum as $\sum_{\mathbf{k}}^{*}$ with the additional restriction that $k_{m+1} = 0$). For $k \geq 2$, we clearly have, by induction,

$$\begin{aligned}
r_n(2k) &= o(n^k) + \sum_{\mathbf{k}}^{**} \binom{2k}{2k_1, \ldots, 2k_m} \frac{1}{\binom{n}{m-1}} \sum_{\mathbf{j}} \tilde{\mu}_{j_1}(2k_1) \cdots \tilde{\mu}_{j_m}(2k_m) \\
&= o(n^k) + \sum_{\mathbf{k}}^{**} \binom{2k}{2k_1, \ldots, 2k_m} \frac{1}{\binom{n}{m-1}} \sum_{\mathbf{j}} \frac{(2k)!}{2^{k_1} k_1!} \sigma^{2k_1} j_1^{k_1} \cdots \frac{(2k_m)!}{2^{k_m} k_m!} \sigma^{2k_m} j_m^{k_m} \\
&= o(n^k) + \frac{(2k)!}{2^k k!} \sigma^{2k} n^k \sum_{\mathbf{k}}^{**} \binom{k}{k_1, \ldots, k_m} \frac{1}{\binom{n}{m-1}} \sum_{\mathbf{j}} \left(\frac{j_1}{n}\right)^{k_1} \cdots \left(\frac{j_m}{n}\right)^{k_m}.
\end{aligned}$$

But

$$\begin{aligned}&\frac{1}{\binom{n}{m-1}}\sum_{\mathbf{j}}\left(\frac{j_1}{n}\right)^{k_1}\cdots\left(\frac{j_m}{n}\right)^{k_m}\\&\to(m-1)!\int x_1^{k_1}\cdots x_{m-1}^{k_m-1}(1-x_1-\cdots-x_{m-1})^{k_m}\,dx_1\cdots dx_{m-1}\\&=(m-1)!\frac{\Gamma(k_1+1)\cdots\Gamma(k_m+1)}{\Gamma(k+m)}=\frac{1}{\binom{k}{k_1,\dots,k_m}\binom{k+m-1}{m-1}},\end{aligned}$$

where the above integral is over $(x_1,\dots,x_{m-1})\in[0,1]^{m-1}$ with sum not exceeding unity. Since the number of terms in $\sum_{\mathbf{k}}^{**}$ is $\binom{k+m-1}{m-1}-m$, we therefore have

$$r_n(2k)=\frac{(2k)!}{2^k k!}\sigma^{2k}n^k\frac{\binom{k+m-1}{m-1}-m}{\binom{k+m-1}{m-1}}+o(n^k)=\frac{(2k)!}{2^k k!}\sigma^{2k}n^k\left[1-\frac{m!\Gamma(k+1)}{\Gamma(k+m)}\right]+o(n^k),\qquad k\geq 2.$$

Similarly,

$$r_n(2k-1)=o(n^{k-\frac{1}{2}}),\qquad k\geq 2.$$

Now part (c) of the ATT (Theorem 3.2.1) implies (6.1.7). ☐

The following lemma lies at the heart of the proof of Theorem 6.1.1.

Lemma 6.1.2. *In the context of Theorem 6.1.1, the sequence* (r_n) *defined at* (6.1.5) *satisfies the conditions* (3.2.1).

Proof. Clearly

$$r_n=\frac{1}{\binom{n}{m-1}}\sum_{\mathbf{j}}[t_n+\tilde{\mu}_{j_1}+\cdots+\tilde{\mu}_{j_m}-\tilde{\mu}_n]^2,\qquad n\geq m-1,\tag{6.1.9}$$

with

$$\tilde{\mu}_n:=\mu_n-\mu(n+1),\tag{6.1.10}$$

which is $o(\sqrt{n})$ by (6.1.2). Recall the inequality

$$\left[\sum_{i=1}^{k}\xi_i\right]^2\leq k\sum_{i=1}^{k}\xi^2\tag{6.1.11}$$

for real numbers $\xi_1,\dots,\xi_k$. Applying this to (6.1.9),

$$\frac{r_n}{m+2}\leq t_n^2+\tilde{\mu}_n^2+\frac{m}{\binom{n}{m-1}}\sum_{j=0}^{n-(m-1)}\binom{n-1-j}{m-2}\tilde{\mu}_j^2,\tag{6.1.12}$$

from which it is clear that $r_n=o(n)$.

To establish the summability of r_n/n^2 we need only establish that of $\tilde{\mu}_n^2/n^2$. Indeed we may then use (6.1.12) again, together with the fact that (6.1.1)(b) implies

$$\sum^{\infty}\frac{t_n^2}{n^2}<\infty\tag{6.1.13}$$

and the estimate

$$\sum_{n=m-1}^{\infty} n^{-2} \frac{m}{\binom{n}{m-1}} \sum_{j=0}^{n-(m-1)} \binom{n-1-j}{m-2} \tilde{\mu}_j^2 = m(m-1) \sum_{j=0}^{\infty} \tilde{\mu}_j^2 \sum_{n=j+m-1}^{\infty} \frac{(n-1-j)^{\underline{m-2}}}{n^2 n^{\underline{m-1}}}$$
$$\leq m(m-1) \sum_{j=0}^{\infty} \tilde{\mu}_j^2 \sum_{n=j+m-1}^{\infty} n^{-3} = O\left(\sum \frac{\tilde{\mu}_j^2}{j^2}\right) < \infty.$$

To establish the summability of $\tilde{\mu}_n^2/n^2$, we recall from (3.2.15) and (3.2.8) that

$$\tilde{\mu}_n = O(n^{\alpha-1}) + t_n - \frac{1}{H_m - 1}(n+1) \sum_{k=n}^{\infty} \frac{\hat{t}_k}{(k+1)(k+2)} + \sum_{j=2}^{m-1} O\left(n^{\alpha_j - 1} \sum_{k=0}^{n-1} \frac{|\hat{t}_k|}{(k+1)^{\alpha_j}}\right), \tag{6.1.14}$$

writing $\alpha_j := \Re(\lambda_j)$ (with $\alpha = \alpha_2 < 3/2$, since $m \leq 26$). Using (6.1.11), we need only establish the summability of n^{-2} times the square of each of the four terms on the right in (6.1.14). The first of these verifications is trivial, and the second was carried out at (6.1.13). For the third we apply the Cauchy–Schwarz inequality to give

$$\left[\sum_{k=n}^{\infty} \frac{\hat{t}_k}{(k+1)(k+2)}\right]^2 \leq \frac{1}{n}\left[\sum_{k=n}^{\infty} \sqrt{n} \frac{\sqrt{k}}{(k+1)(k+2)} \frac{|t_k|}{\sqrt{k}}\right]^2$$
$$= O\left(\frac{1}{n} \sum_{k=n}^{\infty} \sqrt{n} k^{-3/2} \frac{t_k^2}{k}\right) = O\left(n^{-1/2} \sum_{k=n}^{\infty} k^{-5/2} t_k^2\right),$$

whence

$$\sum_n \left[\sum_{k=n}^{\infty} \frac{\hat{t}_k}{(k+1)(k+2)}\right]^2 = O\left(\sum_n n^{-1/2} \sum_{k=n}^{\infty} k^{-5/2} t_k^2\right)$$
$$= O\left(\sum_k k^{-5/2} t_k^2 k^{1/2}\right) = O\left(\sum_k \frac{t_k^2}{k^2}\right) < \infty$$

by (6.1.13) again.

We pause to note that when $m = 2$ the proof is finished here, and that up to now we have used only (6.1.13), not the stronger assumption (6.1.1)(b).

For our fourth and final verification, it suffices [again by invoking (6.1.11)] to establish the summability of

$$n^{2\rho-4} \left[\sum_{k=1}^{n-1} \frac{|t_k|}{k^\rho}\right]^2 \tag{6.1.15}$$

for any real $\rho < 3/2$. To do this we break the sum into $\sum_{k<n^\delta}$ and $\sum_{n^\delta \leq k < n}$ and once again invoke (6.1.11). In the range $\sum_{k<n^\delta}$ we simply use $t_k = O(\sqrt{k})$ and note that

$$n^{2\rho-4} \left[\sum_{k<n^\delta} O\left(k^{\frac{1}{2}-\rho}\right)\right]^2 = O\left(n^{2\rho-4} (n^\delta)^{3-2\rho}\right) = O(n^\tau)$$

with $\tau < -1$. In the range $\sum_{n^\delta \le k < n}$ we use Cauchy–Schwarz again:

$$n^{2\rho-4}\left[\sum_{n^\delta \le k<n} \frac{|t_k|}{k^\rho}\right]^2 = n^{2\rho-4} n^{3-2\rho}\left[\sum_{n^\delta \le k<n} \frac{k^{\frac{1}{2}-\rho}}{n^{\frac{3}{2}-\rho}} \frac{|t_k|}{k^{1/2}}\right]^2$$

$$= O\left(n^{-1} \sum_{n^\delta \le k<n} \frac{k^{\frac{1}{2}-\rho}}{n^{\frac{3}{2}-\rho}} \frac{t_k^2}{k}\right) = O\left(n^{-1} \max_{n^\delta \le k<n} \frac{t_k^2}{k}\right),$$

which is summable by assumption (6.1.1)(b). □

Remark 6.1.3. (a) We have already noted in the proof above that condition (6.1.1)(b) trivially implies

$$\sum^\infty \frac{t_n^2}{n^2} < \infty, \tag{6.1.16}$$

which in turn implies that (3.2.1) holds with absolute convergence; indeed, since the nonnegative numbers $[(n+1)(n+2)]^{-1}$, $n \ge 0$, sum to unity, we have

$$\left[\sum_{n=0}^\infty \frac{|t_n|}{(n+1)(n+2)}\right]^2 \le \sum_{n=0}^\infty \frac{t_n^2}{(n+1)(n+2)} < \infty. \tag{6.1.17}$$

(b) If

$$|t_n| = O(\tilde{t}_n) \quad \text{with} \quad 0 \le \frac{\tilde{t}_n}{\sqrt{n}} \downarrow \quad \text{and} \quad \sum^\infty \frac{\tilde{t}_n^2}{n^2} < \infty, \tag{6.1.18}$$

then we claim that (6.1.1) holds, and then as a corollary

$$\frac{t_n}{\sqrt{n}} \downarrow 0 \quad \text{and} \quad \sum^\infty \frac{t_n^2}{n^2} < \infty$$

implies (6.1.1). To see this claim, first observe that the condition (6.1.18) implies (6.1.1)(a); moreover, we observe that the series (say, over $2 \le n < \infty$) in (6.1.1)(b) is bounded by a constant times

$$\sum_{n=2}^\infty n^{-1} \max_{n^\delta \le k \le n} \frac{\tilde{t}_k^2}{k} = \sum_{n=2}^\infty n^{-1} \frac{\tilde{t}_{\lceil n^\delta \rceil}^2}{\lceil n^\delta \rceil} = \sum_{k=2}^\infty \frac{\tilde{t}_k^2}{k} \sum_{(k-1)^{1/\delta} < n \le k^{1/\delta}} n^{-1} = O\left(\sum_{k=2}^\infty \frac{\tilde{t}_k^2}{k^2}\right) < \infty.$$

(c) We have also observed that, when $m = 2$, the proof of Theorem 6.1.1 requires only (6.1.1)(a) and (6.1.16). In that case we obtain a strengthening of "Case S1" of Theorem 2 in [38] (for deterministic toll sequences); they required $t_n = O(\sqrt{n}/(\ln n)^{\frac{1}{2}+\epsilon})$ for some $\epsilon > 0$.

We can also obtain asymptotic normality in the "borderline small" case. The definition of slowly varying is available as Definition 3.2.6.

Theorem 6.1.4. *If* $2 \le m \le 26$ *and the real toll sequence* (t_n) *satisfies*

$$t_n \sim \sqrt{n} L(n) \tag{6.1.19}$$

with L *slowly varying, then the mean* μ_n *of the corresponding additive functional* X_n *on* m*-ary* *search trees with the random permutation model satisfies*

$$\mu_n = \frac{K_1}{H_m - 1} n - \frac{(3/2)^{\overline{m-1}}}{m! - (3/2)^{\overline{m-1}}} \sqrt{n} L(n) + o(\sqrt{n} L(n)), \tag{6.1.20}$$

with K_1 *defined at* (6.1.3)*.*

1. *If* $\sum^{\infty} \frac{L^2(k)}{k} < \infty$, *then the variance* σ_n^2 *satisfies* (6.1.4)*–*(6.1.5) *and we define*

$$s^2(n) := \sigma^2 n.$$

2. *If* $\sum^{\infty} \frac{L^2(k)}{k} = \infty$, *then*

$$\sigma_n^2 \sim s^2(n) := \sigma^2 n \sum_{k \le n} \frac{L^2(k)}{k}, \tag{6.1.21}$$

where in this case we define

$$\sigma^2 := \frac{((3/2)^{\overline{m-1}})^2 [\frac{\pi}{4}(m-1) + 1] - (m!)^2}{(H_m - 1)[m! - (3/2)^{\overline{m-1}}]^2}. \tag{6.1.22}$$

Moreover, in either case

$$\frac{X_n - \mu n}{s(n)} \xrightarrow{\mathcal{L}} \mathcal{N}(0,1)$$

and there is convergence of moments of every order.

Proof sketch. Given the similarity to the proof of Theorem 6.1.1, we will be brief here. Again we use the method of moments together with asymptotic transfer results.

Equation (6.1.20) simply repeats the transfer result (3.2.12). As before, (σ_n^2) satisfies the recurrence (2.3.3) with (b_n) replaced by (r_n) of (6.1.9)–(6.1.10), where again $\mu := K_1/(H_m - 1)$ and $r_j := 0$ for $0 \le j \le m-2$. Here the proofs diverge somewhat. The analogue of Lemma 6.1.2 is Lemma 6.1.5 below. Then the asymptotic variance assertions of the theorem follow immediately from Theorem 3.2.7(c).

If $\sum^{\infty} \frac{L^2(k)}{k} < \infty$, then from (3.2.10) applied to L^2 it follows that (6.1.6) satisfies (6.1.7) for $k = 1$. Then higher moments are treated exactly as in the proof of Theorem 6.1.1 to complete the proof.

If $\sum^{\infty} \frac{L^2(k)}{k} = \infty$, then one uses (3.2.10), Theorem 3.2.7(d), and induction to show that the moments (6.1.6) satisfy

$$\tilde{\mu}_n(2k) \sim \frac{(2k)!}{2^k k!} s^{2k}(n) \quad \text{and} \quad \tilde{\mu}_n(2k-1) = o(s^{2k-1}(n)), \qquad k \ge 1 \tag{6.1.23}$$

and thereby complete the proof. We omit the details. □

The following cousin to Lemma 6.1.2 was used in the proof of Theorem 6.1.4 above.

Lemma 6.1.5. *In the context of Theorem 6.1.4, the sequence* (r_n) *defined for* $n \ge m-1$ *by*

$$r_n := \frac{1}{\binom{n}{m-1}} \sum_{\mathbf{j}} [t_n + \tilde{\mu}_{j_1} + \cdots + \tilde{\mu}_{j_m} - \tilde{\mu}_n]^2$$

satisfies

$$r_n \sim (H_m - 1)\sigma^2 n L^2(n).$$

Proof. By (6.1.20), with $\theta := (3/2)^{\overline{m-1}}/(m! - (3/2)^{\overline{m-1}})$, we have

$$r_n \sim \frac{1}{\binom{n}{m-1}} \sum_{\mathbf{j}} [n^{1/2}L(n) - \theta_{j_1}^{1/2}L(j_1) - \cdots - \theta j_m^{1/2}L(j_m) + \theta n^{1/2}L(n)]^2$$

$$\sim nL^2(n)(m-1)! \int \left[(1+\theta) - \theta \sum_{i=1}^{m-1} x_i^{1/2} - \theta \left(1 - \sum_{i=1}^{m-1} x_i\right)^{1/2}\right]^2 dx_1 \cdots dx_{m-1},$$

where the integral, call it J, is over $(x_1, \dots, x_{m-1}) \in [0,1]^{m-1}$ with the sum not exceeding unity. To complete the proof we need only show $J = (H_m - 1)\sigma^2/(m-1)!$, with σ^2 defined at (6.1.22).

Indeed, J equals

$$\int \Bigg[(1+\theta)^2 + \theta^2 \sum_{i=1}^{m-1} x_i + \theta^2 \left(1 - \sum_{i=1}^{m-1} x_i\right) - 2\theta(1+\theta) \sum_{i=1}^{m-1} x_i^{1/2} - 2\theta(1+\theta)\left(1 - \sum_{i=1}^{m-1} x_i\right)^{1/2}$$

$$+ \theta^2 \sum_{i,j:i\neq j} x_i^{1/2} x_j^{1/2} + 2\theta^2 \left(1 - \sum_{i=1}^{m-1} x_i\right)^{1/2} \sum_{i=1}^{m-1} x_i^{1/2}\Bigg] dx_1 \cdots dx_{m-1}$$

$$= [(1+\theta)^2 + \theta^2]\frac{1}{(m-1)!} - 2\theta(1+\theta)m\frac{\Gamma(\frac{3}{2})}{\Gamma(m+\frac{1}{2})} + \theta^2 m(m-1)\frac{[\Gamma(3/2)]^2}{\Gamma(m+1)}$$

$$= [(1+\theta)^2 + \theta^2]\frac{1}{(m-1)!} - 2\theta(1+\theta)m\frac{1}{(3/2)^{\overline{m-1}}} + \theta^2\frac{\pi/4}{(m-2)!}.$$

Plugging in the value of θ and simplifying, we obtain $J = (H_m - 1)\sigma^2/(m-1)!$, as desired. □

Remark 6.1.6. When $m = 2$ the constant σ^2 in (6.1.22) equals $\frac{9}{2}\pi - 14$, and Theorem 6.1.4 reduces to "Case S2" of Theorem 2 in [38] (for deterministic toll sequences): see especially their displays (15) and (17), with $\tau_2 = 1$.

Periodicity for $m \geq 27$

If $t_n = o(\sqrt{n})$ as in Theorem 6.1.1 but $m \geq 27$, then the remainder term $\tilde{\mu}_n := \mu_n - \mu(n+1)$ for the mean—which by (6.1.2) was $o(\sqrt{n})$ when $m \leq 26$—now satisfies, by the ETT (Theorem 3.1.1) and (3.2.9) [compare (3.2.15)]

$$\tilde{\mu}_n = c_2\frac{n^{\lambda_2 - 1}}{\Gamma(\lambda_2)} + c_3\frac{n^{\lambda_3 - 1}}{\Gamma(\lambda_3)} + m!\sum_{j=2}^{m-1}\frac{1}{\psi'(\lambda_j)}[z^n]\left((1-z)^{-\lambda_j}\int_{\zeta=0}^{z}\widehat{T}(\zeta)(1-\zeta)^{\lambda_j - 1}\,d\zeta\right)$$
$$+ o(\sqrt{n}) + O(n^{\Re(\lambda_4)-1}). \quad (6.1.24)$$

Typically this will lead to the negative result that $(\tilde{\mu}_n)$ [and hence also (r_n) and (σ_n^2)] suffers from periodicity and that there is no natural distributional limit for normalized X_n. See, e.g., Corollary 1 of [8].

But it is perhaps difficult to establish a *general* negative result, due to cancellations. For example, suppose $T(z)$ equals $(1-z)^{-1}$, so that $t_n \equiv 1$, as studied by Chern and Hwang [8], except perhaps that the initial values $t_0, \dots, t_{m-2}$ are changed. Then

$$T^{(m-1)}(z) \equiv (m-1)!(1-z)^{-m},$$

whence

$$\begin{aligned} A(z) := \sum_{n=0}^{\infty} \mu_n z^n &= \sum_{j=1}^{m-1} c_j (1-z)^{-\lambda_j} + (m-1)! \sum_{j=1}^{m-1} \frac{(1-z)^{-\lambda_j}}{\psi'(\lambda_j)} \int_{\zeta=0}^{z} (1-\zeta)^{\lambda_j - 2}\, d\zeta \\ &= \sum_{j=1}^{m-1} c_j (1-z)^{-\lambda_j} - \frac{1}{m-1}(1-z)^{-1} - (m-1)! \sum_{j=1}^{m-1} \frac{(1-z)^{-\lambda_j}}{(1-\lambda_j)\psi'(\lambda_j)}. \end{aligned}$$

Now it is possible to choose $t_0, \ldots, t_{m-2}$ so that

$$c_j = \frac{(m-1)!}{(1-\lambda_j)\psi'(\lambda_j)}, \qquad j = 1, \ldots, m-1. \tag{6.1.25}$$

In that case $A(z) = -\frac{1}{m-1}(1-z)^{-c}$, whence $\tilde{\mu}_n \equiv \mu_n \equiv -1/(m-1)$ [and we see that the chosen values of $t_0, \ldots, t_{m-2}$ are all $-1/(m-1)$], and we get linear variance and asymptotic normality, just as in Theorem 6.1.1, for *every* $2 \le m < \infty$.

One might object that the above example is artificial, in that the toll sequence changes sign. But the same calculation show that if the toll sequence is chosen as above ($t_n \equiv 1$) but with inital values

$$t_j := K(j+1) - \frac{1}{m-1}, \qquad 0 \le j \le m-2,$$

then still, for every $m \ge 2$, the sequence $(\tilde{\mu}_n)$ is constant, the variance is linear, and we have asymptotic normality. Further, (t_n) is nonnegative provided $K \ge 1/(m-1)$. [We remark in passing that the choice $K = 1/(m-1)$ leads to the degenerate case mentioned at the beginning of the proof of Theorem 6.1.1.] The sequence (t_n) is also nondecreasing (as in most real examples) provided $K \le m/(m-1)^2$.

6.2 Moderate and large toll functions

In order to describe the limiting distribution of X_n for moderate and large tolls, we will introduce a family of random variables $Y \equiv Y_\beta$ defined for $\beta > 1/2$, $\beta \ne 1$. Anticipating Lemma 6.2.1, we need to consider the distributional equation

$$Y \overset{\mathcal{L}}{=} \sum_{j=1}^{m} S_j^{\beta} Y_j + 1 \tag{6.2.1}$$

Here $(Y_j)_{j=1}^m$ are independent copies of Y and $(S_1, \ldots, S_m)$ is uniformly distributed on the $(m-1)$-simplex, independent of $(Y_j)_{j=1}^m$. Recall that the $(m-1)$-simplex is the set

$$\{(s_1, \ldots, s_m)\colon s_j \ge 0 \text{ for } 1 \le j \le m \text{ and } \mathbf{s}_+ = 1\},$$

where $\mathbf{s}_+$ denotes $\sum_{j=1}^m s_j$.

Let $U_{(1)}, \ldots, U_{(m-1)}$ be the order statistics of a sample of size $m-1$ from $\text{Unif}(0,1)$ with joint density

$$f_{U_{(1)},\ldots,U_{(m-1)}}(x_1, \ldots, x_{m-1}) \equiv (m-1)!\, \mathbf{1}(0 < x_1 < \cdots < x_{m-1} < 1)$$

with respect to Lebesgue measure on $\mathbb{R}^{m-1}$, where $\mathbf{1}(A)$ is the indicator of A.

By a change of variables, we find that the joint distribution of the spacings $S_1, \ldots, S_m$, defined, with $U_{(0)} := 0$ and $U_{(m)} := 1$, by

$$S_i := U_{(i)} - U_{(i-1)},\ i = 1, \ldots, m,$$

is uniform over the $(m-1)$-simplex:

$$f_{S_1,\dots,S_{m-1}}(s_1,\dots,s_{m-1}) \equiv (m-1)!\,\mathbf{1}(s_j>0, j=1,\dots,m-1;\ \mathbf{s}_+<1)\,.$$

When $r_j > -1$ for $1 \le j \le m$, observe that

$$\begin{aligned}\mathbf{E}\left[\prod_{j=1}^m S_j^{r_j}\right] &= (m-1)!\int_{\substack{s_1,\dots,s_{m-1}>0\\ \mathbf{s}_+<1}} s_1^{r_1}\cdots s_{m-1}^{r_{m-1}}(1-\mathbf{s}_+)^{r_m}\,ds_{m-1}\cdots ds_1\\ &=: (m-1)!B(r_1+1,\dots,r_m+1) \qquad (6.2.2)\\ &= (m-1)!\frac{\prod_{j=1}^m \Gamma(r_j+1)}{\Gamma(r_1+\cdots+r_m+m)}.\end{aligned}$$

Lemma 6.2.1. *Fix $\beta > 1/2$ with $\beta \ne 1$. Then there exists a unique distribution $\mathcal{L}(Y) \equiv \mathcal{L}(Y_\beta)$ with finite second moment satisfying the distributional identity* (6.2.1).

Proof. We first observe the that mean of any such distribution is determined by (6.2.1). Indeed, by taking expectations in (6.2.1) and using (6.2.2), we get

$$\mu := \mathbf{E}\,Y = \left(1 - \frac{m!\Gamma(\beta+1)}{\Gamma(\beta+m)}\right)^{-1}$$

since $\beta \ne 1$. Thus we can equivalently consider the distributional identity

$$W \stackrel{\mathcal{L}}{=} \sum_{j=1}^m S_j^\beta W_j + H,$$

where

$$H := 1 - \mu + \mu\sum_{j=1}^m S_j^\beta.$$

Here W is restricted to have mean 0 and finite second moment, $(W_j)_{j=1}^m$ are independent copies of W, and $(S_1,\dots,S_m)$ is uniformly distributed on the $(m-1)$-simplex, independent of $(W_j)_{j=1}^m$.

We now employ a standard contraction-method argument [64, 65]. Let d_2 denote the metric on $\mathcal{M}_2(0)$, the space of probability distributions with mean 0 and finite variance, defined by

$$d_2(G_1,G_2) := \min\|X_2 - X_1\|_2,$$

taking the minimum over all pairs of random variables X_1 and X_2 defined on a common probability space with $\mathcal{L}(X_1) = G_1$ and $\mathcal{L}(X_2) = G_2$. Here $\|\cdot\|_2$ denotes L_2-norm.

Let T be the map

$$T : \mathcal{M}_2(0) \to \mathcal{M}_2(0), \quad G \mapsto \mathcal{L}\left(\sum_{j=1}^m S_j^\beta X_j + H\right),$$

where $(X_j)_{j=1}^m$ are independent with $\mathcal{L}(X_j) = G$, $j = 1,\dots,m$, and $(S_1,\dots,S_m)$ is uniformly distributed on the $(m-1)$-simplex, independent of $(X_j)_{j=1}^m$. We show that T is a contraction on $\mathcal{M}_2(0)$; more precisely, that there exists a $\rho < 1$ such that

$$d_2(T(\mathcal{L}(A)), T(\mathcal{L}(B))) \le \rho d_2(\mathcal{L}(A), \mathcal{L}(B))$$

for all pairs $\mathcal{L}(A)$ and $\mathcal{L}(B)$ in $\mathcal{M}_2(0)$. To bound $d_2(T(\mathcal{L}(A)), T(\mathcal{L}(B)))$, we couple $T(\mathcal{L}(A))$

and $T(\mathcal{L}(B))$ by taking m independent copies (A_j, B_j) of the d_2-optimally coupled (A, B), an independent $(S_1, \dots, S_m)$, and defining

$$A' := \sum_{j=1}^{m} S_j^{\beta} A_j + H \sim T(\mathcal{L}(A)), \quad B' := \sum_{j=1}^{m} S_j^{\beta} B_j + H \sim T(\mathcal{L}(B)).$$

Now, defining $\mathbf{S} := (S_1, \dots, S_m)$ and using the law of total variance,

$$\begin{aligned}
& d_2(\mathcal{L}(T(A)), \mathcal{L}(T(B)))^2 \\
& \le \|B' - A'\|_2^2 = \left\| \sum_{j=1}^{m} S_j^{\beta}(B_j - A_j) \right\|_2^2 = \mathbf{Var} \left[\sum_{j=1}^{m} S_j^{\beta}(B_j - A_j) \right] \\
& = \mathbf{E}\ \mathbf{Var} \left[\sum_{j=1}^{m} S_j^{\beta}(B_j - A_j) \middle| \mathbf{S} \right] + \mathbf{Var}\ \mathbf{E} \left[\sum_{j=1}^{m} S_j^{\beta}(B_j - A_j) \middle| \mathbf{S} \right] \\
& = \sum_{j=1}^{m} (\mathbf{E}\, S_j^{2\beta}) \mathbf{Var}\ [B_j - A_j] = d_2(\mathcal{L}(A), \mathcal{L}(B))^2 \sum_{j=1}^{m} \mathbf{E}\, S_j^{2\beta} = m! \frac{\Gamma(2\beta + 1)}{\Gamma(2\beta + m)} d_2(A, B)^2.
\end{aligned}$$

We need only verify that

$$\rho^2 := m! \frac{\Gamma(2\beta + 1)}{\Gamma(2\beta + m)} = \frac{m!}{(2\beta + m - 1) \cdots (2\beta + 1)} < 1,$$

which is true when $\beta > 1/2$. The existence and uniqueness of $\mathcal{L}(Y)$ now follows from the Banach fixed point theorem [65, Theorem 2]. □

Moderate toll functions

In the case of moderate toll functions, convergence in distribution and convergence of all moments can be stated as

Theorem 6.2.2. *If the real toll sequence* (t_n) *satisfies*

$$t_n \sim n^{\beta}, \qquad 1/2 < \beta < 1,$$

and $\alpha < 1 + \beta$, *then the mean of the corresponding linear functional* X_n *on* m*-ary search trees with the random permutation model satisfies*

$$\mu_n = \mu n - \frac{(1 + \beta)^{\overline{m-1}}}{m! - (1 + \beta)^{\overline{m-1}}} n^{\beta} + o(n^{\beta}), \qquad \mu := \frac{K_1}{H_m - 1}, \tag{6.2.3}$$

with K_1 *defined at* (6.1.3)*. Moreover,*

$$\frac{X_n - \mu n}{n^{\beta}} \xrightarrow{\mathcal{L}} Y_{\beta},$$

with convergence of all moments.

Proof. We use the notation introduced in the proof of Theorem 6.1.1. Equation (6.2.3) is simply a restatement of the asymptotic transfer result (3.2.12).

We show that the moments $\tilde{\mu}_n(k)$ satisfy

$$\tilde{\mu}_n(k) = g_k n^{k\beta} + o(n^{k\beta}) \text{ as } n \to \infty. \tag{6.2.4}$$

The claim holds for $k = 1$ with

$$g_1 := -\frac{(1+\beta)^{\overline{m-1}}}{m! - (1+\beta)^{\overline{m-1}}} = \left(1 - \frac{m!\Gamma(\beta+1)}{\Gamma(\beta+m)}\right)^{-1}. \tag{6.2.5}$$

Using (6.1.9) and (6.1.8), by induction we get, for $k \geq 2$,

$$\begin{aligned} r_n(k) &= o(n^{k\beta}) + \sum_{\mathbf{k}}{}^{*} \binom{k}{k_1, \ldots, k_m, k_{m+1}} (n^\beta)^{k_{m+1}} \frac{1}{\binom{n}{m-1}} \sum_{\mathbf{j}} g_{k_1} j_1^{k_1\beta} \cdots g_{k_m} j_m^{k_m\beta} \\ &= o(n^{k\beta}) + \sum_{\mathbf{k}}{}^{*} \binom{k}{k_1, \ldots, k_m, k_{m+1}} n^{k\beta} g_{k_1} \cdots g_{k_m} \frac{1}{\binom{n}{m-1}} \sum_{\mathbf{j}} \left(\frac{j_1}{n}\right)^{k_1\beta} \cdots \left(\frac{j_m}{n}\right)^{k_m\beta}. \end{aligned}$$

But

$$\frac{1}{\binom{n}{m-1}} \sum_{\mathbf{j}} \left(\frac{j_1}{n}\right)^{k_1\beta} \cdots \left(\frac{j_m}{n}\right)^{k_m\beta} \to (m-1)! B(k_1\beta + 1, \ldots k_m\beta + 1)$$

so that

$$r_n(k) = o(n^{k\beta}) + n^{k\beta}(m-1)! \sum_{\mathbf{k}}{}^{*} \binom{k}{k_1, \ldots, k_m, k_{m+1}} g_{k_1} \cdots g_{k_m} B(k_1\beta + 1, \ldots, k_m\beta + 1).$$

Using Theorem 3.2.1 (the ATT), with $v = k\beta > 1$, we get

$$\tilde{\mu}_n(k) = o(n^{k\beta}) + n^{k\beta} \frac{(m-1)!}{1 - \frac{m!\Gamma(k\beta+1)}{\Gamma(k\beta+m)}} \sum_{\mathbf{k}}{}^{*} \binom{k}{k_1, \ldots, k_m, k_{m+1}} g_{k_1} \cdots g_{k_m} B(k_1\beta + 1, \ldots, k_m\beta + 1).$$

Thus, defining g_k recursively as

$$g_k = \frac{(m-1)!}{1 - \frac{m!\Gamma(k\beta+1)}{\Gamma(k\beta+m)}} \sum_{\mathbf{k}}{}^{*} \binom{k}{k_1, \ldots, k_{m+1}} g_{k_1} \cdots g_{k_m} B(k_1\beta + 1, \ldots, k_m\beta + 1), \tag{6.2.6}$$

we see that (6.2.4) holds for all $k \geq 1$.

By Lemma 6.2.3 (to follow) and the method of moments (cf., e.g., [9, Sections 4.4 and 4.5]), the g_k's are the moment of a uniquely determined distribution, say $\mathcal{L}(\widehat{Y})$, and

$$\frac{X_n - \mu n}{n^\beta} \xrightarrow{\mathcal{L}} \widehat{Y}$$

with convergence of all moments. It remains to show that $\widehat{Y} \overset{\mathcal{L}}{=} Y_\beta$.

Define

$$\widetilde{Y} := \sum_{j=1}^{m} S_j^\beta \widehat{Y}_j + 1, \tag{6.2.7}$$

where $(\widehat{Y}_j)_{j=1}^m$ are independent copies of $\widehat{Y}$ and $(S_1, \ldots, S_m)$ is uniformly distributed on the $(m-1)$-simplex, independent of $(\widehat{Y}_j)_{j=1}^m$. We will show that $\widehat{Y} \overset{\mathcal{L}}{=} \widetilde{Y}$, and then, by (6.2.7), $\mathcal{L}(\widehat{Y})$ satisfies the distributional identity (6.2.1) and has finite second moment. By Lemma 6.2.1, $\widehat{Y} \overset{\mathcal{L}}{=} Y_\beta$, as desired.

To show $\widehat{Y} \overset{\mathcal{L}}{=} \widetilde{Y}$, it suffices to show that $\widehat{Y}$ and $\widetilde{Y}$ have the same moments. Letting $\sum_{\mathbf{k}}$ denote (as before) the sum over $(m+1)$-tuples $(k_1, \ldots, k_{m+1})$ of nonnegative integers summing to k, and using (6.2.7), (6.2.2), and (6.2.6),

$$\begin{aligned}\mathbf{E}\,\widetilde{Y}^k &= \sum_{\mathbf{k}} \binom{k}{k_1,\dots,k_{m+1}} \mathbf{E}\left[\prod_{j=1}^m (S_j^\beta \widehat{Y}_j)^{k_j}\right] \\ &= \sum_{\mathbf{k}} \binom{k}{k_1,\dots,k_{m+1}} (m-1)!B(k_1\beta+1,\dots k_m\beta+1) g_{k_1}\cdots g_{k_m} \\ &= (m-1)! \sum_{\mathbf{k}}^{*} \binom{k}{k_1,\dots,k_{m+1}} B(k_1\beta+1,\dots k_m\beta+1) g_{k_1}\cdots g_{k_m} \\ &\quad + m!B(k\beta+1,1,\dots,1)g_k \\ &= \left[1-\frac{m!\Gamma(k\beta+1)}{\Gamma(k\beta+m)}\right] g_k + \frac{m!\Gamma(k\beta+1)}{\Gamma(k\beta+m)} g_k = g_k = \mathbf{E}\,\widehat{Y}^k,\end{aligned}$$

where (as before) $\sum_{\mathbf{k}}^*$ denotes the same sum as $\sum_{\mathbf{k}}$ with the additional restriction that $k_i < k$ for $i = 1,\dots,m$. ☐

Lemma 6.2.3. *The moments (g_k) uniquely determine the distribution $\mathcal{L}(Y)$.*

Proof. Define $\gamma_k := g_k/k!$. It suffices to show that there exists an M such that $\gamma_k \le M^k$ for all k sufficiently large. We proceed by induction. Indeed, by (6.2.6) we know

$$\begin{aligned}\gamma_k &= \frac{(m-1)!}{1-\frac{m!\Gamma(k\beta+1)}{\Gamma(k\beta+m)}} \sum_{\mathbf{k}}^{*} \frac{1}{k_{m+1}!}\left(\prod_{j=1}^m \gamma_{k_j}\right) B(k_1\beta+1,\dots,k_m\beta+1) \\ &\le M^k \frac{(m-1)!}{1-\frac{m!\Gamma(k\beta+1)}{\Gamma(k\beta+m)}} \sum_{\mathbf{k}}^{*} \frac{M^{-k_{m+1}}}{k_{m+1}!} B(k_1\beta+1,\dots,k_m\beta+1)\end{aligned}$$

by the induction hypothesis. So it is certainly sufficient to show that

$$\begin{aligned}&\sum_{\mathbf{k}}^{*} \frac{M^{-k_{m+1}}}{k_{m+1}!} B(k_1\beta+1,\dots,k_m\beta+1) \\ &= \sum_{k_{m+1}=0}^{k} \frac{M^{-k_{m+1}}}{k_{m+1}!} \Gamma((k-k_{m+1})\beta+m)^{-1} \sum_{\substack{0\le k_1,\dots,k_m<k \\ k_1+\cdots+k_m=k-k_{m+1}}} \prod_{j=1}^m \Gamma(k_j\beta+1) \\ &\to 0 \quad \text{as } k\to\infty. \end{aligned} \tag{6.2.8}$$

For this, fix a value of $k_{m+1} \in \{0,1,2,\dots\}$, and consider the sum

$$\sum_{\substack{0\le k_1,\dots,k_m<k \\ k_1+\cdots+k_m=k-k_{m+1}}} \prod_{j=1}^m \Gamma(k_j\beta+1). \tag{6.2.9}$$

By log-convexity of Γ [1, 6.4.1], taking I to be $(0,\infty)$ and g to be $\ln\Gamma$ in Proposition 3.C.1 of [52], the logarithm of the product in (6.2.9) is Schur-convex on $(0,\infty)^m$. Thus applying Proposition 5.C.2 of [52] with $m = 0$ there, the biggest terms in the sum correspond to $k_j = k - k_{m+1}$ for j equal to some j_0 and $k_j = 0$ otherwise; together, these m terms contribute $m\Gamma((k-k_{m+1})\beta+1)$ to the sum. If $k > k_{m+1}$, there are other terms, the biggest of which corresponds to having one of the k_j's be $k - k_{m+1} - 1$, one be 1, and the rest be 0. (This follows from Proposition 5.C.1 of [52] with $m = 0$ and $M = k - k_{m+1} - 1$ there.) The total number of terms in the sum (6.2.9) is at most $\binom{m-1+(k-k_{m+1})}{m-1}$. So the remaining contribution to (6.2.9) is at most

$$\binom{m-1+(k-k_{m+1})}{m-1}\Gamma((k-k_{m+1}-1)\beta+1)\Gamma(\beta+1).$$

We have found that the left side of (6.2.8) is bounded by

$$\sum_{k_{m+1}=0}^{\infty} \frac{M^{-k_{m+1}}}{k_{m+1}!}\mathbf{1}(k_{m+1} \le k)\, f(k-k_{m+1}),$$

where

$$\begin{aligned} f(k) &\le \frac{m}{(k\beta+1)\cdots(k\beta+m-1)} + \Gamma(\beta+1)\frac{\binom{m-1+k}{m-1}\Gamma((k-1)\beta+1)}{\Gamma(k\beta+m)} && (6.2.10)\\ &\le \frac{\binom{m-1+k}{m-1}\Gamma(k\beta+1)}{\Gamma(k\beta+m)} = \frac{1}{(m-1)!}\frac{(k+1)\cdots(k+(m-1))}{(k\beta+1)\cdots(k\beta+(m-1))}, && (6.2.11) \end{aligned}$$

which is a bounded function of k. To apply the dominated convergence theorem, it suffices to show that the right side of (6.2.10) tends to 0 as $k \to \infty$, which follows from Stirling's approximation and the fact that $\beta > 0$. □

When t_n satisfies the conditions in Theorem 6.2.2 but $\alpha \ge 1+\beta$ then [compare (6.1.24)],

$$\begin{aligned} \tilde{\mu}_n = c_2\frac{n^{\lambda_2-1}}{\Gamma(\lambda_2)} + c_3\frac{n^{\lambda_3-1}}{\Gamma(\lambda_3)} + m!\sum_{j=1}^{m-1}\frac{1}{\psi'(\lambda_j)}[z^n]\left((1-z)^{-\lambda_j}\int_{\zeta=0}^{z}\widehat{T}(\zeta)(1-\zeta)^{\lambda_j-1}\,d\zeta\right)\\ + o(n^{\beta}) + O(n^{\Re(\lambda_4-1)}), \end{aligned}$$

and typically this leads to periodicity.

It is known that $\alpha < 3/2$ for $m \le 26$. Computations show that α increases with m, at least for $m \le 10000$ [51]. This suggests that for a fixed β, the condition $\alpha < 1+\beta$ (resp. $\alpha \ge 1+\beta$) is equivalent to $m \le m_0$ (resp. $m > m_0$) for some $m_0 > 26$.

Large toll functions

If $t_n \sim n^{\beta}$, where $\beta > 1$, then we have convergence in distribution for all values of m. We state the result, omitting the proof, as it is very similar to that of Theorem 6.2.2.

Theorem 6.2.4. *If the real toll sequence* (t_n) *satisfies*

$$t_n \sim n^{\beta}, \qquad \beta > 1,$$

then

$$\frac{X_n}{n^{\beta}} \xrightarrow{\mathcal{L}} Y_{\beta}$$

with convergence of all moments, where $\mathcal{L}(Y_{\beta})$ *is the unique distribution satisfying* (6.2.1).

The borderline case $t_n \sim n$ [more specifically, $t_n \equiv n-(m-1)$] corresponds to the well-studied total path length of a random m-ary search tree. The corresponding additive functional measures the number of key comparisons in m-ary Quicksort. As is well known [63], the number of key comparisons has mean $\Theta(n \ln n)$ and standard deviation $\Theta(n)$. See [36] for details in this case.

Chapter 7

Catalan model: the toll function n^α

We will now use the extended singularity analysis toolkit from Chapter 4 to derive limiting distributions for the additive functionals on binary search trees under the uniform model for the class of toll functions of the form n^α where $\alpha > 0$.

7.1 Preliminaries

In the sequel the notation $[\cdots]$ is used both for Iverson's convention [43, see 1.2.3(16)] and the coefficient of certain terms in the succeeding expression. The interpretation will be clear from the context. For example, $[\alpha > 0]$ has the value 1 when $\alpha > 0$ and the value 0 otherwise. In contrast, $[z^n]F(z)$ denotes the coefficient of z^n in the series expansion of $F(z)$. For notational convenience we define $L(z) := \ln((1-z)^{-1})$. Unless otherwise noted, henceforth all our singular expansions are valid as $z \to 1$ in the sector $-\pi + \epsilon < \arg(1-z) < \pi - \epsilon$, for any $\epsilon > 0$.

We will make extensive use of the following consequences of the singular expansion of the generalized polylogarithm. Note that neither this lemma nor the ones following make any claims about uniformity in α or r.

Lemma 7.1.1. *For real $\alpha < 1$ and r a nonnegative integer,*

$$\operatorname{Li}_{\alpha,r}(z) = \sum_{k=0}^{r} \lambda_k^{(\alpha,r)} (1-z)^{\alpha-1} L^{r-k}(z) + O(|1-z|^{\alpha-\epsilon}) + (-1)^r \zeta^{(r)}(\alpha)[\alpha > 0],$$

where $\lambda_k^{(\alpha,r)} \equiv \binom{r}{k}\Gamma^{(k)}(1-\alpha)$ and $\epsilon > 0$ is arbitrarily small.

Proof. By Theorem 4.1.3,

$$\operatorname{Li}_{\alpha,0}(z) \sim \Gamma(1-\alpha)t^{\alpha-1} + \sum_{j\geq 0} \frac{(-1)^j}{j!}\zeta(\alpha-j)t^j, \qquad t = -\ln z = \sum_{l=1}^{\infty} \frac{(1-z)^l}{l}, \tag{7.1.1}$$

and for r a positive integer,

$$\operatorname{Li}_{\alpha,r}(z) = (-1)^r \frac{\partial^r}{\partial \alpha^r} \operatorname{Li}_{\alpha,0}(z).$$

Moreover, the singular expansion for $\operatorname{Li}_{\alpha,r}$ is obtained by performing the indicated differentiation of (7.1.1) term-by-term. To establish the claim we set $f = \Gamma(1-\alpha)$ and $g = t^{\alpha-1}$ in the general formula for the rth derivative of a product:

$$(fg)^{(r)} = \sum_{k=0}^{r} \binom{r}{k} f^{(k)} g^{(r-k)}$$

to first obtain

$$(-1)^r \frac{\partial^r}{\partial \alpha^r}[\Gamma(1-\alpha)t^{\alpha-1}] = (-1)^r \sum_{k=0}^{r} \binom{r}{k} (-1)^k \Gamma^{(k)}(1-\alpha) t^{\alpha-1} (\ln t)^{r-k}$$

The claim then follows easily. ☐

The following “inverse” of Lemma 7.1.1 is very useful for computing with Hadamard products.

Lemma 7.1.2. *For real* $\alpha < 1$ *and* r *a nonnegative integer,*

$$(1-z)^{\alpha-1}L^r(z) = \sum_{k=0}^{r} \mu_k^{(\alpha,r)} \operatorname{Li}_{\alpha,r-k}(z) + O(|1-z|^{\alpha-\epsilon}) + c_r(\alpha)[\alpha>0],$$

where $\mu_0^{(\alpha,r)} = 1/\Gamma(1-\alpha)$, $c_r(\alpha)$ *is a constant, and* $\epsilon > 0$ *is arbitrarily small.*

Proof. We use induction on r. For $r=0$ we have

$$\operatorname{Li}_{\alpha,0}(z) = \Gamma(1-\alpha)(1-z)^{\alpha-1} + O(|1-z|^{\alpha-\epsilon}) + \zeta(\alpha)[\alpha>0]$$

and the claim is verified with

$$\mu_0^{(\alpha,0)} = \frac{1}{\Gamma(1-\alpha)} \qquad \text{and} \qquad c_0(\alpha) = -\frac{\zeta(\alpha)}{\Gamma(1-\alpha)}.$$

Let $r \geq 1$. Then using Lemma 7.1.1 and the induction hypothesis we get

$$\begin{aligned}
\operatorname{Li}_{\alpha,r}(z) &= \Gamma(1-\alpha)(1-z)^{\alpha-1}L^r(z) \\
&\quad + \sum_{k=1}^{r} \lambda_k^{(\alpha,r)} \left[\sum_{l=0}^{r-k} \mu_l^{(\alpha,r-k)} \operatorname{Li}_{\alpha,r-k-l}(z) + O(|1-z|^{\alpha-\epsilon}) + c_{r-k}(\alpha)[\alpha>0]\right] \\
&\quad + O(|1-z|^{\alpha-\epsilon}) + (-1)^r \zeta^{(r)}(\alpha)[\alpha>0] \\
&= \Gamma(1-\alpha)(1-z)^{\alpha-1}L^r(z) + \sum_{k=1}^{r} \lambda_k^{(\alpha,r)} \sum_{s=0}^{r-k} \mu_{r-k-s}^{(\alpha,r-k)} \operatorname{Li}_{\alpha,s}(z) + O(|1-z|^{\alpha-\epsilon}) \\
&\quad + \left(\sum_{k=1}^{r} \lambda_k^{(\alpha,r)} c_{r-k}(\alpha) + (-1)^r \zeta^{(r)}(\alpha)\right)[\alpha>0] \\
&= \Gamma(1-\alpha)(1-z)^{\alpha-1}L^r(z) + \sum_{s=0}^{r-1} \nu_s^{(\alpha,r)} \operatorname{Li}_{\alpha,s}(z) + O(|1-z|^{\alpha-\epsilon}) + \gamma_r(\alpha)[\alpha>0],
\end{aligned}$$

where, for $0 \leq s \leq r-1$,

$$\nu_s^{(\alpha,r)} := \sum_{k=1}^{r-s} \lambda_k^{(\alpha,r)} \mu_{r-s-k}^{(\alpha,r-k)},$$

and where

$$\gamma_r(\alpha) := \sum_{k=1}^{r} \lambda_k^{(\alpha,r)} c_{r-k}(\alpha) + (-1)^r \zeta^{(r)}(\alpha).$$

Setting

$$\mu_0^{(\alpha,r)} = \frac{1}{\Gamma(1-\alpha)}, \qquad \mu_k^{(\alpha,r)} = -\frac{\nu_{r-k}^{(\alpha,r)}}{\Gamma(1-\alpha)}, \quad 1 \leq k \leq r,$$

and

$$c_r(\alpha) = -\frac{\gamma_r(\alpha)}{\Gamma(1-\alpha)},$$

the result follows. ☐

For the calculation of the mean, the following refinement of a special case of Lemma 7.1.1 is required. It is a simple consequence of Theorem 4.1.3.

Lemma 7.1.3. *When* $\alpha < 0$,

$$\mathrm{Li}_{\alpha,0}(z) = \Gamma(1-\alpha)(1-z)^{\alpha-1} - \Gamma(1-\alpha)\frac{1-\alpha}{2}(1-z)^{\alpha} + O(|1-z|^{\alpha+1}) + \zeta(\alpha)[\alpha > -1].$$

7.2 Asymptotics of the mean

We start by reviewing some notation. Let β_n denote the nth Catalan number

$$\beta_n := \frac{1}{n+1}\binom{2n}{n},$$

with generating function

$$\mathrm{CAT}(z) := \sum_{n=0}^{\infty} \beta_n z^n = \frac{1}{2z}(1-\sqrt{1-4z}).$$

In our subsequent analysis we will make use of the identity

$$z\,\mathrm{CAT}^2(z) = \mathrm{CAT}(z) - 1. \tag{7.2.1}$$

The mean of the cost function $a_n := \mathbf{E}\, X_n$ can be obtained recursively by conditioning on the key stored at the root as

$$a_n = \sum_{j=1}^{n} \frac{\beta_{j-1}\beta_{n-j}}{\beta_n}(a_{j-1} + a_{n-j}) + t_n, \qquad n \geq 1.$$

This recurrence can be rewritten as

$$(\beta_n a_n) = 2\sum_{j=1}^{n} (\beta_{j-1}a_{j-1})\beta_{n-j} + (\beta_n t_n), \qquad n \geq 1. \tag{7.2.2}$$

Multiplying (7.2.2) by $z^n/4^n$ and summing over $n \geq 1$ we get

$$A(z) \odot \mathrm{CAT}(z/4) = \frac{T(z) \odot \mathrm{CAT}(z/4)}{\sqrt{1-z}} + \frac{a_0 - t_0}{\sqrt{1-z}}, \tag{7.2.3}$$

where $A(z)$ and $T(z)$ are the ordinary generating functions of (a_n) and (t_n) respectively. In our analysis we will assume $a_0 = t_0$ so that the second term in (7.2.3) does not contribute. (Our results can be easily adjusted when this is not the case.) We will use (7.2.3) (with $a_0 = t_0$) as the starting point for our analysis.

Since $t_n = n^{\alpha}$, by definition $T = \mathrm{Li}_{-\alpha,0}$. Thus, by Lemma 7.1.3,

$$T(z) = \Gamma(1+\alpha)(1-z)^{-\alpha-1} - \Gamma(1+\alpha)\frac{\alpha+1}{2}(1-z)^{-\alpha} + O(|1-z|^{-\alpha+1}) + \zeta(-\alpha)[\alpha < 1].$$

We will now use (7.2.3) to obtain the asymptotics of the mean.

First we treat the case $\alpha < 1/2$. Since $\mathrm{CAT}(z/4) = 2 + O(|1-z|^{1/2})$ as $z \to 1$ in a suitable indented crown, we have, by Theorem 4.3.5(b),

$$T(z) \odot \mathrm{CAT}(z/4) = C_0 + O(|1-z|^{-\alpha+\frac{1}{2}}),$$

where

$$C_0 := T(z) \odot \mathrm{CAT}(z/4)\Big|_{z=1} = \sum_{n=1}^{\infty} n^{\alpha} \frac{\beta_n}{4^n}.$$

We now already know the constant term in the singular expansion of $T(z) \odot \mathrm{CAT}(z/4)$ at $z = 1$ and henceforth we need only compute lower-order terms. The constant $\bar{c}$ is used in the sequel to denote an unspecified (possibly 0) constant, possibly different at each appearance.

Let us write $T(z) = L_1(z) + R_1(z)$, and $\mathrm{CAT}(z/4) = L_2(z) + R_2(z)$, where

$$\begin{aligned} L_1(z) &:= \Gamma(1+\alpha)(1-z)^{-\alpha-1} - \Gamma(1+\alpha)\frac{\alpha+1}{2}(1-z)^{-\alpha} + \zeta(-\alpha), \\ R_1(z) &:= T(z) - L_1(z) = O(|1-z|^{1-\alpha}), \\ L_2(z) &:= 2(1-(1-z)^{1/2}), \\ R_2(z) &:= \mathrm{CAT}(z/4) - L_2(z) = O(|1-z|). \end{aligned}$$

We will analyze each of the four Hadamard products separately. First,

$$L_1(z) \odot L_2(z) = -2\Gamma(1+\alpha)\big[(1-z)^{-\alpha-1} \odot (1-z)^{1/2}\big] + 2\Gamma(1+\alpha)\frac{\alpha+1}{2}\big[(1-z)^{-\alpha} \odot (1-z)^{1/2}\big] + \bar{c}.$$

By Theorem 4.3.5,

$$(1-z)^{-\alpha-1} \odot (1-z)^{1/2} = \bar{c} + \frac{\Gamma(\alpha-\frac{1}{2})}{\Gamma(\alpha+1)\Gamma(-1/2)}(1-z)^{-\alpha+\frac{1}{2}} + O(|1-z|),$$

and

$$(1-z)^{-\alpha} \odot (1-z)^{1/2} = \bar{c} + O(|1-z|)$$

by another application of Theorem 4.3.5(b), this time with $k = 1$. Hence

$$L_1(z) \odot L_2(z) = \big[L_1(z) \odot L_2(z)\big]\Big|_{z=1} + \frac{\Gamma(\alpha-\frac{1}{2})}{\sqrt{\pi}}(1-z)^{-\alpha+\frac{1}{2}} + O(|1-z|).$$

The other three Hadamard products are easily handled as

$$\begin{aligned} L_1(z) \odot R_2(z) &= \big[L_1(z) \odot R_2(z)\big]\Big|_{z=1} + O(|1-z|^{-\alpha+1}), \\ L_2(z) \odot R_1(z) &= \big[L_2(z) \odot R_1(z)\big]\Big|_{z=1} + O(|1-z|), \\ R_1(z) \odot R_2(z) &= \big[R_1(z) \odot R_2(z)\big]\Big|_{z=1} + O(|1-z|). \end{aligned}$$

Putting everything together, we get

$$T(z) \odot \mathrm{CAT}(z/4) = C_0 + \frac{\Gamma(\alpha-\frac{1}{2})}{\sqrt{\pi}}(1-z)^{-\alpha+\frac{1}{2}} + O(|1-z|^{-\alpha+1}).$$

Using this in (7.2.3), we get

$$A(z) \odot \mathrm{CAT}(z/4) = C_0(1-z)^{-1/2} + \frac{\Gamma(\alpha-\frac{1}{2})}{\sqrt{\pi}}(1-z)^{-\alpha} + O(|1-z|^{-\alpha+\frac{1}{2}}). \tag{7.2.4}$$

To treat the case $\alpha \geq 1/2$ we make use of the estimate

$$(1-z)^{1/2} = \frac{1}{\Gamma(-1/2)}[\mathrm{Li}_{3/2,0}(z) - \zeta(3/2)] + O(|1-z|), \tag{7.2.5}$$

a consequence of Theorem 4.1.3, so that

$$T(z) \odot (1-z)^{1/2} = \mathrm{Li}_{-\alpha,0}(z) \odot (1-z)^{1/2} = \frac{1}{\Gamma(-1/2)} \mathrm{Li}_{\frac{3}{2}-\alpha,0}(z) + R(z),$$

where

$$R(z) = \begin{cases} \bar{c} + O(|1-z|^{1-\alpha}) & 1/2 \leq \alpha < 1 \\ O(|L(z)|) & \alpha = 1 \\ O(|1-z|^{1-\alpha}) & \alpha > 1. \end{cases} \tag{7.2.6}$$

Hence

$$T(z) \odot \mathrm{CAT}(z/4) = -\frac{2}{\Gamma(-1/2)} \mathrm{Li}_{\frac{3}{2}-\alpha,0}(z) + \widetilde{R}(z),$$

where $\widetilde{R}$, like R, satisfies (7.2.6) (with a possibly different $\bar{c}$). When $\alpha = 1/2$, this gives us

$$T(z) \odot \mathrm{CAT}(z/4) = -\frac{2}{\Gamma(-1/2)} L(z) + \bar{c} + O(|1-z|^{1/2}),$$

so that

$$A(z) \odot \mathrm{CAT}(z/4) = \frac{1}{\sqrt{\pi}}(1-z)^{-1/2} L(z) + \bar{c}(1-z)^{-1/2} + O(1). \tag{7.2.7}$$

For $\alpha > 1/2$ another singular expansion leads to the conclusion that

$$A(z) \odot \mathrm{CAT}(z/4) = \frac{\Gamma(\alpha - \frac{1}{2})}{\sqrt{\pi}}(1-z)^{-\alpha} + \widehat{R}(z), \tag{7.2.8}$$

where

$$\widehat{R}(z) = \begin{cases} O(|1-z|^{-\frac{1}{2}}) & 1/2 < \alpha < 1 \\ O(|1-z|^{-\frac{1}{2}}|L(z)|) & \alpha = 1 \\ O(|1-z|^{-\alpha+\frac{1}{2}}) & \alpha > 1. \end{cases}$$

We defer deriving the asymptotics of a_n until Section 7.4.

7.3 Higher moments

We will analyze separately the cases $0 < \alpha < 1/2$, $\alpha = 1/2$, and $\alpha > 1/2$. The reason for this will become evident soon; though the technique used to derive the asymptotics is induction in each case, the induction hypothesis is different for each of these cases.

Small toll functions $(0 < \alpha < 1/2)$

We start by restricting ourselves to tolls of the form n^α where $0 < \alpha < 1/2$. In this case we observe that by singularity analysis applied to (7.2.4),

$$\frac{a_n \beta_n}{4^n} = \frac{C_0}{\sqrt{\pi}} n^{-1/2} + O(n^{-3/2}) + O(n^{\alpha-1}) = \frac{C_0}{\sqrt{\pi}} n^{-1/2} + O(n^{\alpha-1}),$$

so

$$a_n = n^{\frac{3}{2}}[1 + O(n^{-1})][C_0 n^{-\frac{1}{2}} + O(n^{\alpha-1})] = C_0 n + O(n^{\alpha+\frac{1}{2}}) = (C_0 + o(1))(n+1).$$

The lead order of the mean $a_n = \mathbf{E}\, X_n$ does not depend on α. We perform an approximate centering to get to the dependence on α.

Define $\widetilde{X}_n := X_n - C_0(n+1)$, with $X_0 := 0$; $\tilde{\mu}_n(k) := \mathbf{E}\, \widetilde{X}_n^k$, with $\tilde{\mu}_n(0) = 1$ for all $n \geq 0$; and $\hat{\mu}_n(k) := \beta_n \tilde{\mu}_n(k)/4^n$. Let $\widehat{M}_k(z)$ denote the ordinary generating function of $\hat{\mu}_n(k)$ in the argument n.

By an argument similar to the one that led to (7.2.2), we get, for $k \geq 2$,

$$\hat{\mu}_n(k) = \frac{1}{2} \sum_{j=1}^{n} \frac{\beta_{n-j}}{4^{n-j}} \hat{\mu}_{j-1}(k) + \hat{r}_n(k), \qquad n \geq 1,$$

where

$$\begin{aligned} \hat{r}_n(k) &:= \frac{1}{4} \sum_{j=1}^{n} \sum_{\substack{k_1+k_2+k_3=k \\ k_1,k_2<k}} \binom{k}{k_1, k_2, k_3} \hat{\mu}_{j-1}(k_1) \hat{\mu}_{n-j}(k_2) t_n^{k_3} \\ &= \frac{1}{4} \sum_{\substack{k_1+k_2+k_3=k \\ k_1,k_2<k}} \binom{k}{k_1, k_2, k_3} t_n^{k_3} \sum_{j=1}^{n} \hat{\mu}_{j-1}(k_1) \hat{\mu}_{n-j}(k_2), \end{aligned}$$

for $n \geq 1$ and $\hat{r}_0(k) := \hat{\mu}_0(k) = \tilde{\mu}_0(k) = (-1)^k C_0^k$. Let $\widehat{R}_k(z)$ denote the ordinary generating function of $\hat{r}_n(k)$ in the argument n. Then, mimicking (7.2.3),

$$\widehat{M}_k(z) = \frac{\widehat{R}_k(z)}{\sqrt{1-z}} \tag{7.3.1}$$

with

$$\widehat{R}_k(z) = (-1)^k C_0^k + \sum_{\substack{k_1+k_2+k_3=k \\ k_1,k_2<k}} \binom{k}{k_1, k_2, k_3} \big(T(z)^{\odot k_3}\big) \odot \big[\frac{z}{4} \widehat{M}_{k_1}(z) \widehat{M}_{k_2}(z)\big], \tag{7.3.2}$$

where for k a nonnegative integer

$$T(z)^{\odot k} := \underbrace{T(z) \odot \cdots \odot T(z)}_{k}.$$

Note that $\widehat{M}_0(z) = \mathrm{CAT}(z/4)$.

We analyze separately the cases $0 < \alpha \leq 1/4$ and $1/4 < \alpha < 1/2$. The proof technique in either case is induction, though the latter case is slightly simpler.

Proposition 7.3.1. *In the notation introduced above, when $0 < \alpha \leq 1/4$, for $k \geq 1$,*

$$\widehat{M}_k(z) = C_k (1-z)^{-k(\alpha+\frac{1}{2})+\frac{1}{2}} + O(|1-z|^{-k(\alpha+\frac{1}{2})+\frac{1}{2}+(2\alpha-\epsilon)}),$$

as $z \to 1$ in the sector $-\pi + \epsilon' < \arg(1-z) < \pi - \epsilon'$, where $\epsilon > 0$ and $\epsilon' > 0$ are arbitrarily *small. The C_k's here are defined by the recurrence*

$$C_k = \frac{1}{4}\sum_{j=1}^{k-1}\binom{k}{j}C_jC_{k-j} + kC_{k-1}\frac{\Gamma(k\alpha+\frac{k}{2}-1)}{\Gamma((k-1)\alpha+\frac{k}{2}-1)}, \quad k \geq 2; \qquad C_1 = \frac{\Gamma(\alpha-\frac{1}{2})}{\sqrt{\pi}}. \quad (7.3.3)$$

Proof. For $k=1$ the claim is true as shown in (7.2.4) with C_1 as defined in (7.3.3). We will now analyze each term in (7.3.2) for $k \geq 2$. For notational convenience, define $\alpha' := \alpha + \frac{1}{2}$. Also, observe that

$$T(z)^{\odot k} = \mathrm{Li}_{-k\alpha,0}(z) = \Gamma(1+k\alpha)(1-z)^{-k\alpha-1} + O(|1-z|^{-k\alpha-\epsilon})$$

by Lemma 7.1.1.

For this paragraph, consider the case that k_1 and k_2 are both nonzero. It follows from the induction hypothesis that

$$\begin{aligned}\frac{z}{4}\widehat{M}_{k_1}(z)\widehat{M}_{k_2}(z) &= \frac{1}{4}(1-(1-z))\left[C_{k_1}(1-z)^{-k_1\alpha'+\frac{1}{2}} + O(|1-z|^{-k_1\alpha'+\frac{1}{2}+(2\alpha-\epsilon)})\right] \\ &\quad \times \left[C_{k_2}(1-z)^{-k_2\alpha'+\frac{1}{2}} + O(|1-z|^{-k_2\alpha'+\frac{1}{2}+(2\alpha-\epsilon)})\right] \\ &= \frac{1}{4}C_{k_1}C_{k_2}(1-z)^{-(k_1+k_2)\alpha'+1} + O(|1-z|^{-(k_1+k_2)\alpha'+1+(2\alpha-\epsilon)}).\end{aligned}$$

If $k_3 = 0$ then the corresponding contribution to $\widehat{R}_k(z)$ is

$$\frac{1}{4}\binom{k}{k_1}C_{k_1}C_{k_2}(1-z)^{-k\alpha'+1} + O(|1-z|^{-k\alpha'+1+(2\alpha-\epsilon)}).$$

If $k_3 \neq 0$ we use Lemma 7.1.2 to express

$$\begin{aligned}\frac{z}{4}\widehat{M}_{k_1}(z)\widehat{M}_{k_2}(z) &= \frac{C_{k_1}C_{k_2}}{4\Gamma((k_1+k_2)\alpha'-1)}\,\mathrm{Li}_{-(k_1+k_2)\alpha'+2,0}(z) \\ &\quad + O(|1-z|^{-(k_1+k_2)\alpha'+1+(2\alpha-\epsilon)}) - \frac{C_{k_1}C_{k_2}}{4}[(k_1+k_2)\alpha' < 2]\frac{\zeta(-(k_1+k_2)\alpha'+2)}{\Gamma((k_1+k_2)\alpha'-1)}.\end{aligned}$$

The corresponding contribution to $\widehat{R}_k(z)$ is then $\binom{k}{k_1,k_2,k_3}$ times:

$$\frac{C_{k_1}C_{k_2}}{4\Gamma((k_1+k_2)\alpha'-1)}\,\mathrm{Li}_{-k\alpha'+\frac{k_3}{2}+2,0}(z) + \mathrm{Li}_{-k_3\alpha,0}(z) \odot O(|1-z|^{-(k_1+k_2)\alpha'+1+(2\alpha-\epsilon)}).$$

Now $k_3 \leq k-2$ so $-k\alpha' + \frac{k_3}{2} + 2 < 1$. Hence the contribution when $k_3 \neq 0$ is

$$O(|1-z|^{-k\alpha'+\frac{k_3}{2}+1}) = O(|1-z|^{-k\alpha'+\frac{3}{2}}) = O(|1-z|^{-k\alpha'+1+(2\alpha-\epsilon)}).$$

Next we consider the case when k_1 is nonzero but $k_2 = 0$. In this case using the induction hypothesis we see that

$$\begin{aligned}\frac{z}{4}\widehat{M}_{k_1}(z)\widehat{M}_{k_2}(z) &= \frac{z}{4}\,\mathrm{CAT}(z/4)\widehat{M}_{k_1}(z) \\ &= \frac{1-(1-z)^{1/2}}{2}\left[C_{k_1}(1-z)^{-k_1\alpha'+\frac{1}{2}}\right] + O(|1-z|^{-k_1\alpha'+\frac{1}{2}+(2\alpha-\epsilon)}) \\ &= \frac{C_{k_1}}{2}(1-z)^{-k_1\alpha'+\frac{1}{2}} + O(|1-z|^{-k_1\alpha'+\frac{1}{2}+(2\alpha-\epsilon)}).\end{aligned}$$

Applying Lemma 7.1.2 to the last expression we get

$$\frac{z}{4}\widehat{M}_{k_1}(z)\widehat{M}_{k_2}(z) = \frac{C_{k_1}}{2\Gamma(k_1\alpha' - \frac{1}{2})}\operatorname{Li}_{-k_1\alpha'+\frac{3}{2},0}(z) \\ + O(|1-z|^{-k_1\alpha'+\frac{1}{2}+(2\alpha-\epsilon)}) - \frac{C_{k_1}}{2}[k_1\alpha' - \tfrac{1}{2} < 1]\frac{\zeta(-k_1\alpha'+\frac{3}{2})}{\Gamma(k_1\alpha'-\frac{1}{2})}.$$

The contribution to $\widehat{R}_k(z)$ is hence $\binom{k}{k_1}$ times:

$$\frac{C_{k_1}}{2\Gamma(k_1\alpha' - \frac{1}{2})}\operatorname{Li}_{-k\alpha'+\frac{k_3}{2}+\frac{3}{2},0}(z) + \operatorname{Li}_{-k_3\alpha,0}(z) \odot O(|1-z|^{-k_1\alpha'+\frac{1}{2}+(2\alpha-\epsilon)}).$$

Using the fact that $\alpha > 0$ and $k_3 \leq k-1$, we conclude that $-k\alpha' + \frac{k_3}{2} + \frac{3}{2} < 1$ so that, by Lemma 7.1.1 and Theorem 4.3.5(a), the contribution is

$$O(|1-z|^{-k\alpha'+\frac{k_3}{2}+\frac{1}{2}}) = O(|1-z|^{-k\alpha'+\frac{3}{2}})$$

where the displayed equality holds unless $k_3 = 1$. When $k_3 = 1$ we get a corresponding contribution to $\widehat{R}_k(z)$ of $\binom{k}{k-1}$ times:

$$\frac{C_{k-1}\Gamma(k\alpha'-1)}{2\Gamma((k-1)\alpha'-\frac{1}{2})}(1-z)^{-k\alpha'+1} + O(|1-z|^{-k\alpha'+1+(2\alpha-\epsilon)}),$$

since for $k \geq 2$ we have $k\alpha' > 1 + (2\alpha - \epsilon)$. The introduction of ϵ handles the case when $k\alpha' = 1 + 2\alpha$, which would have otherwise, according to Theorem 4.3.5(c) introduced a logarithmic remainder. In either case the remainder is $O(|1-z|^{-k\alpha'+1+(2\alpha-\epsilon)})$. The case when k_2 is nonzero but $k_1 = 0$ is handled similarly by exchanging the roles of k_1 and k_2.

The final contribution comes from the single term where both k_1 and k_2 are zero. In this case the contribution to $\widehat{R}_k(z)$ is, recalling (7.2.1),

$$\operatorname{Li}_{-k\alpha,0}(z) \odot [\frac{z}{4}\operatorname{CAT}^2(z/4)] = \operatorname{Li}_{-k\alpha,0}(z) \odot (\operatorname{CAT}(z/4) - 1) = \operatorname{Li}_{-k\alpha,0}(z) \odot \operatorname{CAT}(z/4). \quad (7.3.4)$$

Now, using Theorem 4.1.3,

$$\operatorname{CAT}(z/4) = 2 - 2(1-z)^{1/2} + O(|1-z|) = 2 + 2\frac{\zeta(3/2)}{\Gamma(-1/2)} - \frac{2}{\Gamma(-1/2)}\operatorname{Li}_{3/2,0}(z) + O(|1-z|),$$

so that (7.3.4) is

$$-\frac{2}{\Gamma(-1/2)}\operatorname{Li}_{\frac{3}{2}-k\alpha,0}(z) + O(|1-z|^{1-k\alpha}) + \begin{cases} 0 & 1-k\alpha < 0, \\ O(|1-z|^{-\epsilon}) & 1-k\alpha = 0, \\ O(1) & 1-k\alpha > 0. \end{cases}$$

When $\frac{3}{2} - k\alpha < 1$ this is $O(|1-z|^{-k\alpha+\frac{1}{2}})$; when $\frac{3}{2} - k\alpha \geq 1$, it is $O(1)$. In either case we get a contribution which is $O(|1-z|^{-k\alpha'+1+(2\alpha-\epsilon)})$.

Hence

$$\begin{aligned}\widehat{R}_k(z) &= O(1) + \left[\sum_{\substack{k_1+k_2=k\\ k_1,k_2<k}} \binom{k}{k_1}\frac{C_{k_1}C_{k_2}}{4} + 2k\frac{C_{k-1}}{2}\frac{\Gamma(k\alpha+\frac{k}{2}-1)}{\Gamma((k-1)\alpha+\frac{k}{2}-1)}\right](1-z)^{-k\alpha'+1}\\ &\quad + O(|1-z|^{-k\alpha'+1+(2\alpha-\epsilon)})\\ &= C_k(1-z)^{-k\alpha'+1} + O(|1-z|^{-k\alpha'+1+(2\alpha-\epsilon)}),\end{aligned}$$

with the C_k's defined by the recurrence (7.3.3). Now using (7.3.1), the claim follows. ☐

Next, we handle the case when $1/4 < \alpha < 1/2$.

Proposition 7.3.2. *In the notation introduced above Proposition 7.3.1, when* $1/4 < \alpha < 1/2$,

$$\widehat{M}_k(z) = C_k(1-z)^{-k(\alpha+\frac{1}{2})+\frac{1}{2}} + O(|1-z|^{-k(\alpha+\frac{1}{2})+1}),$$

for $k \geq 1$, *where the* C_k*'s are defined by the recurrence* (7.3.3).

Proof. For $k = 1$ the claim is true as shown in Section 7.2 with C_1 defined at (7.3.3). We will now analyze each term in (7.3.2) for $k \geq 2$ in a manner similar to that in the proof of Proposition 7.3.1.

First we consider the case that k_1 and k_2 are both nonzero. It follows from the induction hypothesis that

$$\frac{z}{4}\widehat{M}_{k_1}(z)\widehat{M}_{k_2}(z) = \frac{C_{k_1}C_{k_2}}{4}(1-z)^{-(k_1+k_2)\alpha'+1} + O(|1-z|^{-(k_1+k_2)\alpha'+\frac{3}{2}}).$$

If $k_3 = 0$ then the corresponding contribution to $\widehat{R}_k(z)$ is

$$\frac{1}{4}\binom{k}{k_1}C_{k_1}C_{k_2}(1-z)^{-k\alpha'+1} + O(|1-z|^{-k\alpha'+\frac{3}{2}}).$$

If $k_3 \neq 0$ we use Lemma 7.1.2 to express

$$\begin{aligned}\frac{z}{4}\widehat{M}_{k_1}(z)\widehat{M}_{k_2}(z) &= \frac{C_{k_1}C_{k_2}}{4\Gamma((k_1+k_2)\alpha'-1)}\operatorname{Li}_{-(k_1+k_2)\alpha'+2,0}(z)\\ &\quad + O(|1-z|^{-(k_1+k_2)\alpha'+\frac{3}{2}}) - \frac{C_{k_1}C_{k_2}}{4}[(k_1+k_2)\alpha' < 2]\frac{\zeta(-(k_1+k_2)\alpha'+2)}{\Gamma((k_1+k_2)\alpha'-1)}.\end{aligned}$$

The contribution to $\widehat{R}_k(z)$ is then $\binom{k}{k_1,k_2,k_3}$ times:

$$\frac{C_{k_1}C_{k_2}}{4\Gamma((k_1+k_2)\alpha'-1)}\operatorname{Li}_{-k\alpha'+\frac{k_3}{2}+2,0}(z) + \operatorname{Li}_{-k_3\alpha,0}(z) \odot O(|1-z|^{-(k_1+k_2)\alpha'+\frac{3}{2}}).$$

Now $k_3 \leq k-2$ so $-k\alpha' + \frac{k_3}{2} + 2 < 1$. Hence the contribution when $k_3 \neq 0$ is

$$O(|1-z|^{-k\alpha'+\frac{k_3}{2}+1}) = O(|1-z|^{-k\alpha'+\frac{3}{2}})$$

by another application of Theorem 4.3.5(a).

Next we consider the case when k_1 is nonzero but $k_2 = 0$. In this case using the induction hypothesis we see that

$$\frac{z}{4}\widehat{M}_{k_1}(z)\widehat{M}_{k_2}(z) = \frac{z}{4}\operatorname{CAT}(z/4)\widehat{M}_{k_1}(z) = \frac{C_{k_1}}{2}(1-z)^{-k_1\alpha'+\frac{1}{2}} + O(|1-z|^{-k_1\alpha'+1}).$$

Applying Lemma 7.1.2 to the last expression we get

$$\frac{z}{4}\widehat{M}_{k_1}(z)\widehat{M}_{k_2}(z) = \frac{C_{k_1}}{2\Gamma(k_1\alpha' - \frac{1}{2})}\operatorname{Li}_{-k_1\alpha'+\frac{3}{2},0}(z) + O(|1-z|^{-k_1\alpha'+1}) - \frac{C_{k_1}}{2}[k_1\alpha' - \tfrac{1}{2} < 1]\frac{\zeta(-k_1\alpha' + \frac{3}{2})}{\Gamma(k_1\alpha' - \frac{1}{2})}$$

The contribution to $\widehat{R}_k(z)$ is hence $\binom{k}{k_1}$ times:

$$\frac{C_{k_1}}{2\Gamma(k_1\alpha' - \frac{1}{2})}\operatorname{Li}_{-k\alpha'+\frac{k_3}{2}+\frac{3}{2},0}(z) + \operatorname{Li}_{-k_3\alpha,0}(z) \odot O(|1-z|^{-k_1\alpha'+1}).$$

Using the fact that $\alpha > 1/4$ and $k_3 \le k-1$, we conclude that $-k\alpha' + \frac{k_3}{2} + \frac{3}{2} < \frac{1}{2}$ so that the contribution is

$$O(|1-z|^{-k\alpha'+\frac{k_3}{2}+\frac{1}{2}}) = O(|1-z|^{-k\alpha'+\frac{3}{2}})$$

by Theorem 4.3.5(a), unless $k_3 = 1$. When $k_3 = 1$ we get a contribution of

$$\binom{k}{k-1}\frac{C_{k-1}\Gamma(k\alpha' - 1)}{2\Gamma((k-1)\alpha' - \frac{1}{2})}(1-z)^{-k\alpha'+1} + O(|1-z|^{-k\alpha'+\frac{3}{2}}),$$

since $k \ge 2$ and $\alpha > 1/4$ imply $k\alpha' > 3/2$. The case when k_2 is nonzero but $k_1 = 0$ is handled similarly.

The final contribution comes from the single term where both k_1 and k_2 are zero. In this case the contribution to $\widehat{R}_k(z)$ is

$$\operatorname{Li}_{-k\alpha,0}(z) \odot [\frac{z}{4}\operatorname{CAT}^2(z/4)] = \operatorname{Li}_{-k\alpha,0}(z) \odot (\operatorname{CAT}(z/4) - 1) = \operatorname{Li}_{-k\alpha,0}(z) \odot \operatorname{CAT}(z/4)$$

which is easily seen to be $O(|1-z|^{-k\alpha'+\frac{3}{2}})$.

Hence, as in the proof of Proposition 7.3.1, we see that

$$\widehat{R}_k(z) = C_k(1-z)^{-k(\alpha+\frac{1}{2})+1} + O(|1-z|^{-k(\alpha+\frac{1}{2})+\frac{3}{2}})$$

with the C_k's defined by the recurrence (7.3.3), and the claim follows using (7.3.1). □

Large toll functions ($\alpha \ge 1/2$)

When $\alpha \ge 1/2$ there is no need to apply the centering techinques. Define $\mu_n(k) := \mathbf{E}\,X_n^k$ and $\bar{\mu}_n(k) := \beta_n\mu_n(k)/4^n$. Let $\overline{M}_k(z)$ denote the ordinary generating function of $\bar{\mu}_n(k)$ in n. Observe that $\overline{M}_0(z) = \operatorname{CAT}(z/4)$. As earlier, conditioning on the key stored at the root, we get, for $k \ge 2$,

$$\bar{\mu}_n(k) = \frac{1}{2}\sum_{j=1}^{n}\frac{\beta_{n-j}}{4^{n-j}}\bar{\mu}_{j-1}(k) + \bar{r}_n(k), \qquad n \ge 1,$$

where

$$\bar{r}_n(k) := \frac{1}{4}\sum_{\substack{k_1+k_2+k_3=k\\ k_1,k_2<k}}\binom{k}{k_1,k_2,k_3}t_n^{k_3}\sum_{j=1}^{n}\bar{\mu}_{j-1}(k_1)\bar{\mu}_{n-j}(k_2),$$

for $n \geq 1$ and $\bar{r}_0(k) := \bar{\mu}_0(k) = \mu_0(k) = 0$. Let $\overline{R}_k(z)$ denote the ordinary generating function of $\bar{r}_n(k)$ in n. Then

$$\overline{M}_k(z) = \frac{\overline{R}_k(z)}{\sqrt{1-z}}$$

and

$$\overline{R}_k(z) = \sum_{\substack{k_1+k_2+k_3=k \\ k_1,k_2<k}} \binom{k}{k_1, k_2, k_3} \big(T(z)^{\odot k_3}\big) \odot \big[\frac{z}{4}\overline{M}_{k_1}(z)\overline{M}_{k_2}(z)\big]. \tag{7.3.5}$$

We can now state the result about the asymptotics of the generating function $\overline{M}_k$ when $\alpha > 1/2$. The case $\alpha = 1/2$ will be handled subsequently, in Proposition 7.3.4.

Proposition 7.3.3. *Let $\epsilon > 0$ be arbitrary, and define*

$$c := \begin{cases} \alpha - \frac{1}{2} & \frac{1}{2} < \alpha < 1 \\ \frac{1}{2} - \epsilon & \alpha = 1 \\ \frac{1}{2} & \alpha > 1. \end{cases} \tag{7.3.6}$$

Then the generating function $\overline{M}_k(z)$ of $\bar{\mu}_n(k)$ satisfies

$$\overline{M}_k(z) = C_k(1-z)^{-k(\alpha+\frac{1}{2})+\frac{1}{2}} + O(|1-z|^{-k(\alpha+\frac{1}{2})+\frac{1}{2}+c})$$

for $k \geq 1$, where the C_k's are defined by the recurrence (7.3.3).

Proof. The proof is very similar to those of Propositions 7.3.1 and 7.3.2. We present a sketch. The reader is invited to compare the cases enumerated below to those in the earlier proofs.

When $k = 1$ the claim is true by (7.2.8). We analyze the various terms in (7.3.5) for $k \geq 2$, employing the notational convenience $\alpha' := \alpha + \frac{1}{2}$.

When both k_1 and k_2 are nonzero then the contribution to $\overline{R}_k(z)$ is

$$\frac{1}{4}\binom{k}{k_1} C_{k_1} C_{k_2} (1-z)^{-k\alpha'+1} + O(|1-z|^{-k\alpha'+c+1})$$

when $k_3 = 0$ and is $O(|1-z|^{-k\alpha'+c+1})$ otherwise.

When k_1 is nonzero and $k_2 = 0$ the contribution to $\overline{R}_k(z)$ is

$$k\frac{C_{k-1}\Gamma(k\alpha'-1)}{2\Gamma((k-1)\alpha'-\frac{1}{2})}(1-z)^{-k\alpha'+1} + O(|1-z|^{-k\alpha'+c+1})$$

when $k_3 = 1$ and $O(|1-z|^{-k\alpha'+c+1})$ otherwise. The case when k_2 is nonzero and $k_1 = 0$ is identical.

The final contribution comes from the single term when both k_1 and k_2 are zero. In this case we get a contribution of $O(|1-z|^{-k\alpha+\frac{1}{2}})$ which is $O(|1-z|^{-k\alpha'+c+1})$. Adding all these contributions yields the desired result. ☐

The result when $\alpha = 1/2$ is as follows. Recall that $L(z) := \ln((1-z)^{-1})$.

Proposition 7.3.4. *Let $\alpha = 1/2$. In the notation of Proposition 7.3.3,*

$$\overline{M}_k(z) = (1-z)^{-k+\frac{1}{2}} \sum_{l=0}^{k} C_{k,l} L^{k-l}(z) + O(|1-z|^{-k+1-\epsilon})$$

for $k \geq 1$ and any $\epsilon > 0$, where the $C_{k,l}$'s are constants. The constant multiplying the lead-order term is given by

$$C_{k,0} = \frac{(2k-2)!}{2^{2k-2}(k-1)!\pi^{k/2}}. \tag{7.3.7}$$

Proof. For $k = 1$ the claim is true by (7.2.7). For $k \geq 2$ we estimate each term in (7.3.5).

For this paragraph, suppose that k_1 and k_2 are both nonzero. Then using the induction hypothesis we get

$$\begin{aligned}\frac{z}{4}\overline{M}_{k_1}(z)\overline{M}_{k_2}(z) = \frac{1}{4}(1-z)^{-(k_1+k_2)+1}\sum_{l=0}^{k_1}\sum_{m=0}^{k_2} C_{k_1,l}C_{k_2,m}L^{(k_1+k_2)-(l+m)}(z)& \\ + O(|1-z|^{-(k_1+k_2)+\frac{3}{2}-2\epsilon}).&\end{aligned} \tag{7.3.8}$$

If $k_3 = 0$ the corresponding contribution to $\overline{R}_k(z)$ is

$$\frac{1}{4}\binom{k}{k_1}(1-z)^{-k+1}\sum_{l=0}^{k} A_{k,l}L^{k-l}(z) + O(|1-z|^{-k+\frac{3}{2}-2\epsilon}),$$

where $A_{k,0} = C_{k_1,0}C_{k-k_1,0}$ and $(A_{k,l})_{l=1}^{k}$ can be determined easily from (7.3.8). If $k_3 \neq 0$ then we use Lemma 7.1.2 once again to get

$$\frac{z}{4}\overline{M}_{k_1}(z)\overline{M}_{k_2}(z) = \sum_{l=0}^{k_1+k_2} \widetilde{A}_{k_1+k_2,l}\,\mathrm{Li}_{-(k_1+k_2)+2,k_1+k_2-l}(z) + O(|1-z|^{-(k_1+k_2)+\frac{3}{2}-2\epsilon}),$$

where $(\widetilde{A}_{k_1+k_2,l})_{l=0}^{k_1+k_2}$ can be determined using (7.3.8) and Lemma 7.1.2. The corresponding contribution to $\overline{R}_k(z)$ is then $\binom{k}{k_1,k_2,k_3}$ times:

$$\sum_{l=0}^{k_1+k_2} \widetilde{A}_{k_1+k_2,l}\,\mathrm{Li}_{-k+\frac{k_3}{2}+2,k_1+k_2-l}(z) + \mathrm{Li}_{-\frac{k_3}{2},0}(z) \odot O(|1-z|^{-(k_1+k_2)+\frac{3}{2}-2\epsilon}).$$

Invoking Lemma 7.1.1 we see that the contribution when $k_3 \neq 0$ is $O(|1-z|^{-k+\frac{k_3}{2}+1-\epsilon})$, which is $O(|1-z|^{-k+\frac{3}{2}-\epsilon})$.

Next suppose that k_1 is nonzero but $k_2 = 0$. In this case using the induction hypothesis and we see that

$$\frac{z}{4}\overline{M}_{k_1}(z)\overline{M}_{k_2}(z) = \frac{1}{2}(1-z)^{-k_1+\frac{1}{2}}\sum_{l=0}^{k_1} C_{k_1,l}L^{k_1-l}(z) + O(|1-z|^{-k_1+1-\epsilon}).$$

By Lemma 7.1.2 this is

$$\sum_{l=0}^{k_1} B_{k_1,l}\,\mathrm{Li}_{-k_1+\frac{3}{2},k_1-l}(z) + O(|1-z|^{-k_1+1-\epsilon}),$$

where $(B_{k_1,l})_{l=0}^{k_1}$ are constants. The corresponding contribution to $\overline{R}_k(z)$ is then $\binom{k}{k_1}$ times:

$$\sum_{l=0}^{k_1} B_{k_1,l}\,\mathrm{Li}_{-k+\frac{k_3}{2}+\frac{3}{2},k_1-l}(z) + \mathrm{Li}_{-\frac{k_3}{2},0}(z) \odot O(|1-z|^{-k_1+1-\epsilon}).$$

When $k_3 \geq 2$ the contribution is $O(|1-z|^{-k+\frac{k_3}{2}+\frac{1}{2}-\epsilon})$, which is $O(|1-z|^{-k+\frac{3}{2}-\epsilon})$. When $k_3 = 1$ the contribution to $\overline{R}_k(z)$ is

$$(1-z)^{-k+1} \sum_{l=0}^{k-1} \widetilde{B}_{k-1,l} L^{k-1-l}(z) + O(|1-z|^{-k+\frac{3}{2}-\epsilon}),$$

for some suitably defined constants $\widetilde{B}_{k,l}$. Note that there is no lead-order contribution from this term. The terms where $k_1 = 0$ and k_2 is nonzero are similarly handled.

The final contribution from the single term with $k_1 = k_2 = 0$ is $O(|1-z|^{-\frac{k}{2}+\frac{1}{2}})$, which is $O(|1-z|^{-k+\frac{3}{2}})$ for $k \geq 2$.

Summing all the contributions we get

$$\overline{R}_k(z) = (1-z)^{-k+1} \sum_{l=0}^{k} C_{k,l} L^{k-l}(z) + O(|1-z|^{-k+\frac{3}{2}-\epsilon}),$$

where, for $k \geq 2$,

$$C_{k,0} = \frac{1}{4} \sum_{k_1=1}^{k-1} \binom{k}{k_1} C_{k_1,0} C_{k-k_1,0}, \qquad C_{1,0} = \frac{1}{\sqrt{\pi}},$$

$$C_{k,l} = \frac{1}{4} \sum_{k_1=1}^{k-1} \binom{k}{k_1} A_{k,l} + 2\widetilde{B}_{k-1,l}, \qquad l \geq 1.$$

The recurrence for $C_{k,0}$ can be easily solved to yield

$$C_{k,0} = 2 \cdot k!\, [z^k] \left[1 - \left(1 - \frac{z}{\sqrt{\pi}}\right)^{1/2} \right] = \frac{(2k-2)!}{2^{2k-2}(k-1)!\pi^{k/2}}.$$

The result follows immediately. □

7.4 Asymptotics of moments

For $0 < \alpha < 1/2$, we have seen in Propositions 7.3.1 and 7.3.2 that the generating function $\widehat{M}_k(z)$ of $\hat{\mu}_n(k) = \beta_n \tilde{\mu}_n(k)/4^n$ satisfies

$$\widehat{M}_k(z) = C_k (1-z)^{-k(\alpha+\frac{1}{2})+\frac{1}{2}} + O(|1-z|^{-k(\alpha+\frac{1}{2})+\frac{1}{2}+c}),$$

where $c := \min\{2\alpha - \epsilon, 1/2\}$. By Theorem 4.1.2,

$$\frac{\beta_n \tilde{\mu}_n(k)}{4^n} = C_k \frac{n^{k(\alpha+\frac{1}{2})-\frac{3}{2}}}{\Gamma(k(\alpha+\frac{1}{2})-\frac{1}{2})} + O(n^{k(\alpha+\frac{1}{2})-\frac{3}{2}-c}).$$

Recall that

$$\beta_n = \frac{4^n}{\sqrt{\pi} n^{3/2}} \left(1 + O(\tfrac{1}{n})\right),$$

so that

$$\tilde{\mu}_n(k) = \frac{C_k \sqrt{\pi}}{\Gamma(k(\alpha+\frac{1}{2})-\frac{1}{2})} n^{k(\alpha+\frac{1}{2})} + O(n^{k(\alpha+\frac{1}{2})-c}). \tag{7.4.1}$$

For $\alpha > 1/2$ a similar analysis using Proposition 7.3.3 yields

$$\mu_n(k) = \frac{C_k\sqrt{\pi}}{\Gamma(k(\alpha+\frac{1}{2})-\frac{1}{2})} n^{k(\alpha+\frac{1}{2})} + O(n^{k(\alpha+\frac{1}{2})-c}), \tag{7.4.2}$$

with now c as defined at (7.3.6). Finally, when $\alpha = 1/2$ the asymptotics of the moments are given by

$$\mu_n(k) = \left(\frac{1}{\sqrt{\pi}}\right)^k (n \ln n)^k + O(n^k(\ln n)^{k-1}). \tag{7.4.3}$$

7.5 The limiting distributions

In Section 7.5.1 we will use our moment estimates (7.4.1) and (7.4.2) with the method of moments to derive limiting distributions for our additive functionals. The case $\alpha = 1/2$ requires a somewhat delicate analysis, which we will present separately in Section 7.5.2.

7.5.1 $\alpha \neq 1/2$

We first handle the case $0 < \alpha < 1/2$. (We assume this restriction until just before Proposition 7.5.2.) We have

$$\tilde{\mu}_n(1) = \mathbf{E}\,\widetilde{X}_n = \mathbf{E}\,[X_n - C_0(n+1)] = \frac{C_1\sqrt{\pi}}{\Gamma(\alpha)} n^{\alpha+\frac{1}{2}} + O(n^{\alpha+\frac{1}{2}-c}) \tag{7.5.1}$$

with $c := \min\{2\alpha - \epsilon, 1/2\}$ and

$$\tilde{\mu}_n(2) = \mathbf{E}\,\widetilde{X}_n^2 = \frac{C_2\sqrt{\pi}}{\Gamma(2\alpha+\frac{1}{2})} n^{2\alpha+1} + O(n^{2\alpha+1-c}).$$

So

$$\mathbf{Var}\,X_n = \mathbf{Var}\,\widetilde{X}_n = \tilde{\mu}_n(2) - [\tilde{\mu}_n(1)]^2 = \sigma^2 n^{2\alpha+1} + O(n^{2\alpha+1-c}), \tag{7.5.2}$$

where

$$\sigma^2 := \frac{C_2\sqrt{\pi}}{\Gamma(2\alpha+\frac{1}{2})} - \frac{C_1^2\pi}{\Gamma^2(\alpha)}. \tag{7.5.3}$$

We also have, for $k \geq 1$,

$$\mathbf{E}\left[\frac{\widetilde{X}_n}{n^{\alpha+\frac{1}{2}}}\right]^k = \frac{\tilde{\mu}_n(k)}{n^{k(\alpha+\frac{1}{2})}} = \frac{C_k\sqrt{\pi}}{\Gamma(k(\alpha+\frac{1}{2})-\frac{1}{2})} + O(n^{-c}). \tag{7.5.4}$$

The following lemma provides a sufficient bound on the moments facilitating the use of the method of moments.

Lemma 7.5.1. *Define $\alpha' := \alpha + \frac{1}{2}$. There exists a constant $A < \infty$ depending only on α such that*

$$\left|\frac{C_k}{k!}\right| \leq A^k k^{\alpha' k}$$

for all $k \geq 1$.

Proof. The proof is by induction. For the basis case $k = 1$, we simply choose A large enough so that the claim is true. For the induction step, assume $k \geq 2$. Define $s_k := C_k/k!$. Dividing (7.3.3) by $k!$ we get

$$s_k = \frac{1}{4}\sum_{j=1}^{k-1} s_j s_{k-j} + s_{k-1}\frac{\Gamma(k\alpha' - 1)}{\Gamma(k\alpha' - 1 - \alpha)}. \tag{7.5.5}$$

Using Stirling's approximation we can find a constant $\gamma < \infty$ depending only on α so that, for all $k \geq 2$,

$$\left|\frac{\Gamma(k\alpha' - 1)}{\Gamma(k\alpha' - 1 - \alpha)}\right| \leq \gamma k^{\alpha}.$$

Using this in (7.5.5) we get

$$|s_k| \leq \frac{1}{4}\sum_{j=1}^{k-1} |s_j||s_{k-j}| + \gamma|s_{k-1}|k^{\alpha}$$

By the induction hypothesis

$$|s_k| \leq \frac{A^k}{4}\sum_{j=1}^{k-1}[j^j(k-j)^{k-j}]^{\alpha'} + \gamma A^{k-1}(k-1)^{\alpha'(k-1)}k^{\alpha}.$$

Now $x^x(k-x)^{k-x}$ decreases as x increases for $0 < x \leq k/2$ so that we can bound the sum above by the sum of two times the $j = 1$ term and k times the $j = 2$ term. This and trivial bounds yield

$$\begin{aligned}
|s_k| &\leq \frac{A^k}{2}\left[(k-1)^{k-1} + \frac{k}{2} \times 4(k-2)^{k-2}\right]^{\alpha'} + \gamma A^{k-1}k^{\alpha' k - \frac{1}{2}} \\
&\leq \frac{A^k}{2}[k^{k-1} + 2k^{k-1}]^{\alpha'} + A^k\frac{\gamma}{A}k^{\alpha' k - \frac{1}{2}} \\
&= \frac{A^k}{2}(3k^{k-1})^{\alpha'} + A^k\frac{\gamma}{A}k^{\alpha' k - \frac{1}{2}} \\
&= \left[\frac{3^{\alpha'}}{2}k^{-\alpha'} + \frac{\gamma}{A}k^{-\frac{1}{2}}\right]A^k k^{\alpha' k} \leq A^k k^{\alpha' k},
\end{aligned}$$

the last inequality justified for all $k \geq 2$ by choosing A large enough. This proves the claim. ☐

It follows from Lemma 7.5.1 and Stirling's approximation that

$$\left|\frac{C_k\sqrt{\pi}}{k!\Gamma(k(\alpha + \frac{1}{2}) - \frac{1}{2})}\right| \leq B^k \tag{7.5.6}$$

for large enough B depending only on α. Using standard arguments [3, Theorem 30.1] it follows that X_n suitably normalized has a limiting distribution that is characterized by its moments. Before we state the result, we observe that the argument presented above can be adapted with minor modifications to treat the case $\alpha > 1/2$, with $\widetilde{X}_n$ replaced by X_n. We can now state a result for $\alpha \neq 1/2$. We will use the notation $\xrightarrow{\mathcal{L}}$ to denote convergence in law (or distribution).

Proposition 7.5.2. *Let X_n denote the additive functional on Catalan trees induced by the toll* *sequence* $(n^{\alpha})_{n\geq 0}$*. Define the random variable* Y_n *as follows:*

$$Y_n := \begin{cases} \dfrac{X_n - C_0(n+1)}{n^{\alpha+\frac{1}{2}}} & 0<\alpha<1/2, \\ \dfrac{X_n}{n^{\alpha+\frac{1}{2}}} & \alpha > 1/2, \end{cases}$$

where

$$C_0 := \sum_{n=0}^{\infty} n^{\alpha}\frac{\beta_n}{4^n}, \qquad \beta_n = \frac{1}{n+1}\binom{2n}{n}.$$

Then

$$Y_n \xrightarrow{\mathcal{L}} Y;$$

here Y *is a random variable with the unique distribution whose moments are*

$$\mathbf{E}\, Y^k = \frac{C_k\sqrt{\pi}}{\Gamma(k(\alpha+\frac{1}{2})-\frac{1}{2})}, \tag{7.5.7}$$

where the C_k*'s satisfy the recurrence*

$$C_k = \frac{1}{4}\sum_{j=1}^{k-1}\binom{k}{j}C_jC_{k-j} + k\frac{\Gamma(k\alpha+\frac{k}{2}-1)}{\Gamma((k-1)\alpha+\frac{k}{2}-1)}C_{k-1}, k\geq 2; \quad C_1 = \frac{\Gamma(\alpha-\frac{1}{2})}{\sqrt{\pi}}.$$

The case $\alpha = 1/2$ is handled in Section 7.5.2, leading to Proposition 7.5.5, and a unified result for all cases is stated as Theorem 7.5.7.

Remark 7.5.3. We now consider some properties of the limiting random variable $Y \equiv Y(\alpha)$ defined by its moments at (7.5.7) for $\alpha \neq 1/2$.

(a) When $\alpha = 1$ setting $\Omega_k := C_k/2$ we see immediately that

$$\mathbf{E}\, Y^k = \frac{-\Gamma(-1/2)}{\Gamma((3k-1)/2)}\Omega_k,$$

where

$$2\Omega_k = \sum_{j=1}^{k-1}\binom{k}{j}\Omega_j\Omega_{k-j} + k(3k-4)\Omega_{k-1}, \qquad \Omega_1 = \frac{1}{2}.$$

Thus Y has the ubiquitous Airy distribution and we have recovered the limiting distribution of path length in Catalan trees [70–72]. The Airy distribution arises in many contexts including parking allocations, hashing tables, trees, discrete random walks, mergesorting, etc.—see, e.g., the introduction of [30] which contains numerous references to the Airy distribution.

(b) When $\alpha = 2$, setting $\eta := Y/\sqrt{2}$ and $a_{0,l} := 2^{2l-1}C_l$, we see that

$$\mathbf{E}\, \eta^l = \frac{\sqrt{\pi}}{2^{(5l-2)/2}\Gamma((5l-1)/2)}a_{0,l},$$

where

$$a_{0,l} = \frac{1}{2}\sum_{j=1}^{l-1}\binom{l}{j}a_{0,j}a_{0,l-j} + l(5l-4)(5l-6), \qquad a_{0,1} = 1.$$

We have thus recovered the recurrence for the moments of the distribution $\mathcal{L}(\eta)$, which arises in the study of the Wiener index of Catalan trees [41, proof of Theorem 3.3 in Section 5].

(c) Consider the variance σ^2 defined at (7.5.3).

   (i) Figure 7.5.1, plotted using `Mathematica`, suggests that σ^2 is positive for all $\alpha > 0$. We will prove this fact in Theorem 7.5.7. There is also numerical evidence that σ^2 is unimodal with $\max_\alpha \sigma^2(\alpha) \doteq 0.198946$ achieved at $\alpha \doteq 0.682607$. (Here $\doteq$ denotes approximate equality.)



Figure 7.5.1: σ^2 of (7.5.3) as a function of α.

   (ii) As $\alpha \to \infty$, using Stirling's approximation one can show that $\sigma^2 \sim (\sqrt{2}-1)\alpha^{-1}$.

   (iii) As $\alpha \downarrow 0$, using a Laurent series expansion of $\Gamma(\alpha)$ we see that $\sigma^2 \sim 4(1 - \ln 2)\alpha$.

   (iv) Though the random variable $Y(\alpha)$ has been defined only for $\alpha \neq 1/2$, the variance σ^2 has a limit at $\alpha = 1/2$:

$$\lim_{\alpha \to 1/2} \sigma^2(\alpha) = \frac{8 \ln 2}{\pi} - \frac{\pi}{2}. \tag{7.5.8}$$

(d) Figure 7.5.2 shows the third central moment $\mathbf{E}\,[Y - \mathbf{E}\,Y]^3$ as a function of α. The plot suggests that the third central moment is positive for each $\alpha > 0$, which would also establish that $Y(\alpha)$ is not normal for any $\alpha > 0$. However we do not know a proof of this positive skewness. [Of course, the law of $Y(\alpha)$ is not normal for any $\alpha > 1/2$, since its support is a subset of $[0, \infty)$.]

(e) When $\alpha = 0$, the additive functional with toll sequence $(n^\alpha = 1)_{n \geq 1}$ is n for all trees with n nodes. However, if one considers the random variable $\alpha^{-1/2} Y(\alpha)$ as $\alpha \downarrow 0$, using (7.5.7) and induction one can show that $\alpha^{-1/2} Y(\alpha)$ converges in distribution to the normal distribution with mean 0 and variance $4(1 - \ln 2)$.

(f) Finally, if one considers the random variable $\alpha^{1/2} Y(\alpha)$ as $\alpha \to \infty$, again using (7.5.7) and induction we find that $\alpha^{1/2} Y(\alpha)$ converges in distribution to the unique distribution with kth moment $\sqrt{k!}$.

Remark 7.5.4. Let Y be the unique distribution whose kth moment is $\sqrt{k!}$ for $k = 1, 2, \ldots$. Taking Y^* to be an independent copy of Y and defining $X := YY^*$, we see immediately that X is Exponential with unit mean. It follows by taking logarithms that the distribution of $\ln Y$ is



Figure 7.5.2: $\mathbf{E}\,[Y - \mathbf{E}\,Y]^3$ of Proposition 7.5.2 as a function of α.

a convolution square root of the distribution of $\ln X$. In particular, the characteristic function ϕ of $\ln Y$ has square equal to $\Gamma(1+it)$ at $t \in (-\infty, \infty)$. By exponential decay of $\Gamma(1+it)$ as $t \to \pm\infty$ and standard theory (see, e.g., [20, Chapter XV]), $\ln Y$ has an infinitely smooth density on $(-\infty, \infty)$, and the density and each of its derivatives are bounded.

So Y has an infinitely smooth density on $(0, \infty)$. By change of variables, the density f_Y of Y satisfies

$$f_Y(y) = \frac{f_{\ln Y}(\ln y)}{y}.$$

Clearly $f_Y(y)$ is bounded for y *not* near 0. (We shall drop further consideration of derivatives.) To determine the behavior near 0, we need to know the behavior of $f_{\ln Y}(\ln y)/y$ as $y \to 0$. Using the Fourier inversion formula, we may equivalently study

$$e^x f_{\ln Y}(-x) = \frac{1}{2\pi} \int_{-\infty}^{\infty} e^{(1+it)x} \phi(t)\, dt,$$

as $x \to \infty$. By an application of the method of steepest descents [(7.2.11) in [5], with $g_0 = 1$, $\beta = 1/2$, w the identity map, $z_0 = 0$, and $\alpha = 0$], we get

$$f_Y(y) \sim \frac{1}{\sqrt{\pi \ln(1/y)}} \quad \text{as } y \downarrow 0.$$

Hence f_Y is bounded everywhere.

Using the Cauchy integral formula and simple estimates, it is easy to show that

$$f_Y(y) = o(e^{-My}) \quad \text{as } y \to \infty$$

for any $M < \infty$. Computations using the WKB method [34] suggest

$$f_Y(y) \sim (2/\pi)^{1/4} y^{1/2} \exp(-y^2/2) \quad \text{as } y \to \infty, \tag{7.5.9}$$

in agreement with numerical calculations using `Mathematica`. [In fact, the right-side of (7.5.9) appears to be a highly accurate approximation to $f_Y(y)$ for all $y \geq 1$.] Figure 7.5.3 depicts the salient features of f_Y. In particular, note the steep descent of $f_Y(y)$ to 0 as $y \downarrow 0$ and the quasi-Gaussian tail.



Figure 7.5.3: f_Y of Remark 7.5.4.

7.5.2 $\alpha = 1/2$

For $\alpha = 1/2$, from (7.4.3) we see immediately that

$$\mathbf{E}\left[\frac{X_n}{n \ln n}\right]^k = \left(\frac{1}{\sqrt{\pi}}\right)^k + O\left(\frac{1}{\ln n}\right).$$

Thus the random variable $X_n/(n \ln n)$ converges in distribution to the degenerate random variable $1/\sqrt{\pi}$. To get a nondegenerate distribution, we carry out an analysis similar to the one that led to (7.2.7), getting more precise asymptotics for the mean of X_n. The refinement of (7.2.7) that we need is the following, whose proof we omit:

$$A(z) \odot \mathrm{CAT}(z/4) = \frac{1}{\sqrt{\pi}}(1-z)^{-1/2}L(z) + D_0(1-z)^{-1/2} + O(|1-z|^{\frac{1}{2}-\epsilon}),$$

where D_0 is a constant whose value we will not need in our subsequent analysis. By Theorem 4.1.2, this leads to

$$\mathbf{E}\,X_n = \frac{1}{\sqrt{\pi}} n \ln n + D_1 n + O(n^{\epsilon}), \tag{7.5.10}$$

where D_1 is another constant whose value we will not need in our subsequent analysis. Now analyzing the random variable $X_n - \pi^{-1/2} n \ln n$ in a manner similar to that of Section 7.3 we obtain

$$\mathbf{Var}\,[X_n - \pi^{-1/2} n \ln n] = \left(\frac{8}{\pi}\ln 2 - \frac{\pi}{2}\right) n^2 + O(n^{\frac{3}{2}+\epsilon}). \tag{7.5.11}$$

Using (7.5.10) and (7.5.11) we conclude that

$$\mathbf{E}\left[\frac{X_n - \pi^{-1/2} n \ln n - D_1 n}{n}\right] = o(1)$$

and

$$\mathbf{Var}\left[\frac{X_n - \pi^{-1/2} n \ln n - D_1 n}{n}\right] = \frac{8}{\pi}\ln 2 - \frac{\pi}{2} = \lim_{\alpha \to 1/2} \sigma^2(\alpha), \tag{7.5.12}$$

where $\sigma^2 \equiv \sigma^2(\alpha)$ is defined at (7.5.3) for $\alpha \neq 1/2$. [Recall (7.5.8) of Remark 7.5.3.]

It is possible to carry out a program similar to that of Section 7.3 to derive asymptotics of higher order moments using singularity analysis. However we choose to sidestep this arduous, albeit mechanical, computation. Instead we will derive the asymptotics of higher moments using a somewhat more direct approach akin to the one employed in [21]. The approach involves approximation of sums by Riemann integrals. To that end, define

$$\widetilde{X}_n := X_n - \pi^{-1/2}(n+1)\ln(n+1) - D_1(n+1), \qquad \text{and} \qquad \hat{\mu}_n(k) := \frac{\beta_n}{4^{n+1}}\mathbf{E}\,\widetilde{X}_n^k. \tag{7.5.13}$$

Note that $\widetilde{X}_0 = -D_1$, $\hat{\mu}_n(0) = \beta_n/4^{n+1}$, and $\hat{\mu}_0(k) = (-D_1)^k/4$. Then, in a now familiar manner, for $n \geq 1$ we find

$$\hat{\mu}_n(k) = 2\sum_{j=1}^{n} \frac{\beta_{j-1}}{4^j}\hat{\mu}_{n-j}(k) + \hat{r}_n(k),$$

where now we define

$$\hat{r}_n(k) := \sum_{\substack{k_1+k_2+k_3=k\\ k_1,k_2<k}} \binom{k}{k_1,k_2,k_3} \sum_{j=1}^{n} \hat{\mu}_{j-1}(k_1)\hat{\mu}_{n-j}(k_2)$$
$$\times \left[\frac{1}{\sqrt{\pi}}(j\ln j + (n+1-j)\ln(n+1-j) - (n+1)\ln(n+1) + \sqrt{\pi}n^{1/2})\right]^{k_3}$$

Passing to generating functions and then back to sequences one gets, for $n \geq 0$,

$$\hat{\mu}_n(k) = \sum_{j=0}^{n}(j+1)\frac{\beta_j}{4^j}\hat{r}_{n-j}(k).$$

Using induction on k, we can approximate $\hat{r}_n(k)$ and $\hat{\mu}_n(k)$ above by integrals and obtain the following result. We omit the proof, leaving it as an exercise for the ambitious reader.

Proposition 7.5.5. *Let X_n be the additive functional induced by the toll sequence $(n^{1/2})_{n\geq 1}$ on Catalan trees. Define $\widetilde{X}_n$ as in (7.5.13). Then*

$$\mathbf{E}\,[\widetilde{X}_n/n]^k = m_k + o(1) \text{ as } n \to \infty,$$

where $m_0 = 1$, $m_1 = 0$, and, for $k \geq 2$,

$$m_k = \frac{1}{4\sqrt{\pi}}\frac{\Gamma(k-1)}{\Gamma(k-\frac{1}{2})}\left[\sum_{\substack{k_1+k_2+k_3=k\\ k_1,k_2<k}} \binom{k}{k_1,k_2,k_3} m_{k_1}m_{k_2}\left(\frac{1}{\sqrt{\pi}}\right)^{k_3} J_{k_1,k_2,k_3} + 4\sqrt{\pi}km_{k-1}\right], \tag{7.5.14}$$

where

$$J_{k_1,k_2,k_3} := \int_{x=0}^{1} x^{k_1-\frac{3}{2}}(1-x)^{k_2-\frac{3}{2}}[x\ln x + (1-x)\ln(1-x)]^{k_3}\,dx.$$

Furthermore $\widetilde{X}_n/(n+1) \xrightarrow{\mathcal{L}} Y$, where Y is a random variable with the unique distribution whose moments are $\mathbf{E}\,Y^k = m_k$, $k \geq 0$.

7.5.3 A unified result

The approach outlined in the preceding section can also be used for the case $\alpha \neq 1/2$. For completeness, we state the result for that case here (without proof).

Proposition 7.5.6. *Let X_n be the additive functional induced by the toll sequence $(n^{\alpha})_{n\geq 1}$ on*

Catalan trees. Let $\alpha' := \alpha + \frac{1}{2}$. *Define* $\widetilde{X}_n$ *as*

$$\widetilde{X}_n := \begin{cases} X_n - C_0(n+1) - \dfrac{\Gamma(\alpha - \frac{1}{2})}{\Gamma(\alpha)}(n+1)^{\alpha'} & 0 < \alpha < 1/2, \\ X_n - \dfrac{\Gamma(\alpha - \frac{1}{2})}{\Gamma(\alpha)}(n+1)^{\alpha'} & \alpha > 1/2, \end{cases} \tag{7.5.15}$$

where

$$C_0 := \sum_{n=1}^{\infty} n^{\alpha} \frac{\beta_n}{4^n}.$$

Then, for $k = 0, 1, 2, \ldots$,

$$\mathbf{E}\left[\widetilde{X}_n / n^{\alpha'}\right]^k = m_k + o(1) \text{ as } n \to \infty,$$

where $m_0 = 1$, $m_1 = 0$, *and, for* $k \geq 2$,

$$m_k = \frac{1}{4\sqrt{\pi}} \frac{\Gamma(k\alpha' - 1)}{\Gamma(k\alpha' - \frac{1}{2})} \times \left[\sum_{\substack{k_1+k_2+k_3=k \\ k_1,k_2<k}} \binom{k}{k_1, k_2, k_3} m_{k_1} m_{k_2} \left(\frac{\Gamma(\alpha - \frac{1}{2})}{\Gamma(\alpha)} \right)^{k_3} J_{k_1,k_2,k_3} + 4\sqrt{\pi} k m_{k-1} \right], \tag{7.5.16}$$

with

$$J_{k_1,k_2,k_3} := \int_0^1 x^{k_1\alpha' - \frac{3}{2}} (1-x)^{k_2\alpha' - \frac{3}{2}} [x^{\alpha'} + (1-x)^{\alpha'} - 1]^{k_3} \, dx.$$

Furthermore, $\widetilde{X}_n \xrightarrow{\mathcal{L}} Y$, *where* Y *is a random variable with the unique distribution whose moments are* $\mathbf{E}\, Y^k = m_k$.

[The reader may wonder as to why we have chosen to state Proposition 7.5.6 using several instances of $n+1$, rather than n, in (7.5.15). The reason is that use of $n+1$ is somewhat more natural in the calculations that establish the proposition.]

In light of Propositions 7.5.2, 7.5.5, and 7.5.6, there are a variety of ways to state a unified result. We state one such version here.

Theorem 7.5.7. *Let* X_n *denote the additive functional induced by the toll sequence* $(n^{\alpha})_{n \geq 1}$ *on Catalan trees. Then*

$$\frac{X_n - \mathbf{E}\, X_n}{\sqrt{\mathbf{Var}\, X_n}} \xrightarrow{\mathcal{L}} W,$$

where the distribution of W *is described as follows:*

(a) For $\alpha \neq 1/2$,

$$W = \frac{1}{\sigma}\left(Y - \frac{C_1\sqrt{\pi}}{\Gamma(\alpha)} \right), \qquad \text{with} \qquad \sigma^2 := \frac{C_2\sqrt{\pi}}{\Gamma(2\alpha + \frac{1}{2})} - \frac{C_1^2 \pi}{\Gamma^2(\alpha)} > 0,$$

where Y *is a random variable with the unique distribution whose moments are*

$$\mathbf{E}\, Y^k = \frac{C_k\sqrt{\pi}}{\Gamma(k(\alpha + \frac{1}{2}) - \frac{1}{2})},$$

and the C_k*'s satisfy the recurrence (7.3.3).*

(b) For $\alpha = 1/2$,

$$W = \frac{Y}{\sigma}, \qquad \textit{with} \quad \sigma^2 := \frac{8}{\pi}\ln 2 - \frac{\pi}{2},$$

where Y *is a random variable with the unique distribution whose moments* $m_k = \mathbf{E}\, Y^k$ *are given by (7.5.14).*

Proof. Define

$$W_n := \frac{X_n - \mathbf{E}\, X_n}{\sqrt{\mathbf{Var}\, X_n}}$$

(a) Consider first the case $\alpha < 1/2$ and let $\alpha' := \alpha + \frac{1}{2}$. By (7.5.1),

$$\mathbf{E}\, X_n = C_0(n+1) + \frac{C_1\sqrt{\pi}}{\Gamma(\alpha)} n^{\alpha'} + o(n^{\alpha'}). \tag{7.5.17}$$

Since $\widetilde{X}_n$ defined at (7.5.15) and X_n differ by a deterministic amount, $\mathbf{Var}\, X_n = \mathbf{Var}\, \widetilde{X}_n$. Now by Proposition 7.5.6,

$$\mathbf{Var}\, \widetilde{X}_n = \mathbf{E}\, \widetilde{X}_n^2 - (\mathbf{E}\, \widetilde{X}_n)^2 = (m_2 + o(1))n^{2\alpha'} - (m_1^2 + o(1))n^{2\alpha'} = (m_2 + o(1))n^{2\alpha'}. \tag{7.5.18}$$

So σ^2 equals m_2 defined at (7.5.16), namely,

$$\frac{1}{4\sqrt{\pi}} \frac{\Gamma(2\alpha' - 1)}{\Gamma(2\alpha' - \frac{1}{2})} \left(\frac{\Gamma(\alpha - \frac{1}{2})}{\Gamma(\alpha)} \right)^2 J_{0,0,2}.$$

Thus to show $\sigma^2 > 0$ it is enough to show that $J_{0,0,2} > 0$. But

$$J_{0,0,2} = \int_x^1 x^{-3/2}(1-x)^{-3/2}[x^{\alpha'} + (1-x)^{\alpha'} - 1]^2\, dx,$$

which is clearly positive. Using (7.5.17) and (7.5.18),

$$W_n = \frac{X_n - C_0(n+1) - \frac{C_1\sqrt{\pi}}{\Gamma(\alpha)} n^{\alpha'} + o(n^{\alpha'})}{(1 + o(1))\sigma n^{\alpha'}},$$

so, by Proposition 7.5.2 and Slutsky's theorem [3, Theorem 25.4], the claim follows.

The case $\alpha > 1/2$ follows similarly.

(b) When $\alpha = 1/2$,

$$\mathbf{E}\, X_n = \frac{1}{\sqrt{\pi}} n \ln n + D_1 n + o(n)$$

by (7.5.10) and

$$\mathbf{Var}\, X_n = \left(\frac{8}{\pi}\ln 2 - \frac{\pi}{2} + o(1) \right) n^2$$

by (7.5.12). The claim then follows easily from Proposition 7.5.5 and Slutsky's theorem. □

Chapter 8

Catalan model: the shape functional

We now turn our attention to the shape functional for Catalan trees. The shape functional is the cost induced by the toll function $t_n \equiv \log n$, $n \geq 1$. For background and results on the shape functional, we refer the reader to [21] and [55].

In the sequel we will improve on the mean and variance estimates obtained in [21] and derive a central limit theorem for the shape functional for Catalan trees. The technique employed is singularity analysis followed by the method of moments.

8.1 Mean

We use the notation and techniques of Section 7.2 again. Observe that now $T(z) = \mathrm{Li}_{0,1}(z)$ and (7.2.5) gives

$$\mathrm{CAT}(z/4) = 2 - \frac{2}{\Gamma(-1/2)}[\mathrm{Li}_{3/2,0}(z) - \zeta(3/2)] + 2\left(1 - \frac{\zeta(1/2)}{\Gamma(-1/2)}\right)(1-z) + O(|1-z|^{3/2}).$$

So

$$T(z) \odot \mathrm{CAT}(z/4) = -\frac{2}{\Gamma(-1/2)}\,\mathrm{Li}_{3/2,1}(z) + \bar{c} + \bar{\bar{c}}(1-z) + O(|1-z|^{\frac{3}{2}-\epsilon}),$$

where $\bar{c}$ and $\bar{\bar{c}}$ denote unspecified (possibly 0) constants. The constant term in the singular expansion of $T(z) \odot \mathrm{CAT}(z/4)$ is already known to be, using Theorem 4.3.5,

$$C_0 = T(z) \odot \mathrm{CAT}(z/4)\Big|_{z=1} = \sum_{n=1}^{\infty} (\log n)\frac{\beta_n}{4^n}.$$

Now using Theorem 4.1.3 to obtain the singular expansion of $\mathrm{Li}_{3/2,1}(z)$, we get

$$T(z) \odot \mathrm{CAT}(z/4) = C_0 - 2(1-z)^{1/2}L(z) - 2(2(1-\log(2)) - \gamma)(1-z)^{1/2} + O(|1-z|),$$

so that

$$A(z) \odot \mathrm{CAT}(z/4) = C_0(1-z)^{-1/2} - 2L(z) - 2(2(1-\log 2) - \gamma) + O(|1-z|^{1/2}). \qquad (8.1.1)$$

Using Theorem 4.1.2 and the asymptotics of the Catalan numbers we get that the mean a_n of the shape functional is given by

$$a_n = C_0(n+1) - 2\sqrt{\pi}n^{1/2} + O(1), \qquad (8.1.2)$$

which agrees with the estimate in Theorem 3.1 of [21] and improves the remainder estimate.

8.2 Second moment and variance

We now derive the asymptotics of the approximately centered second moment and the variance of the shape functional. These estimates will serve as the basis for the induction to follow. We will use the notation of Section 7.3, centering the cost function as before by $C_0(n+1)$.

It is clear from (8.1.1) that

$$\widehat{M_1}(z) = -2L(z) - 2(2(1-\log 2) - \gamma) + O(|1-z|^{1/2}), \tag{8.2.1}$$

and (7.3.2) with $k = 2$ gives us, recalling (7.2.1),

$$\widehat{R}_2(z) = C_0^2 + \mathrm{CAT}(z/4) \odot \mathrm{Li}_{0,2}(z) + 4\,\mathrm{Li}_{0,1}(z) \odot [\frac{z}{4}\,\mathrm{CAT}(z/4)\widehat{M_1}(z)] + \frac{z}{2}\widehat{M}_1^2(z). \tag{8.2.2}$$

We analyze each of the terms in this sum. For the last term, observe that $z/2 \to 1/2$ as $z \to 1$, so that

$$\frac{z}{2}\widehat{M}_1^2(z) = 2L^2(z) + 4(2(1-\log 2) - \gamma)L(z) + 2(2(1-\log 2) - \gamma)^2 + O(|1-z|^{\frac{1}{2}-\epsilon}),$$

the ϵ introduced to avoid logarithmic remainders. The first term is easily seen to be

$$\mathrm{CAT}(z/4) \odot \mathrm{Li}_{0,2}(z) = K + O(|1-z|^{\frac{1}{2}-\epsilon})$$

by Theorem 4.3.5, where

$$K := \sum_{n=1}^{\infty} (\log n)^2 \frac{\beta_n}{4^n}.$$

For the middle term, first observe that

$$\frac{z}{4}\,\mathrm{CAT}(z/4)\widehat{M_1}(z) = -L(z) - (2(1-\log 2) - \gamma) + (1-z)^{1/2}L(z) + O(|1-z|^{1/2})$$

and that $L(z) = \mathrm{Li}_{1,0}(z)$. Thus the third term on the right in (8.2.2) is 4 times:

$$-\mathrm{Li}_{1,1}(z) + \bar{c} + O(|1-z|^{\frac{1}{2}-2\epsilon}) = -\frac{1}{2}L^2(z) + \gamma L(z) + \bar{c} + O(|1-z|^{\frac{1}{2}-\epsilon}).$$

[The singular expansion for $\mathrm{Li}_{1,1}(z)$ was obtained using (4.1.4). We state it here for the reader's convenience:

$$\mathrm{Li}_{1,1}(z) = \frac{1}{2}L^2(z) - \gamma L(z) + \bar{c} + O(|1-z|),$$

where $\bar{c}$ is again an unspecified constant.] Hence

$$\widehat{R}_2(z) = 8(1-\log 2)L(z) + \bar{c} + O(|1-z|^{\frac{1}{2}-\epsilon}),$$

which leads to

$$\widehat{M_2}(z) = 8(1-\log 2)(1-z)^{-1/2}L(z) + \bar{c}(1-z)^{-1/2} + O(|1-z|^{-\epsilon}). \tag{8.2.3}$$

We draw the attention of the reader to the cancellation of the ostensible lead-order term $L^2(z)$. This kind of cancellation will appear again in the next section when we deal with higher moments.

Now using singularity analysis and estimates for the Catalan numbers we get

$$\tilde{\mu}_n(2) = 8(1-\log 2)n\log n + \bar{c}n + O(n^{\frac{1}{2}+\epsilon}). \tag{8.2.4}$$

Using (8.1.2),

$$\mathbf{Var}\, X_n = \tilde{\mu}_n(2) - \tilde{\mu}_n(1)^2 = 8(1-\log 2)n\log n + \bar{c}n + O(n^{\frac{1}{2}+\epsilon}),$$

which agrees with Theorem 3.1 of [21] (after a correction pointed out in [55]) and improves the remainder estimate. In our subsequent analysis we will not need to evaluate the unspecified constant $\bar{c}$.

8.3 Higher moments

We now turn our attention to deriving the asymptotics of higher moments of the shape functional. The main result is as follows.

Proposition 8.3.1. *Define* $\widetilde{X}_n := X_n - C_0(n+1)$*, with* $X_0 := 0$*;* $\tilde{\mu}_n(k) := \mathbf{E}\,\widetilde{X}_n^k$*, with* $\tilde{\mu}_n(0) = 1$ *for all* $n \geq 0$*; and* $\hat{\mu}_n(k) := \beta_n \tilde{\mu}_n(k)/4^n$*. Let* $\widehat{M}_k(z)$ *denote the ordinary generating function of* $\hat{\mu}_n(k)$ *in the argument* n*. For* $k \geq 2$*,* $\widehat{M}_k(z)$ *satisfies*

$$\widehat{M}_k(z) = (1-z)^{-\frac{k-1}{2}} \sum_{j=0}^{\lfloor k/2 \rfloor} C_{k,j} L^{\lfloor k/2 \rfloor - j}(z) + O(|1-z|^{-\frac{k}{2}+1-\epsilon}),$$

with

$$C_{2l,0} = \frac{1}{4} \sum_{j=1}^{l-1} \binom{2l}{2j} C_{2j,0} C_{2l-2j,0}, \qquad C_{2,0} = 8(1-\log 2).$$

Proof. The proof is by induction. For $k = 2$ the claim is true by (8.2.3). We note that the claim is *not* true for $k = 1$. Instead, recalling (8.2.1),

$$\widehat{M}_1(z) = -2L(z) - 2(2(1-\log 2) - \gamma) + O(|1-z|^{1/2}). \tag{8.3.1}$$

For the induction step, let $k \geq 3$. We will first get the asymptotics of $\widehat{R}_k(z)$ defined at (7.3.2) with $T(z) = \mathrm{Li}_{0,1}(z)$. In order to do that we will obtain the asymptotics of each term in the defining sum. We remind the reader that we are only interested in the form of the asymptotic expansion of $\widehat{R}_k(z)$ and the coefficient of the lead order term when k is even. This allows us to "define away" all other constants, their determination delayed to the time when the need arises.

For this paragraph suppose that $k_1 \geq 2$ and $k_2 \geq 2$. Then by the induction hypothesis

$$\frac{z}{4}\widehat{M}_{k_1}(z)\widehat{M}_{k_2}(z) = \frac{1}{4}(1-z)^{-\frac{k_1+k_2}{2}+1} \sum_{l=0}^{\lfloor k_1/2 \rfloor + \lfloor k_2/2 \rfloor} A_{k_1,k_2,l} L^{\lfloor k_1/2 \rfloor + \lfloor k_2/2 \rfloor - l}(z) + O(|1-z|^{-\frac{k_1+k_2}{2}+\frac{3}{2}-\epsilon}), \tag{8.3.2}$$

where $A_{k_1,k_2,0} = C_{k_1,0}C_{k_2,0}$. (a) If $k_3 = 0$ then $k_1 + k_2 = k$ and the corresponding contribution to $\widehat{R}_k(z)$ is given by

$$\frac{1}{4}\binom{k}{k_1}(1-z)^{-\frac{k}{2}+1} \sum_{l=0}^{\lfloor k_1/2 \rfloor + \lfloor (k-k_1)/2 \rfloor} A_{k_1,k-k_1,l} L^{\lfloor k_1/2 \rfloor + \lfloor (k-k_1)/2 \rfloor - l}(z) + O(|1-z|^{-\frac{k}{2}+\frac{3}{2}-\epsilon}). \tag{8.3.3}$$

Observe that if k is even and k_1 is odd the highest power of $L(z)$ in (8.3.3) is $\lfloor k/2 \rfloor - 1$. In all other cases the the highest power of $L(z)$ in (8.3.3) is $\lfloor k/2 \rfloor$. (b) If $k_3 \neq 0$ then we use Lemma 7.1.2 to express (8.3.2) as a linear combination of

$$\left\{ \mathrm{Li}_{-\frac{k_1+k_2}{2}+2,l}(z) \right\}_{l=0}^{\lfloor k_1/2 \rfloor + \lfloor k_2/2 \rfloor}$$

with a remainder that is $O(|1-z|^{-\frac{k_1+k_2}{2}+\frac{3}{2}-\epsilon})$. When we take the Hadamard product of such a term with $\mathrm{Li}_{0,k_3}(z)$ we will get a linear combination of

$$\left\{\mathrm{Li}_{-\frac{k_1+k_2}{2}+2,l+k_3}(z)\right\}_{l=0}^{\lfloor k_1/2\rfloor+\lfloor k_2/2\rfloor}$$

and a smaller remainder. Such terms are all $O(|1-z|^{-\frac{k_1+k_2}{2}+1-\epsilon})$, which is $O(|1-z|^{-\frac{k}{2}+\frac{3}{2}-\epsilon})$.

Next, consider the case when $k_1=1$ and $k_2\geq 2$. Using the induction hypothesis and (8.3.1) we get

$$\frac{z}{4}\widehat{M}_{k_1}(z)\widehat{M}_{k_2}(z) = -\frac{1}{2}(1-z)^{-\frac{k_2-1}{2}}\sum_{j=0}^{\lfloor k_2/2\rfloor+1} B_{k_2,j}L^{\left\lfloor\frac{k_2}{2}\right\rfloor+1-j}(z)+O(|1-z|^{-\frac{k_2}{2}+1-2\epsilon}), \quad (8.3.4)$$

with $B_{k_2,0}=C_{k_2,0}$. (a) If $k_3=0$ then $k_2=k-1$ and the corresponding contribution to $\widehat{R}_k(z)$ is given by

$$-\frac{k}{2}(1-z)^{-\frac{k}{2}+1}\sum_{j=0}^{\lfloor (k-1)/2\rfloor+1} B_{k-1,j}L^{\left\lfloor\frac{k-1}{2}\right\rfloor+1-j}(z)+O(|1-z|^{-\frac{k}{2}+\frac{3}{2}-2\epsilon}). \quad (8.3.5)$$

(b) If $k_3\neq 0$ then Lemma 7.1.2 can be used once again to express (8.3.4) in terms of generalized polylogarithms, whence an argument similar to that at the end of the preceding paragraph yields that the contributions from such terms is $O(|1-z|^{-\frac{k_2-1}{2}-\epsilon})$, which is $O(|1-z|^{-\frac{k}{2}+\frac{3}{2}-\epsilon})$. The case when $k_1\geq 2$ and $k_2=1$ is handled symmetrically.

When $k_1=k_2=1$ then $(z/4)\widehat{M}_{k_1}(z)\widehat{M}_{k_2}(z)$ is $O(|1-z|^{-\epsilon})$ and when one takes the Hadamard product of this term with $\mathrm{Li}_{0,k_3}(z)$ the contribution will be $O(|1-z|^{-2\epsilon})$.

Now consider the case when $k_1=0$ and $k_2\geq 2$. Now $\widehat{M}_0(z)=\mathrm{CAT}(z/4)$, so that

$$\frac{z}{4}\widehat{M}_{k_1}(z)\widehat{M}_{k_2}(z) = \frac{1}{2}(1-z)^{-\frac{k_2-1}{2}}\sum_{j=0}^{\lfloor k_2/2\rfloor} C_{k_2,j}L^{\lfloor k_2/2\rfloor-j}(z)+O(|1-z|^{-\frac{k_2}{2}+1-\epsilon}). \quad (8.3.6)$$

By Lemma 7.1.2 this can be expressed as a linear combination of

$$\left\{\mathrm{Li}_{-\frac{k_2-1}{2}+1,j}(z)\right\}_{j=0}^{\lfloor k_2/2\rfloor}$$

with a $O(|1-z|^{-\frac{k_2}{2}+1-\epsilon})$ remainder. When we take the Hadamard product of such a term with $\mathrm{Li}_{0,k_3}(z)$ we will get a linear combination, call it $S(z)$, of

$$\left\{\mathrm{Li}_{-\frac{k_2-1}{2}+1,j+k_3}(z)\right\}_{j=0}^{\lfloor k_2/2\rfloor}$$

with a remainder of $O(|1-z|^{-\frac{k_2}{2}+1-2\epsilon})$, which is $O(|1-z|^{-\frac{k}{2}+\frac{3}{2}-2\epsilon})$ unless $k_2=k-1$. When $k_2=k-1$, by Lemma 7.1.2 the constant multiplying the lead-order term $\mathrm{Li}_{-\frac{k}{2}+2,\lfloor\frac{k-1}{2}\rfloor}(z)$ in $S(z)$ is $\frac{C_{k-1,0}}{2}\mu_0^{(-\frac{k}{2}+2,\lfloor\frac{k-1}{2}\rfloor)}$. When we take the Hadamard product of this term with $\mathrm{Li}_{0,k_3}(z)$

we get a lead-order term of

$$\frac{C_{k-1,0}}{2}\mu_0^{(-\frac{k}{2}+2,\lfloor\frac{k-1}{2}\rfloor)}\,\mathrm{Li}_{-\frac{k}{2}+2,\lfloor\frac{k-1}{2}\rfloor+1}(z).$$

Now we use Lemma 7.1.1 and the observation that $\lambda_0^{(\alpha,r)}\mu_0^{(\alpha,s)} = 1$ to conclude that the contribution to $\widehat{R}_k(z)$ from the term with $k_1 = 0$ and $k_2 = k-1$ is

$$\frac{k}{2}(1-z)^{-\frac{k}{2}+1}\sum_{j=0}^{\lfloor\frac{k-1}{2}\rfloor+1} D_{k,j}L^{\lfloor\frac{k-1}{2}\rfloor+1-j}(z) + O(|1-z|^{-\frac{k}{2}+\frac{3}{2}-\epsilon}), \tag{8.3.7}$$

with $D_{k,0} = C_{k-1,0}$. Notice that the lead order from this contribution is precisely that from (8.3.5) but with opposite sign; thus the two contributions cancel each other to lead order. The case $k_2 = 0$ and $k_1 \geq 2$ is handled symmetrically.

The last two cases are $k_1 = 0$, $k_2 = 1$ (or vice-versa) and $k_1 = k_2 = 0$. The contribution from these cases can be easily seen to be $O(|1-z|^{-\frac{k}{2}+\frac{3}{2}-2\epsilon})$.

We can now deduce the asymptotic behavior of $\widehat{R}_k(z)$. The three contributions are (8.3.3), (8.3.5), and (8.3.7), with only (8.3.3) contributing in net a term of the form $(1-z)^{-\frac{k}{2}+1}L^{\lfloor k/2\rfloor}(z)$ when k is even. The coefficient of this term when k is even is given by

$$\frac{1}{4}\sum_{\substack{0<k_1<k\\ k_1 \text{ even}}}\binom{k}{k_1}C_{k_1,0}C_{k_2,0}.$$

Finally we can sum up the rest of the contribution, define $C_{k,j}$ appropriately and use (7.3.1) to claim the result. □

8.4 A central limit theorem

Proposition 8.3.1 and singularity analysis (Theorem 4.1.2) allows us to get the asymptotics of the moments of the "approximately centered" shape functional. Using arguments identical to those in Section 7.4 it is clear that for $k \geq 2$

$$\tilde{\mu}_n(k) = \frac{C_{k,0}\sqrt{\pi}}{\Gamma(\frac{k-1}{2})}n^{k/2}[\log n]^{\lfloor k/2\rfloor} + O(n^{k/2}[\log n]^{\lfloor k/2\rfloor-1}).$$

This and the asymptotics of the mean derived in Section 8.1 give us, for $k \geq 1$,

$$E\left[\frac{\tilde{X}_n}{\sqrt{n\log n}}\right]^{2k} \to \frac{C_{2k,0}\sqrt{\pi}}{\Gamma(k-\frac{1}{2})}, \qquad E\left[\frac{\tilde{X}_n}{\sqrt{n\log n}}\right]^{2k-1} = o(1)$$

as $n \to \infty$. The recurrence for $C_{2k,0}$ can be solved easily to yield, for $k \geq 1$,

$$C_{2k,0} = \frac{(2k)!(2k-2)!}{2^k 2^{2k-2}k!(k-1)!}\sigma^{2k},$$

where $\sigma^2 := 8(1-\log 2)$. Then using the identity

$$\frac{\Gamma(k-\frac{1}{2})}{\sqrt{\pi}} = \left(\frac{1}{2}\right)^{\overline{k-1}} = \left[2^{2k-2}\frac{(k-1)!}{(2k-2)!}\right]^{-1}$$

we get

$$\frac{C_{2k,0}\sqrt{\pi}}{\Gamma(k-\frac{1}{2})} = \frac{(2k)!}{2^k k!}\sigma^{2k}.$$

It is clear now that both the "approximately centered" and the normalized shape functional are asymptotically normal.

Theorem 8.4.1. *Let* X_n *denote the shape functional, induced by the toll sequence* $(\log n)_{n\geq 1}$*, for Catalan trees. Then*

$$\frac{X_n - C_0(n+1)}{\sqrt{n\log n}} \xrightarrow{\mathcal{L}} \mathcal{N}(0,\sigma^2) \quad \text{and} \quad \frac{X_n - \mathbf{E}\,X_n}{\sqrt{\mathbf{Var}\,X_n}} \xrightarrow{\mathcal{L}} \mathcal{N}(0,1),$$

where

$$C_0 := \sum_{n=1}^{\infty}(\log n)\frac{\beta_n}{4^n}, \qquad \beta_n = \frac{1}{n+1}\binom{2n}{n},$$

and $\sigma^2 := 8(1-\log 2)$.

8.5 Sufficient conditions for asymptotic normality

In this speculative final section we briefly examine the behavior of a general additive functional X_n induced by a given "small" toll sequence (t_n). We have seen evidence [Remark 7.5.3(d)] that if (t_n) is the "large" toll sequence n^α for any fixed $\alpha > 0$, then the limiting behavior is non-normal. When $t_n = \log n$ (or $t_n = n^\alpha$ and $\alpha \downarrow 0$), the (limiting) random variable is normal. Where is the interface between normal and non-normal asymptotics? We have carried out arguments similar to those leading to Propositions 7.5.5 and 7.5.6 (see also [21]) that suggest a sufficient condition for asymptotic normality, but our "proof" is somewhat heuristic, and further technical conditions on (t_n) may be required. Nevertheless, to inspire further work, we present our preliminary indications.

We assume that $t_n \equiv t(n)$, where $t(\cdot)$ is a function of a nonnegative real argument. Suppose that $x^{-3/2}t(x)$ is (ultimately) nonincreasing and that $xt'(x)$ is slowly varying at infinity. Then

$$\mathbf{E}\,X_n = C_0(n+1) - (1+o(1))2\sqrt{\pi}n^{3/2}t'(n),$$

where

$$C_0 = \sum_{n=1}^{\infty} t_n \frac{\beta_n}{4^n}.$$

Furthermore,

$$\mathbf{Var}\,X_n \sim 8(1-\log 2)[nt'(n)]^2 n\log n,$$

and

$$\frac{X_n - C_0(n+1)}{nt'(n)\sqrt{n\log n}} \xrightarrow{\mathcal{L}} \mathcal{N}(0,\sigma^2), \text{ where } \sigma^2 = 8(1-\log 2).$$

This asymptotic normality can also be stated in the form

$$\frac{X_n - \mathbf{E}\,X_n}{\sqrt{\mathbf{Var}\,X_n}} \xrightarrow{\mathcal{L}} \mathcal{N}(0,1).$$

Part IV

Appendices and Bibliography

Appendix A

Solution of an Euler differential equation

We now provide the proof of Theorem 2.3.2, which states the general solution of the differential equation (2.3.4) with initial conditions $a_j = b_j$, $0 \le j \le m-2$. This linear differential equation can be written in the form

$$\mathbf{L}g = h \tag{A.1}$$

where the operator $\mathbf{L}$ is defined as

$$(\mathbf{L}g)(z) := g^{(m-1)}(z) - m!(1-z)^{-(m-1)}g(z). \tag{A.2}$$

We seek the solution $g = A$ corresponding to input $h = B^{(m-1)}$.

Equations of the form (A.1)–(A.2) are members of a class known as *Euler differential equations*. In this appendix we discuss a general method for solving Euler equations, restricting attention, for the sake of definiteness and practicality, to (A.1)–(A.2). We assume that the reader is familiar with the theory of linear differential equations with constant coefficients (see, e.g., [6]).

The homogeneous solution

The technique for solving $\mathbf{L}g = 0$ is quite easily summarized: make the change of variable $z = 1 - e^{-x}$, that is, $x = \ln\left((1-z)^{-1}\right)$. For notational convenience we will abbreviate $\ln\left((1-z)^{-1}\right)$ as $L(z)$. Then consider the function $\tilde{g}$ defined by

$$\tilde{g}(x) := g(1 - e^{-x}), \qquad \text{i.e.,} \qquad g(z) = \tilde{g}(L(z)). \tag{A.3}$$

Lemma A.1. *The derivatives of g are related to those of $\tilde{g}$ by*

$$g^{(k)}(z) = (1-z)^{-k} \sum_{j=0}^{k} \begin{bmatrix} k \\ j \end{bmatrix} \tilde{g}^{(j)}(L(z)), \tag{A.4}$$

where $\left[{k \atop j}\right]$ denotes a signless Stirling number of the first kind.

Proof. Since $\left[{0 \atop 0}\right] = 1$, the basis of the induction is (A.3). For the induction step we differentiate the induction hypothesis (A.4) to get

$$\begin{aligned} g^{(k+1)}(z) &= (1-z)^{-(k+1)} \left[k \sum_{j=0}^{k} \begin{bmatrix} k \\ j \end{bmatrix} \tilde{g}^{(j)}(L(z)) + \sum_{j=0}^{k} \begin{bmatrix} k \\ j \end{bmatrix} \tilde{g}^{(j+1)}(L(z)) \right] \\ &= (1-z)^{-(k+1)} \left\{ k \begin{bmatrix} k \\ 0 \end{bmatrix} \tilde{g}^{(0)}(L(z)) \right. \\ &\qquad \left. + \sum_{j=1}^{k} \left(k \begin{bmatrix} k \\ j \end{bmatrix} + \begin{bmatrix} k \\ j-1 \end{bmatrix} \right) \tilde{g}^{(j)}(L(z)) + \begin{bmatrix} k \\ k \end{bmatrix} \tilde{g}^{(k+1)}(L(z)) \right\}. \end{aligned}$$

We now use standard identities for the Stirling numbers [43, §1.2.6]:

$$k\begin{bmatrix}k\\0\end{bmatrix} = 0 = \begin{bmatrix}k+1\\0\end{bmatrix},\ k \ge 0, \qquad \begin{bmatrix}k\\k\end{bmatrix} = 1 = \begin{bmatrix}k+1\\k+1\end{bmatrix},\ k \ge 0,$$

and

$$k\begin{bmatrix}k\\j\end{bmatrix} + \begin{bmatrix}k\\j-1\end{bmatrix} = \begin{bmatrix}k+1\\j\end{bmatrix},\ 1 \le j \le k.$$

Thus

$$g^{(k+1)}(z) = (1-z)^{-(k+1)} \sum_{j=0}^{k+1} \begin{bmatrix}k+1\\j\end{bmatrix} \tilde{g}^{(j)}(L(z)),$$

completing the induction. □

The left-hand side of (A.1) can hence be expressed as

$$(\mathbf{L}g)(z) = (1-z)^{-(m-1)} \left\{ \sum_{j=0}^{m-1} \begin{bmatrix}m-1\\j\end{bmatrix} \tilde{g}^{(j)}(L(z)) - m!\tilde{g}(L(z)) \right\}$$

so that solving $\mathbf{L}g = 0$ is equivalent to solving $\widetilde{\mathbf{L}}\tilde{g} = 0$, where

$$\widetilde{\mathbf{L}}\tilde{g}(x) := \sum_{j=0}^{m-1} \begin{bmatrix}m-1\\j\end{bmatrix} \tilde{g}^{(j)}(x) - m!\tilde{g}(x). \tag{A.5}$$

But this is a linear differential equation with constant coefficients. Its *indicial polynomial*, or *characteristic polynomial*, is

$$\psi_m(\lambda) \equiv \psi(\lambda) := \sum_{j=0}^{m-1} \begin{bmatrix}m-1\\j\end{bmatrix} \lambda^j - m! = \lambda^{\overline{m-1}} - m!, \tag{A.6}$$

the last equality following from [43, 1.2.6-(44)]. This polynomial is discussed in some detail in [49, Chapter 3] and [51]; some useful identities involving it are stated in Appendix B. From the discussion in [49] it follows that there are $m-1$ distinct (and nonzero) roots of ψ, call them $\lambda_1, \ldots, \lambda_{m-1}$ arranged in nonincreasing order of real parts. Thus the functions $\exp(\lambda_j x)$ are $m-1$ linearly independent solutions of (A.5) and hence the functions $(1-z)^{-\lambda_j}$ form a basis of solutions to $\mathbf{L}g = 0$.

A particular solution

We shall find it convenient to continue working in the x-domain of (A.5). There the differential equation equivalent to $\mathbf{L}g = h$ is $\widetilde{\mathbf{L}}\tilde{g} = \tilde{h}$ where

$$\tilde{h}(x) := e^{-(m-1)x} h(1-e^{-x}), \quad \text{i.e.,} \quad h(z) = (1-z)^{-(m-1)} \tilde{h}(L(z)). \tag{A.7}$$

From the elementary theory of linear differential equations (see, e.g., [6, §5.5]), a particular solution $\tilde{g}_p(x)$ to $\widetilde{\mathbf{L}}\tilde{g} = \tilde{h}$ in an interval near $x = 0$ is given by

$$\tilde{g}_p(x) = \sum_{j=1}^{m-1} y_j(x) \int_{t=0}^{x} \frac{\tilde{h}(t)\widetilde{W}_j(t)}{\widetilde{W}(t)}\, dt,$$

where $y_1, \ldots, y_{m-1}$ form any basis of solutions to $\widetilde{\mathbf{L}}\tilde{g} = 0$, $\widetilde{W}(x)$ is the Wronskian

$$\det\left(\left(y_j^{(i-1)}(x)\right)_{i,j=1,\ldots,m-1}\right),$$

and $\widetilde{W}_j(x)$ is the determinant obtained from $\widetilde{W}(x)$ by replacing the jth column by the column $\mathbf{e}_{m-1} := (0, 0, \ldots, 0, 1)^T$. By Cramer's rule, the column vector $(\widetilde{W}_j(x)/\widetilde{W}(x))_{j=1,\ldots,m-1}$ is the unique solution $\mathbf{a}$ to $\mathcal{W}(x)\mathbf{a} = \mathbf{e}_{m-1}$, i.e.,

$$\mathbf{a} = \mathcal{W}(x)^{-1}\mathbf{e}_{m-1} \tag{A.8}$$

where $\mathcal{W}(x) := \left(y_j^{(i-1)}(x)\right)_{i,j=1,\ldots,m-1}$. The general theory also shows that the initial conditions for $\tilde{g}_p$ all vanish, i.e.,

$$\tilde{g}_p^{(k)}(0) = 0, \quad 0 \le k \le m-2. \tag{A.9}$$

Here $y_j(x) = \exp(\lambda_j x)$, so that $y_j^{(i-1)}(x) = \lambda_j^{i-1}\exp(\lambda_j x)$ for $i, j = 1, \ldots, m-1$. It follows that

$$\mathcal{W}(x)^{-1} = (v^{ij}e^{-\lambda_i x})_{i,j=1,\ldots.m-1},$$

where

$$V := (v_{ij}) := (\lambda_j^{i-1})_{i,j=1,\ldots.m-1} \tag{A.10}$$

is the Vandermonde matrix and $V^{-1} =: (v^{ij})_{i,j=1,\ldots,m-1}$ is its inverse. The inverse is well known. By Exercise 1.2.3-40 in [43] (but note slight differences in definitions),

$$\begin{aligned}
v^{ij} &= [\lambda^{j-1}]\frac{(\lambda_1 - \lambda)\ldots(\lambda_{m-1} - \lambda)}{\lambda_i - \lambda}\Bigg/\prod_{1\le k\le m-1:\, k\ne i}(\lambda_k - \lambda_i)\\
&= [\lambda^{j-1}]\frac{\psi(\lambda)}{\lambda - \lambda_i}\frac{1}{\psi'(\lambda_i)} \qquad \text{[see (B.2) and (B.4)]}\\
&= -\frac{1}{\lambda_i\psi'(\lambda_i)}[\lambda^{j-1}]\left(\psi(\lambda)\left(1 - \frac{\lambda}{\lambda_i}\right)^{-1}\right)\\
&= -\frac{1}{\lambda_i\psi'(\lambda_i)}[\lambda^{j-1}]\left(-m! + \sum_{k=1}^{m}\begin{bmatrix} m-1\\ k\end{bmatrix}\lambda^k\right)\left(\sum_{l=0}^{\infty}\lambda_i^{-l}\lambda^l\right)\\
&= -\frac{1}{\lambda_i\psi'(\lambda_i)}\left(-m!\lambda_i^{-(j-1)} + \sum_{k=1}^{j-1}\begin{bmatrix} m-1\\ k\end{bmatrix}\lambda_i^{-(j-1-k)}\right)\\
&= -\frac{1}{\lambda_i^j\psi'(\lambda_i)}\left(\sum_{k=1}^{j-1}\begin{bmatrix} m-1\\ k\end{bmatrix}\lambda_i^k - m!\right) = \frac{1}{\lambda_i^j\psi'(\lambda_i)}\sum_{k=j}^{m-1}\begin{bmatrix} m-1\\ k\end{bmatrix}\lambda_i^k.
\end{aligned} \tag{A.11}$$

In particular, $\mathbf{a} = (a_i)_{i=1,\ldots,m-1}$ of (A.8) satisfies

$$a_i = v^{i,m-1}e^{-\lambda_i x} = \frac{1}{\lambda_i^{m-1}\psi'(\lambda_i)}\lambda_i^{m-1}e^{-\lambda_i x} = \frac{1}{\psi'(\lambda_i)}e^{-\lambda_i x}.$$

We have thus established

Lemma A.2. *The particular solution to $\widetilde{\mathbf{L}}\tilde{g} = \tilde{h}$ with vanishing initial conditions (through order*

$m-2$*) is*

$$\tilde{g}_p(x) = \sum_{j=1}^{m-1} \frac{e^{\lambda_j x}}{\psi'(\lambda_j)} \int_{t=0}^{x} \tilde{h}(t) e^{-\lambda_j t}\, dt.$$

If desired, this can be transformed back to the z-domain via (A.3), (A.4), and (A.7) and a change of variables in the integral:

Corollary A.3. *The particular solution to* $\mathbf{L}g = h$ *with vanishing initial conditions (through order* $m-2$*) is*

$$g_p(z) = \sum_{j=1}^{m-1} \frac{(1-z)^{-\lambda_j}}{\psi'(\lambda_j)} \int_0^z h(\zeta)(1-\zeta)^{\lambda_j+m-2}\, d\zeta.$$

The initial conditions

Having computed a basis of solutions to the homogeneous equation and a particular solution to the inhomogeneous equation, so far we have established Theorem 2.3.2 modulo determination of the coefficients $c_1, \ldots, c_{m-1}$ at (2.3.7).

The fact that the initial conditions for $\tilde{g}_p$ and for g_p vanish make it simple to solve $\mathbf{L}g = h$ or $\widetilde{\mathbf{L}}\tilde{g} = \tilde{h}$ for specified initial conditions: One need only match up the initial conditions of the homogeneous solutions.

In the x-domain the general complementary solution

$$\tilde{g}_{\mathrm{c}}(x) = \sum_{j=1}^{m-1} A_j e^{\lambda_j x}$$

has initial conditions

$$\tilde{g}_{\mathrm{c}}^{(k-1)}(0) = \sum_{j=1}^{m-1} A_j \lambda_j^{k-1} = \sum_{j=1}^{m-1} v_{kj} A_j, \quad k = 1, \ldots, m-1, \tag{A.12}$$

where V is the Vandermonde matrix at (A.10). In matrix notation, letting

$$\tilde{\mathbf{g}}_{\mathbf{0}} := (\tilde{g}_{\mathrm{c}}^{(0)}(0), \ldots, \tilde{g}_{\mathrm{c}}^{(m-2)}(0))^T$$

and $\mathbf{A} = (A_1, \ldots, A_{m-1})^T$, we can write (A.12) as $\tilde{\mathbf{g}}_{\mathbf{0}} = V\mathbf{A}$. Thus to obtain a specified $\tilde{\mathbf{g}}_{\mathbf{0}}$ we must choose $\mathbf{A} = V^{-1}\tilde{\mathbf{g}}_{\mathbf{0}}$. Recalling (A.11),

$$A_j = \frac{1}{\psi'(\lambda_j)} \sum_{l=1}^{m-1} \left(\sum_{i=l}^{m-1} \begin{bmatrix} m-1 \\ i \end{bmatrix} \lambda_j^{i-l} \right) \tilde{g}_{\mathrm{c}}^{(l-1)}(0), \quad j = 1, \ldots, m-1. \tag{A.13}$$

The general complementary solution to $\mathbf{L}g = h$ is then given by

$$g_{\mathrm{c}}(z) = \sum_{j=1}^{m-1} A_j (1-z)^{-\lambda_j}. \tag{A.14}$$

Finally we turn our attention back to g. According to Lemma A.1, $\mathbf{g}_{\mathbf{0}} = S\tilde{\mathbf{g}}_{\mathbf{0}}$, where

$$\mathbf{g}_{\mathbf{0}} := (g_{\mathrm{c}}^{(0)}(0), \ldots, g_{\mathrm{c}}^{(m-2)}(0))^T$$

are the initial conditions for g and S is the matrix

$$S := (s_{kj}) = \left(\begin{bmatrix} k-1 \\ j-1 \end{bmatrix} \right)_{k,j=1,\dots,m-1}$$

of (signless) Stirling numbers of the first kind. As is well known [43, §1.2.6],

$$S^{-1} := (s^{kj}) = \left((-1)^{j+k} \begin{Bmatrix} k-1 \\ j-1 \end{Bmatrix} \right)_{k,j=1,\dots,m-1}, \tag{A.15}$$

where $\left\{ {\cdot \atop \cdot} \right\}$ denotes (signless) Stirling numbers of the second kind. Thus in order for the general complementary solution (A.14) to have a specified vector $\mathbf{g_0}$ of initial conditions, we must take $\mathbf{A} = V^{-1}\tilde{\mathbf{g}}_{\mathbf{0}} = V^{-1}S^{-1}\mathbf{g_0}$. We can now establish (2.3.7). Observe that if $g_{\mathrm{c}}(z)$ is specified as a power series

$$g_{\mathrm{c}}(z) = \sum_{k=0}^{\infty} g_k z^k$$

then $g_{\mathrm{c}}^{(k)}(0) = k! g_k$ for $k = 0, \dots, m-2$.

Proposition A.4. *The constants A_j at* (A.13) *are given by*

$$A_j = \frac{m!}{\psi'(\lambda_j)} \sum_{k=0}^{m-2} \frac{g_c^{(k)}(0)}{\lambda_j^{\overline{k+1}}}, \quad j = 1, \dots, m-1.$$

Proof. Observe that, by (A.11) and (A.15),

$$\begin{aligned} A_j &= \sum_{k=1}^{m-1} \sum_{l=1}^{m-1} v^{jl} s^{lk} g_{\mathrm{c}}^{(k-1)}(0) = \sum_{k=1}^{m-1} \sum_{l=1}^{m-1} \frac{1}{\psi'(\lambda_j)} \sum_{i=l}^{m-1} \begin{bmatrix} m-1 \\ i \end{bmatrix} \lambda_j^{i-l} (-1)^{l+k} \begin{Bmatrix} l-1 \\ k-1 \end{Bmatrix} g_{\mathrm{c}}^{(k-1)}(0) \\ &= \frac{1}{\psi'(\lambda_j)} \sum_{l=1}^{m-1} \sum_{i=l}^{m-1} \begin{bmatrix} m-1 \\ i \end{bmatrix} \lambda_j^{i-l} \sum_{k=0}^{m-2} (-1)^{l+k+1} \begin{Bmatrix} l-1 \\ k \end{Bmatrix} g_{\mathrm{c}}^{(k)}(0). \end{aligned}$$

Hence it is enough to establish the curious identity

$$\sum_{l=1}^{m-1} \sum_{i=l}^{m-1} \begin{bmatrix} m-1 \\ i \end{bmatrix} \lambda_j^{i-l} (-1)^{l+k+1} \begin{Bmatrix} l-1 \\ k \end{Bmatrix} = \frac{m!}{\lambda_j^{\overline{k+1}}} \tag{A.16}$$

for $k = 0, \dots, m-2$.

For this, denote the left-hand side of (A.16) by f_k (fixing m and j). First, since $\left\{ {n \atop 0} \right\} \equiv \delta_{n,0}$, we have

$$f_0 = \sum_{i=1}^{m-1} \begin{bmatrix} m-1 \\ i \end{bmatrix} \lambda_j^{i-1} = \frac{\lambda_j^{\overline{m-1}}}{\lambda_j} = \frac{m!}{\lambda_j} \tag{A.17}$$

as claimed. For $k \geq 1$, using the recurrence relation [43, §1.2.6]

$$\begin{Bmatrix} n \\ k \end{Bmatrix} = k \begin{Bmatrix} n-1 \\ k \end{Bmatrix} + \begin{Bmatrix} n-1 \\ k-1 \end{Bmatrix}, \quad k \geq 1$$

for Stirling numbers of the second kind and the connection [43, 1.2.6-(47)] between the two kinds of Stirling numbers we see that

$$f_k = -\frac{k}{\lambda_j} f_k + \frac{1}{\lambda_j} f_{k-1} + \delta_{m-1,k}, \quad k \geq 1. \tag{A.18}$$

With initial condition (A.17) the recurrence (A.18) can be solved to give $f_k = m!/\lambda_j^{\overline{k+1}}$ for $k = 0, \ldots, m-2$, as desired. □

APPENDIX B

Identities involving the indicial polynomial

The indicial polynomial

$$\psi(\lambda) \equiv \psi_m(\lambda) := \lambda^{\overline{m-1}} - m! \tag{B.1}$$

plays an important role in the analysis of m-ary search trees under the random permutation model. We will enumerate a few useful identities involving the polynomial in this appendix.

It is well known [49, Chapter 3] that ψ_m has $m-1$ distinct roots $2 = \lambda_1, \lambda_2, \ldots, \lambda_{m-1}$ listed in nonincreasing order of real part. As in [49, Chapter 3] we introduce

$$\alpha := \max_{2 \le j \le m-1} \Re(\lambda_j);$$

that is, α is the real part of the root having the second largest real part among all the roots of the indicial polynomial (B.1). We list some important properties of the roots of (B.1) stated in [49, §3.3]:

1. The number $-m$ is a root if and only if m is odd. All other roots of $\psi(\lambda)$ are simple, non-real roots.
2. No two roots have the same real part unless they are mutually conjugate. (This follows from the strict increasingness of $|(s+it)^{\overline{m-1}}|$ in $|t|$.)

The indicial polynomial can be expressed as

$$\psi(\lambda) = \prod_{j=1}^{m-1} (\lambda - \lambda_j), \tag{B.2}$$

so that

$$\psi'(\lambda) = \sum_{j=1}^{m-1} \prod_{1 \le k \le m-1:\, k \ne j} (\lambda - \lambda_k). \tag{B.3}$$

If $\lambda = \lambda_j$ for some $j = 1, \ldots, m-1$, then from (B.3) we get

$$\psi'(\lambda_j) = \prod_{1 \le k \le m-1:\, k \ne j} (\lambda_j - \lambda_k) \tag{B.4}$$

since all other terms in the sum for $\psi'(\lambda_j)$ contain a factor that is 0.

Identity B.1. When $\lambda \notin \{\lambda_1, \ldots, \lambda_{m-1}\}$,

$$\sum_{j=1}^{m-1} \frac{1}{(\lambda - \lambda_j)\psi'(\lambda_j)} = \frac{1}{\psi(\lambda)}$$

Proof. Using (B.2) and partial fraction expansion we get, for $\lambda \notin \{\lambda_1, \ldots, \lambda_{m-1}\}$,

$$\frac{1}{\psi(\lambda)} = \frac{1}{\prod_{j=1}^{m}(\lambda - \lambda_j)} = \sum_{j=1}^{m-1} \frac{A_j}{\lambda - \lambda_j}$$

where, using (B.4),

$$A_j = \frac{1}{\prod_{1\le k\le m-1:\, k\neq j}(\lambda_j-\lambda_k)} = \frac{1}{\psi'(\lambda_j)},$$

as required. $\square$

For r and n positive integers, let $H_n^{(r)}$ denote the rth-order harmonic number

$$H_n^{(r)} := \sum_{j=1}^{n} \frac{1}{j^r}.$$

When $r=1$ we will use $H_n := H_n^{(1)}$ for the usual (1st-order) harmonic number.

Identity B.2.

$$\sum_{j=1}^{m-1} \frac{1}{(\lambda_j+r)^2\psi'(\lambda_j)} = \begin{cases} \dfrac{(m-1)!H_{m-1}}{[(m-1)!]^2(m-1)^2} = \dfrac{H_{m-1}}{(m-1)!(m-1)^2} & r=-1 \\[2ex] \dfrac{(-1)^r r!(m-2-r)!}{[(-r)^{\overline{m-1}}-m!]^2} & 0\le r\le m-2 \\[2ex] \dfrac{(-1)^m r^{\underline{m-1}}(H_r-H_{r-(m-1)})}{[(-1)^{m-1}r^{\underline{m-1}}-m!]^2} & r\ge m-1. \end{cases}$$

Proof. Differentiate Identity B.1 with respect to λ to get

$$\sum_{j=1}^{m-1} \frac{1}{(\lambda-\lambda_j)^2\psi'(\lambda_j)} = \frac{\psi'(\lambda)}{\psi^2(\lambda)}, \qquad \lambda\notin\{\lambda_1,\dots,\lambda_{m-1}\}. \tag{B.5}$$

Differentiating the defining equation (B.1) we get

$$\psi'(\lambda) = \sum_{j=0}^{m-2}\ \prod_{0\le k\le m-2:\, k\neq j}(\lambda+k) = \lambda^{\overline{m-1}}\sum_{j=0}^{m-2}\frac{1}{\lambda+j}, \tag{B.6}$$

the last equality being valid only for $\lambda\notin\{0,-1,\dots,-(m-2)\}$. Hence for $r\ge -1$,

$$\sum_{j=1}^{m-1}\frac{1}{(\lambda_j+r)^2\psi'(\lambda_j)} = \frac{\psi'(-r)}{\psi^2(-r)} = \frac{\psi'(-r)}{((-r)^{\overline{m-1}}-m!)^2}.$$

Now using (B.6) we see that

$$\psi'(1) = 1^{\overline{m-1}}\sum_{j=0}^{m-2}\frac{1}{1+j} = (m-1)!H_{m-1},$$

$$\psi'(-r) = \prod_{0\le j\le m-2:\, j\neq r}(j-r) = (-1)^r r!(m-2-r)!, \qquad 0\le r\le m-2,$$

$$\psi'(-r) = (-r)^{\overline{m-1}}\sum_{j=0}^{m-2}\frac{1}{j-r} = (-1)^m r^{\underline{m-1}}(H_r-H_{r-(m-1)}), \qquad r\ge m-1,$$

and the claim follows. $\square$

Identity B.3. For $0 \leq k \leq m-3$,

$$\sum_{j=1}^{m-1} \frac{\lambda_j^k}{\psi'(\lambda_j)} = 0.$$

Proof. We take asymptotic expansions of both sides of Identity B.1 as $\lambda \to \infty$ in powers of $1/\lambda$ and equate coefficients. The left-hand side yields

$$\frac{1}{\lambda}\sum_{j=1}^{m-1} \frac{1}{\left(1-\frac{\lambda_j}{\lambda}\right)\psi'(\lambda_j)} = \frac{1}{\lambda}\sum_{j=1}^{m-1} \frac{1}{\psi'(\lambda_j)} \sum_{k=0}^{\infty} \frac{\lambda_j^k}{\lambda^k} = \sum_{k=0}^{\infty} \frac{1}{\lambda^{k+1}} \sum_{j=1}^{m-1} \frac{\lambda_j^k}{\psi'(\lambda_j)}.$$

On the other hand, the right-hand side equals

$$\frac{1}{\lambda^{\overline{m-1}} - m!} = \left(\prod_{k=0}^{m-2} (\lambda+k) - m!\right)^{-1} = \frac{1}{\lambda^{m-1}} \left(\prod_{k=0}^{m-2}\left(1+\frac{k}{\lambda}\right) - \frac{m!}{\lambda^{m-1}}\right)^{-1}.$$

Hence the coefficient of $1/\lambda^{k+1}$ for $k = 0, \ldots, m-3$ in the left-hand side must vanish, proving the claim. ☐

Identity B.4.

$$\sum_{j=2}^{m-1} \frac{1}{(\lambda_j - 2)\psi'(\lambda_j)} = \frac{1}{2(m!)}\left[1 - \frac{H_m^{(2)} - 1}{(H_m - 1)^2}\right].$$

Proof. When λ_k is a root of ψ we have from (B.6)

$$\psi'(\lambda_k) = m! \sum_{j=0}^{m-2} \frac{1}{\lambda_k + j}.$$

Also, differentiating (B.6) once more gives us

$$\psi''(\lambda) = \lambda^{\overline{m-1}} \left[\left(\sum_{j=0}^{m-2} \frac{1}{\lambda+j}\right)^2 - \sum_{j=0}^{m-2} \frac{1}{(\lambda+j)^2}\right], \tag{B.7}$$

so that

$$\psi'(2) = m!(H_m - 1) \qquad \text{and} \qquad \psi''(2) = m![(H_m-1)^2 - (H_m^{(2)} - 1)]. \tag{B.8}$$

Thus, as $\lambda \to 2$,

$$\psi(\lambda) = (\lambda-2)m!(H_m - 1) + \frac{1}{2}m![(H_m-1)^2 - (H_m^{(2)}-1)](\lambda-2)^2 + O(|\lambda-2|^3).$$

By Identity B.1, for $\lambda \notin \{\lambda_1, \ldots, \lambda_{m-1}\}$ we have

$$\begin{aligned}
&\sum_{j=2}^{m-1} \frac{1}{(\lambda-\lambda_j)\psi'(\lambda_j)} \\
&= \frac{1}{\psi(\lambda)} - \frac{1}{(\lambda-2)m!(H_m-1)} \\
&= \frac{1}{(\lambda-2)m!(H_m-1)} \left\{ \left[1 + \frac{1}{2}\left[(H_m-1) - \frac{H_m^{(2)}-1}{H_m-1} \right](\lambda-2) + O(|\lambda-2|^2) \right]^{-1} - 1 \right\} \\
&= \frac{1}{(\lambda-2)m!(H_m-1)} \left\{ -\frac{1}{2}\left[(H_m-1) - \frac{H_m^{(2)}-1}{H_m-1} \right](\lambda-2) + O(|\lambda-2|^2) \right\} \\
&= -\frac{1}{2(m!)}\left[1 - \frac{H_m^{(2)}-1}{(H_m-1)^2} \right] + O(|\lambda-2|);
\end{aligned}$$

in particular, letting $\lambda \to 2$ yields

$$\sum_{j=2}^{m-1} \frac{1}{(\lambda_j-2)\psi'(\lambda_j)} = \frac{1}{2(m!)}\left[1 - \frac{H_m^{(2)}-1}{(H_m-1)^2} \right],$$

as claimed. □

For the next two identities we assume that m is odd. Note that when m is odd, $-m$ is a root of ψ_m—in fact, $\lambda_{m-1} = -m$, as noted at the beginning of this appendix.

Identity B.5. When m is odd,

$$\sum_{j=1}^{m-2} \frac{1}{(\lambda_j+m)\psi'(\lambda_j)} = \frac{1}{2(m!)}\left[1 - \frac{H_m^{(2)}-1}{(H_m-1)^2} \right].$$

Proof. By Identity B.1 the left-hand side above equals

$$\begin{aligned}
&-\lim_{\lambda\to -m}\left[\frac{1}{\psi(\lambda)} - \frac{1}{(\lambda+m)\psi'(-m)} \right] \\
&= -\lim_{\lambda\to -m}\left[\frac{1}{(\lambda+m)\psi'(-m) + \frac{1}{2}(\lambda+m)^2\psi''(-m) + O(|\lambda+m|^3)} - \frac{1}{(\lambda+m)\psi'(-m)} \right] \\
&= \lim_{\lambda\to -m} \frac{1}{(\lambda+m)\psi'(-m)}\left[\frac{1}{2}(\lambda+m)\frac{\psi''(-m)}{\psi'(-m)} + O(|\lambda+m|^2) \right] = \frac{1}{2}\frac{\psi''(-m)}{[\psi'(-m)]^2}.
\end{aligned}$$

Now by (B.6),

$$\psi'(-m) = -m!(H_m-1), \tag{B.9}$$

and by (B.7)

$$\psi''(-m) = m!\left[\left(\sum_{j=0}^{m-2} \frac{1}{j-m} \right)^2 - \sum_{j=0}^{m-2} \frac{1}{(j-m)^2} \right] = m![(H_m-1)^2 - (H_m^{(2)}-1)], \tag{B.10}$$

and the result follows. □

Identity B.6. When m is odd

$$\sum_{j=1}^{m-2} \frac{1}{(\lambda_j+m)^2\psi'(\lambda_j)} = \frac{1}{m!}\left[\frac{1}{12}(H_m-1) - \frac{1}{3}\frac{H_m^{(3)}-1}{(H_m-1)^2} + \frac{1}{4}\frac{(H_m^{(2)}-1)^2}{(H_m-1)^3}\right].$$

Proof. By (B.5) the left-hand-side above equals

$$\begin{aligned}
&\lim_{\lambda\to-m}\left[\frac{\psi'(\lambda)}{\psi^2(\lambda)} - \frac{1}{(\lambda+m)^2\psi'(-m)}\right] \\
&= \lim_{\lambda\to-m}\Bigg[\frac{\psi'(-m)+\psi''(-m)(\lambda+m)+\frac{1}{2}\psi'''(-m)(\lambda+m)^2+O(|\lambda+m|^3)}{[\psi'(-m)(\lambda+m)+\frac{1}{2}\psi''(-m)(\lambda+m)^2+\frac{1}{6}\psi'''(-m)(\lambda+m)^3+O(|\lambda+m)^4|]^2} \\
&\qquad - \frac{1}{(\lambda+m)^2\psi'(-m)}\Bigg] \\
&= \lim_{\lambda\to-m}\Bigg[\frac{1}{(\lambda+m)^2\psi'(-m)} \\
&\qquad \times\left(\frac{1+\frac{\psi''(-m)}{\psi'(-m)}(\lambda+m)+\frac{1}{2}\frac{\psi'''(-m)}{\psi'(-m)}(\lambda+m)^2+O(|\lambda+m|^3)}{[1+\frac{1}{2}\frac{\psi''(-m)}{\psi'(-m)}(\lambda+m)+\frac{1}{6}\frac{\psi'''(-m)}{\psi'(-m)}(\lambda+m)^2+O(|\lambda+m|^3)]^2} - 1\right)\Bigg] \\
&= \lim_{\lambda\to-m}\Bigg[\frac{1}{(\lambda+m)^2\psi'(-m)} \\
&\qquad \times\left(\frac{1+\frac{\psi''(-m)}{\psi'(-m)}(\lambda+m)+\frac{1}{2}\frac{\psi'''(-m)}{\psi'(-m)}(\lambda+m)^2+O(|\lambda+m|^3)}{1+\frac{\psi''(-m)}{\psi'(-m)}(\lambda+m)+\big[\frac{1}{4}(\frac{\psi''(-m)}{\psi'(-m)})^2+\frac{1}{3}\frac{\psi'''(-m)}{\psi'(-m)}\big](\lambda+m)^2+O(|\lambda+m|^3)} - 1\right)\Bigg] \\
&= \lim_{\lambda\to-m}\Bigg[\frac{1}{(\lambda+m)^2\psi'(-m)} \\
&\times\Bigg(\Big[1+\tfrac{\psi''(-m)}{\psi'(-m)}(\lambda+m)+\tfrac{1}{2}\tfrac{\psi'''(-m)}{\psi'(-m)}(\lambda+m)^2+O(|\lambda+m|^3)\Big] \\
&\times\left[1-\frac{\psi''(-m)}{\psi'(-m)}(\lambda+m)+\left[\frac{3}{4}\left(\frac{\psi''(-m)}{\psi'(-m)}\right)^2-\frac{1}{3}\frac{\psi'''(-m)}{\psi'(-m)}\right](\lambda+m)^2+O(|\lambda+m|^3)\right]-1\Bigg)\Bigg] \\
&= \lim_{\lambda\to-m}\left[\frac{1}{(\lambda+m)^2\psi'(-m)}\left(\left(\frac{1}{6}\frac{\psi'''(-m)}{\psi'(-m)}-\frac{1}{4}\left(\frac{\psi''(-m)}{\psi'(-m)}\right)^2\right)(\lambda+m)^2+O(|\lambda+m|^3)\right)\right] \\
&= \frac{1}{6}\frac{\psi'''(-m)}{[\psi'(-m)]^2} - \frac{1}{4}\frac{[\psi''(-m)]^2}{[\psi'(-m)]^3}.
\end{aligned}$$

We have already computed $\psi'(-m)$ and $\psi''(-m)$ at (B.9) and (B.10), respectively. To calculate

$\psi'''(-m)$ we differentiate (B.7) to get, for $\lambda \notin \{0, -1, \ldots, -(m-2)\}$,

$$\begin{aligned}
\psi'''(\lambda) &= \lambda^{\overline{m-1}}\Bigg[2\sum_{j=0}^{m-2}\frac{1}{(\lambda+j)^3} - 2\left(\sum_{j=0}^{m-2}\frac{1}{\lambda+j}\right)\left(\sum_{j=0}^{m-2}\frac{1}{(\lambda+j)^2}\right) \\
&\qquad - \left(\sum_{j=0}^{m-2}\frac{1}{\lambda+j}\right)\left(\sum_{j=0}^{m-2}\frac{1}{(\lambda+j)^2}\right) + \left(\sum_{j=0}^{m-2}\frac{1}{\lambda+j}\right)^3\Bigg] \\
&= \lambda^{\overline{m-1}}\left[2\sum_{j=0}^{m-2}\frac{1}{(\lambda+j)^3} - 3\left(\sum_{j=0}^{m-2}\frac{1}{\lambda+j}\right)\left(\sum_{j=0}^{m-2}\frac{1}{(\lambda+j)^2}\right) + \left(\sum_{j=0}^{m-2}\frac{1}{\lambda+j}\right)^3\right].
\end{aligned}$$

In particular, since $-m$ is a root of ψ when m is odd,

$$\psi'''(-m) = m![-2(H_m^{(3)}-1) + 3(H_m-1)(H_m^{(2)}-1) - (H_m-1)^3].$$

Therefore

$$\begin{aligned}
\sum_{j=1}^{m-2}\frac{1}{(\lambda_j+m)^2\psi'(\lambda_j)} &= \frac{1}{6}\frac{m![-2(H_m^{(3)}-1)+3(H_m-1)(H_m^{(2)}-1)-(H_m-1)^3]}{(m!)^2(H_m-1)^2} \\
&\qquad + \frac{1}{4}\frac{[(H_m-1)^2-(H_m^{(2)}-1)]^2}{m!(H_m-1)^3} \\
&= \frac{1}{6(m!)}\left[-(H_m-1) + \frac{3(H_m^{(2)}-1)}{H_m-1} - \frac{2(H_m^{(3)}-1)}{(H_m-1)^2}\right] \\
&\qquad + \frac{1}{4(m!)}\left[(H_m-1) - \frac{2(H_m^{(2)}-1)}{H_m-1} + \frac{(H_m^{(2)}-1)^2}{(H_m-1)^3}\right] \\
&= \frac{1}{m!}\left[\frac{1}{12}(H_m-1) - \frac{1}{3}\frac{H_m^{(3)}-1}{(H_m-1)^2} + \frac{1}{4}\frac{(H_m^{(2)}-1)^2}{(H_m-1)^3}\right],
\end{aligned}$$

as claimed. $\square$

Bibliography

[1] M. Abramowitz and I. A. Stegun. *Handbook of mathematical functions with formulas, graphs, and mathematical tables*, volume 55 of *National Bureau of Standards Applied Mathematics Series*. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C., 1964.

[2] L. Alonso, P. Chassaing, F. Gillet, S. Janson, E. M. Reingold, and R. Schott. Sorting with unreliable comparisons: a probabilistic analysis. Preprint. Available at `http://www.math.uu.se/~svante/papers/`, 2003.

[3] P. Billingsley. *Probability and measure*. John Wiley & Sons Inc., New York, third edition, 1995. A Wiley-Interscience Publication.

[4] N. H. Bingham, C. M. Goldie, and J. L. Teugels. *Regular variation*, volume 27 of *Encyclopedia of Mathematics and its Applications*. Cambridge University Press, Cambridge, 1989.

[5] N. Bleistein and R. A. Handelsman. *Asymptotic expansions of integrals*. Dover Publications Inc., New York, second edition, 1986.

[6] W. E. Boyce and R. C. DiPrima. *Elementary Differential Equations*. John Wiley & Sons, 4th edition, 1986.

[7] B. Chauvin, M. Drmota, and J. Jabbour-Hattab. The profile of binary search trees. *Ann. Appl. Probab.*, 11(4):1042–1062, 2001.

[8] H.-H. Chern and H.-K. Hwang. Phase changes in random m-ary search trees and generalized quicksort. *Random Structures Algorithms*, 19(3-4):316–358, 2001. Analysis of algorithms (Krynica Morska, 2000).

[9] K. L. Chung. *A course in probability theory*. Academic Press [A subsidiary of Harcourt Brace Jovanovich, Publishers], New York-London, second edition, 1974. Probability and Mathematical Statistics, Vol. 21.

[10] M. Cramer. A note concerning the limit distribution of the quicksort algorithm. *RAIRO Inform. Théor. Appl.*, 30(3):195–207, 1996.

[11] R. De Prisco and A. De Santis. New lower bounds on the cost of binary search trees. *Theoret. Comput. Sci.*, 156(1-2):315–325, 1996.

[12] L. Devroye. A note on the height of binary search trees. *J. Assoc. Comput. Mach.*, 33(3):489–498, 1986.

[13] L. Devroye. On the richness of the collection of subtrees in random binary search trees. *Inform. Process. Lett.*, 65(4):195–199, 1998.

[14] L. Devroye and B. Reed. On the variance of the height of random binary search trees. *SIAM J. Comput.*, 24(6):1157–1162, 1995.

[15] R. P. Dobrow and J. A. Fill. On the Markov chain for the move-to-root rule for binary search trees. *Ann. Appl. Probab.*, 5(1):1–19, 1995.

[16] R. P. Dobrow and J. A. Fill. Rates of convergence for the move-to-root Markov chain for binary search trees. *Ann. Appl. Probab.*, 5(1):20–36, 1995.

[17] R. P. Dobrow and J. A. Fill. Multiway trees of maximum and minimum probability under the random permutation model. *Combin. Probab. Comput.*, 5(4):351–371, 1996.

[18] M. Drmota. An analytic approach to the height of binary search trees. *Algorithmica*, 29(1-2):89–119, 2001. Average-case analysis of algorithms (Princeton, NJ, 1998).

[19] M. Drmota. The variance of the height of binary search trees. *Theoret. Comput. Sci.*, 270(1-2):913–919, 2002.

[20] W. Feller. *An introduction to probability theory and its applications. Vol. II.* Second edition. John Wiley & Sons Inc., New York, 1971.

[21] J. A. Fill. On the distribution of binary search trees under the random permutation model. *Random Structures Algorithms*, 8(1):1–25, 1996.

[22] J. A. Fill and R. P. Dobrow. The number of m-ary search trees on n keys. *Combin. Probab. Comput.*, 6(4):435–453, 1997.

[23] J. A. Fill, P. Flajolet, and N. Kapur. Singularity analysis, Hadamard products, and tree recurrences, 2003. In preparation.

[24] J. A. Fill and S. Janson. A characterization of the set of fixed points of the Quicksort transformation. *Electron. Comm. Probab.*, 5:77–84 (electronic), 2000.

[25] J. A. Fill and S. Janson. Smoothness and decay properties of the limiting Quicksort density function. In *Mathematics and computer science (Versailles, 2000)*, Trends Math., pages 53–64. Birkhäuser, Basel, 2000.

[26] J. A. Fill and S. Janson. Approximating the limiting Quicksort distribution. *Random Structures Algorithms*, 19(3-4):376–406, 2001, arXiv:math.PR/0105246. Analysis of algorithms (Krynica Morska, 2000).

[27] J. A. Fill and S. Janson. Quicksort asymptotics. *J. Algorithms*, 44(1):4–28, 2002, arXiv:math.PR/0105248. Analysis of algorithms.

[28] P. Flajolet. Singularity analysis and asymptotics of Bernoulli sums. *Theoret. Comput. Sci.*, 215(1-2):371–381, 1999.

[29] P. Flajolet, X. Gourdon, and C. Martínez. Patterns in random binary search trees. *Random Structures Algorithms*, 11(3):223–244, 1997.

[30] P. Flajolet and G. Louchard. Analytic variations on the Airy distribution. *Algorithmica*, 31(3):361–377, 2001. Mathematical analysis of algorithms.

[31] P. Flajolet and A. Odlyzko. Singularity analysis of generating functions. *SIAM J. Discrete Math.*, 3(2):216–240, 1990.

[32] P. Flajolet and J.-M. Steyaert. A complexity calculus for recursive tree algorithms. *Math. Systems Theory*, 19(4):301–331, 1987.

[33] M. Foley and C. A. R. Hoare. Proof of a recursive program: quicksort. *Comput. J.*, 14:391–395, 1971.

[34] N. Fröman and P. O. Fröman. *JWKB approximation. Contributions to the theory.* North-Holland Publishing Co., Amsterdam, 1965.

[35] P. Hennequin. Combinatorial analysis of quicksort algorithm. *RAIRO Inform. Théor. Appl.*, 23(3):317–333, 1989.

[36] P. Hennequin. *Analyse en moyenne d'algorithme, tri rapide et arbres de recherche.* PhD thesis, Ecole Polytechnique, 1991.

[37] P. Henrici. *Applied and computational complex analysis. Vol. 2.* Wiley Interscience [John Wiley & Sons], New York, 1977. Special functions—integral transforms—asymptotics—continued fractions.

[38] H.-K. Hwang and R. Neininger. Phase change of limit laws in the quicksort recurrence under varying toll functions. *SIAM J. Comput.*, 31(6):1687–1722, 2002.

[39] J. Jabbour-Hattab. Martingales and large deviations for binary search trees. *Random Structures Algorithms*, 19(2):112–127, 2001.

[40] S. Janson. Ideals in a forest, one-way infinite binary trees and the contraction method. In *Mathematics and computer science, II (Versailles, 2002)*, Trends Math., pages 393–414. Birkhäuser, Basel, 2002.

[41] S. Janson. The Wiener index of simply generated random trees. Preprint. Available at `http://www.math.uu.se/~svante/papers/`, 2002.

[42] C. Knessl and W. Szpankowski. Quicksort algorithm again revisited. *Discrete Math. Theor. Comput. Sci.*, 3(2):43–63 (electronic), 1999.

[43] D. E. Knuth. *The art of computer programming. Volume 1.* Addison-Wesley Publishing Co., Reading, Mass.-London-Don Mills, Ont., 3rd edition, 1997.

[44] D. E. Knuth. *The art of computer programming. Volume 3.* Addison-Wesley Publishing Co., Reading, Mass.-London-Don Mills, Ont., 2nd edition, 1998.

[45] D. E. Knuth and B. Pittel. A recurrence related to trees. *Proc. Amer. Math. Soc.*, 105(2):335–349, 1989.

[46] D. E. Knuth and A. Schönhage. The expected linearity of a simple equivalence algorithm. *Theoret. Comput. Sci.*, 6(3):281–315, 1978.

[47] Y. Koda and F. Ruskey. A Gray code for the ideals of a forest poset. *J. Algorithms*, 15(2):324–340, 1993.

[48] W. Lew and H. M. Mahmoud. The joint distribution of elastic buckets in multiway search trees. *SIAM J. Comput.*, 23(5):1050–1074, 1994.

[49] H. M. Mahmoud. *Evolution of random search trees.* John Wiley & Sons Inc., New York, 1992. A Wiley-Interscience Publication.

[50] H. M. Mahmoud and B. Pittel. Analysis of the space of search trees under the random insertion algorithm. *J. Algorithms*, 10(1):52–75, 1989.

[51] H. M. Mahmoud and R. T. Smythe. Probabilistic analysis of bucket recursive trees. *Theoret. Comput. Sci.*, 144(1-2):221–249, 1995. Special volume on mathematical analysis of algorithms.

[52] A. W. Marshall and I. Olkin. *Inequalities: theory of majorization and its applications.* Academic Press Inc. [Harcourt Brace Jovanovich Publishers], New York, 1979.

[53] C. Martínez and S. Roura. Randomized binary search trees. *J. ACM*, 45(2):288–323, 1998.

[54] A. Meir and J. W. Moon. On the altitude of nodes in random trees. *Canad. J. Math.*, 30(5):997–1015, 1978.

[55] A. Meir and J. W. Moon. On the log-product of the subtree-sizes of random trees. *Random Structures Algorithms*, 12(2):197–212, 1998.

[56] R. R. Muntz and R. C. Uzgalis. Dynamic storage allocation for binary search trees in a two-level memory. In *Proceedings of Princeton Conference on Information Sciences and Systems*, volume 4, pages 345–349, 1971.

[57] R. Neininger. On binary search tree recursions with monomials as toll functions. *J. Comput. Appl. Math.*, 142(1):185–196, 2002. Probabilistic methods in combinatorics and combinatorial optimization.

[58] A. Panholzer and H. Prodinger. An analytic approach for the analysis of rotations in fringe-balanced binary search trees. *Ann. Comb.*, 2(2):173–184, 1998.

[59] A. Panholzer and H. Prodinger. Binary search tree recursions with harmonic toll functions. *J. Comput. Appl. Math.*, 142(1):211–225, 2002. Probabilistic methods in combinatorics and combinatorial optimization.

[60] M. Régnier. A limiting distribution for quicksort. *RAIRO Inform. Théor. Appl.*, 23(3):335–343, 1989.

[61] J. M. Robson. On the concentration of the height of binary search trees. In *Automata, languages and programming (Bologna, 1997)*, volume 1256 of *Lecture Notes in Comput. Sci.*, pages 441–448. Springer, Berlin, 1997.

[62] J. M. Robson. Constant bounds on the moments of the height of binary search trees. *Theoret. Comput. Sci.*, 276(1-2):435–444, 2002.

[63] U. Rösler. A limit theorem for “Quicksort”. *RAIRO Inform. Théor. Appl.*, 25(1):85–100, 1991.

[64] U. Rösler. On the analysis of stochastic divide and conquer algorithms. *Algorithmica*, 29(1-2):238–261, 2001. Average-case analysis of algorithms (Princeton, NJ, 1998).

[65] U. Rösler and L. Rüschendorf. The contraction method for recursive algorithms. *Algorithmica*, 29(1-2):3–33, 2001. Average-case analysis of algorithms (Princeton, NJ, 1998).

[66] W. Schachinger. Asymptotic normality of recursive algorithms via martingale difference arrays. *Discrete Math. Theor. Comput. Sci.*, 4(2):363–397 (electronic), 2001.

[67] R. Sedgewick. Quicksort with equal keys. *SIAM J. Comput.*, 6(2):240–267, 1977.

[68] R. P. Stanley. *Enumerative combinatorics. Vol. 2.* Cambridge University Press, Cambridge, 1999. With a foreword by Gian-Carlo Rota and appendix 1 by Sergey Fomin.

[69] J.-M. Steyaert and P. Flajolet. Patterns and pattern-matching in trees: an analysis. *Inform. and Control*, 58(1-3):19–58, 1983.

[70] L. Takács. A Bernoulli excursion and its various applications. *Adv. in Appl. Probab.*, 23(3):557–585, 1991.

[71] L. Takács. Conditional limit theorems for branching processes. *J. Appl. Math. Stochastic Anal.*, 4(4):263–292, 1991.

[72] L. Takács. On a probability problem connected with railway traffic. *J. Appl. Math. Stochastic Anal.*, 4(1):1–27, 1991.

[73] K. H. Tan and P. Hadjicostas. Some properties of a limiting distribution in Quicksort. *Statist. Probab. Lett.*, 25(1):87–94, 1995.

Vita

Nevin Kapur was born on May 9, 1974 in Mumbai (formerly Bombay), India.

He received a Bachelor of Technology in Electrical Engineering, with a specialization in VLSI design and automation, from the Indian Institute of Technology, Bombay in 1996. From 1996 to 1998 he was part of the Collaborative Benchmarking Laboratory at North Carolina State University (NCSU). He received a Master of Science from the Electrical and Computer Engineering department, specializing in graph algorithms and their applications to automatic place-and-route in VLSI design. At NCSU, while taking courses in design and analysis of algorithms, graph theory, and probability, he became interested in these topics and their intersection. This motivated his joining the Department of Mathematical Sciences at The Johns Hopkins University in 1998.

Nevin plans on pursuing an academic career, as a researcher pursuing problems in discrete mathematics, probability, and their applications to computer science.

Baltimore, 2003